

1 **Manuscript title:**

2 Can headache profile predict future disability: a cohort study.

3

4 **Authors informations:**

5

6 **Mariève Houle, MSc (c)**

7 Department of Human Kinetics, Université du Québec à Trois-Rivières

8 3351 Boul. des Forges, Trois-Rivières, Qc G8Z 4M3

9 Phone : (1) 819-376-5011 ext.3798

10 Email: marieve.houle@uqtr.ca

11

12 **Andrée-Anne Marchand, PhD**

13 Postdoctoral fellow

14 Faculty of Health Sciences, Ontario Tech University

15 2000 Simcoe Street North, Oshawa, ON, L1G 0C5

16 Phone : None

17 andree-anne.marchand@uoit.ca

18

19

20 **Martin Descarreaux, PhD**

21 Department of Human Kinetics, Université du Québec à Trois-Rivières

22 3351 Boul. des Forges, Trois-Rivières, Qc G8Z 4M3

23 Phone : (1) 819-376-5011 ext.3798

24 Email : martin.descarreaux@uqtr.ca

25

26

27 **Conflict of Interest Statement:** There is no conflict of interest.

28

29 **Funding :** This research received no specific grant from any funding agency in the public,

30 commercial, or not-for-profit sectors.

31 **Corresponding author:** Mariève Houle

32

33

34

35 **Abstract:**

36 **Objectives:**

37 The aim of this study was to determine if headache profile can predict future disability in
38 patients with TTH.

39 **Methods**

40 Eighty-three patients with TTH were recruited. To be included in the study participants
41 needed to fulfill the International Headache Society classification's criteria for episodic or
42 chronic TTH form and to be at least 18 years old. Baseline clinical outcomes (headache
43 and neck-related disability, kinesiophobia, self-efficacy and anxiety) and physical
44 outcomes (neck extensors muscles maximum voluntary contraction) were collected for all
45 patients. A prospective data collection of headache characteristics (intensity and frequency)
46 was conducted using daily SMS or e-mail over a 1-month period. Headache-related
47 disability was assessed at the 3-month follow-up and was used as the disability criterion
48 for TTH.

49 **Results**

50 Correlations showed that the number of years with headache ($r=.53$; $p<0.001$, self-
51 reported neck pain intensity ($r=.29$; $p=0.025$), headache frequency ($r=.60$; $p<0.001$) and
52 intensity ($r=.54$; $p<0.001$), anxiety ($r=.28$; $p=0.031$) as well as neck-related disability
53 ($r=.63$; $p<0.001$) were correlated to headache-related disability assessed at 3 months.
54 Multiple regression showed that these determinants can be used to predict headache
55 disability ($R^2= 0.583$). Headache frequency ($\beta=0.284$) was the best individual predictor.

56

57

58 **Discussion**

59 Results showed that TTH frequency and intensity and the presence of concomitant
60 infrequent migraine are predictors of future disability over a 3-month period. Further
61 studies are needed to evaluate the contribution of other potential physical outcomes on
62 headache-related disability.

63 **Keys words:** Tension-type headache; headache profile; neck pain; disability; strength

64 **List of abbreviations**

65 TTH: tension-type headache

66 IETTH: infrequent episodic tension-type headache

67 FETTH: frequent episodic tension-type headache

68 CTTH: chronic tension-type headache

69 IHS: International Headache Society

70 MVC: maximum voluntary contraction

71 HIT-6: 6-item headache impact test

72 NDI: neck disability index

73 VAS: visual analogue scale

74

75

76

77

78

79

80

81 **Can headache profile predict future disability: a cohort study.**

82 **Introduction**

83 Tension-type headache (TTH) is the most common type of headache [1] with a lifetime
84 prevalence in the general population that ranges between 30% to 78% [2-4]. The average
85 age of TTH onset is estimated between 25 and 30 years old with the peak prevalence for
86 both sexes occurring between 30 to 39 years of age, followed by a decline with increasing
87 age [1, 5]. Women are more affected than men with female:male ratios ranging from 1.3:1
88 to 5:4 [1, 5, 6].

89 According to the International Headache Society (IHS), tension-type headache is classified
90 as a primary headache [3] and typically described by a bilateral, pressing, tightening and
91 non-pulsating pain. In addition, intensity of TTH is considered to be mild to moderate
92 without aggravation by routine physical activity such as climbing stairs or walking and is
93 not associated with nausea or vomiting except for mild nausea that can be present in chronic
94 TTH (CTTH) [5]. Furthermore, phonophobia or photophobia can occur during TTH
95 episodes, but both symptoms should not be present at the same time [3, 5]. TTH can be
96 divided into two categories: episodic or chronic form. The episodic form is further divided
97 into two subcategories based on the frequency of episodes: infrequent episodic tension-
98 type headache (IETTH) or frequent episodic tension-type headache (FETTH) [3]. IETTH
99 is characterized by at least 10 episodes occurring less than one day per month (< 12 days
100 per years), FETTH is characterized by 1 to 14 days per month for at least 3 consecutive
101 months (> 12 and < 180 days per year) and CTTH consists of fifteen days or more with
102 headaches per month (> 180 days per year) [1, 3]. In a Danish population-based study, the
103 one-year prevalence of each category at 40 years old was 48.2%, 33.8% and 2.3%

104 respectively and the prevalence was higher in men for the infrequent episodic TTH than in
105 women but frequent episodic and chronic TTH was more frequent in women than in men
106 [7]. Tension-type headache and particularly CTTH are often associated with medical and
107 psychiatric conditions. Indeed, TTH has previously been linked to common comorbidities
108 such as temporomandibular disorders, depression, anxiety and panic disorders [5].

109 To date, only a few studies have evaluated risk factors for the development of TTH or risk
110 factors that leads to a transition from episodic to chronic forms [8-10]. In fact,
111 environmental, genetic and peripheral factors such as tenderness in pericranial muscles,
112 muscle strain, muscle blood flow and other central factors have all been hypothesized as
113 possible contributors to the development of TTH [8]. Moreover, risk factors to transition
114 from the episodic TTH form to the chronic TTH form are divided into two categories: non-
115 modifiable risk factors and modifiable risk factors. On the one hand, non-modifiable risk
116 factors include increasing age, female sex, being Caucasian, previous history of head
117 trauma and low socioeconomic status [9]. On the other hand, modifiable risk factors
118 include sleep disturbances, medication overuse, obesity and the presence of psychological
119 comorbidities [9]. Depression and anxiety are common with TTH, especially in the chronic
120 form [11]. However, there are only few studies with follow-ups that have been conducted
121 in adults with tension-type headache. A study by Lyngberg and colleagues showed that, in
122 adults randomly drawn from the general population, predictors of poor outcomes in patients
123 with tension-type headache were having frequent episodic or chronic tension-type
124 headache at baseline, having coexisting migraine, not being married and having sleep
125 problems. In their study, no association was found with respect to age, gender or
126 educational levels [12].

127 Frequency and intensity of tension-type headache are two factors frequently assessed in
128 studies but are also used to define inclusion criteria for TTH participants (IETTH, FETTH
129 or CTTH group). Moreover, Sauro et al. reported a positive correlation between intensity
130 and headache-related disability as evaluated by the 6-item headache impact test (HIT-6)
131 questionnaire but failed to identify any correlations between frequency and headache-
132 related disability [13] in patients with headache including patients with TTH. In the general
133 population, individuals with higher impact associated to TTH have a lower quality of life,
134 are more frequently absent from work and have lower work performance [14]. A Danish
135 population-based study reported that absence rates were higher in individuals with FTTH
136 than in healthy subjects [15]. Headache-related disability assessed by the HIT-6
137 questionnaire appear to be an important aspect to monitor in TTH patients to evaluate
138 changes over time and to guide futures clinical interventions. The aim of this study was to
139 determine if headache profile can predict future disability in participants with tension-type
140 headache.

141

142 **Materials and Methods**

143 *Study design*

144 This cohort study was conducted at the Laboratory of motor control and neuromechanics
145 located at the University of Québec in Trois-Rivières. Recruitment, testing and follow-up
146 were conducted from August 2016 to November 2017. This study falls within the
147 continuity of a study with controls and TTH participants (Marchand et al. in press in BMJ).

148

149 *Participants' selection*

150 Eighty-three participants with tension-type headache were recruited via social media
151 platforms and from the university community. To be included in the study, participants
152 needed to fulfill the International Headache Society (IHS) classification criteria for IETTH,
153 FETTH or CTTH (see table 1). Participants with concomitant headache and neck pain were
154 included only if neck pain was not the dominant pain perceived. For participants
155 experiencing other concomitant headache type, presentation and symptoms of TTH were
156 discussed at baseline and only patients for which tension-type headache was the main
157 headache type were included. Participants with concomitant migraine were included only
158 if their episodes were infrequent. Distinction between headaches types were clearly
159 highlighted at baseline as the goal of the study was to track only information related to
160 TTH over the study period. However, participants were asked to report when they were
161 affected by another headache type during the follow-up period. Exclusion criteria included
162 having a recent history of cervical spine trauma, recent whiplash, neck fracture, surgery or
163 malignant lesion, infection, medication overuse, having a diagnosis of fibromyalgia and
164 having neurological deficits, spasmodic torticollis, presence of upper limb pain or lack of
165 tension type headache episodes. Participants with neck, head or shoulder pain due to an
166 injury were excluded from this study as well as participants with all forms of pain whose
167 frequency and intensity could interfere with headaches. Participants were not allowed to
168 participate if they were under a course of treatment for headache tension-type headache or
169 for neck pain. Pregnant women were also excluded from the experimentation because of
170 the prone position adopted during the neck extension task.

171


172 Table 1: International Headache Society (IHS) classification criteria for IETTH, FETTH
 173 or CTTH

Infrequent episodic tension-type headache	Frequent episodic tension-type headache	Chronic tension-type headache
A. At least 10 episodes of headache occurring on <1 day/month on average (<12 days/year) and fulfilling criteria B-D	A. At least 10 episodes of headache occurring on 1-14 days per month on average for >3 months (≥ 12 and <180 days per year) and fulfilling criteria B-D	A. Headache occurring on ≥ 15 days per month on average for >3 months (≥ 180 days per year) and fulfilling criteria B-D
B. Lasting from 30 minutes to 7 days	B. Lasting from 30 min to 7 days	B. Lasting hours to days, or unremitting
<p style="text-align: center;">C. At least two of the following four characteristics:</p> <ol style="list-style-type: none"> 1. bilateral location 2. pressing or tightening (non-pulsating) quality 3. mild or moderate intensity 4. not aggravated by routine physical activity such as walking or climbing stairs 		
D. Both of the following: <ol style="list-style-type: none"> 1. no nausea or vomiting 2. no more than one of photophobia or phonophobia 	D. Both of the following: <ol style="list-style-type: none"> 1. no nausea or vomiting 2. no more than one of photophobia or phonophobia 	D. Both of the following: <ol style="list-style-type: none"> 1. no more than one of photophobia, phonophobia or mild nausea 2. neither moderate or severe nausea nor vomiting

174 This study was approved by the University's Research Ethic Committee for human subjects
175 (CER-16-225-07.15). All participants provided informed written consent prior to their
176 participation in the study.

177

178 *Procedures*



179

180 Figure 1 : Timeline of clinical and physical outcomes

181 The first session began with a brief history taking to obtain demographic data as well as
182 information regarding typical episodes of TTH and neck pain over the last month.
183 Headache and neck pain maximum and mean intensities were measured using a 10 cm
184 visual analogue scale (VAS). Participants were inquired about frequency of headache and
185 neck pain episodes over the past month. All clinical and physical outcomes were obtained
186 at baseline. In addition, headache frequency and intensity were monitored daily for 30 days
187 and two self-reported questionnaires (6-item Headache Impact Test and Neck disability

188 Index) were completed electronically using an online survey at the 1- and 3-month follow-
189 ups (see figure 1).

190

191 *Clinical Outcomes*

192 6-item Headache Impact Test

193 The validated French version of HIT-6 questionnaire was used to assess disability related
194 to headaches [16]. The HIT-6 is a 6-item, retrospective and self-reported questionnaire.
195 This questionnaire addresses several quality of life components of such as pain, cognitive
196 functioning, role functioning, vitality, social functioning and psychological distress [13,
197 17]. Participants were asked to complete the questionnaire based on the past four weeks
198 [13]. The total score obtained was calculated by adding scores from each question for a
199 maximum of 78 points [13]. Higher scores reveal greater headache-related disability.

200

201 Neck Disability Index (NDI)

202 The validated French version of NDI was used to evaluate disability related to neck pain
203 [18]. This is a 10-item questionnaire related to cervical pain and the impact on everyday
204 life as, for example, pain intensity, headache, concentration, reading, driving and work
205 [19]. The total score obtained was calculated by adding scores from each question for a
206 maximum of 50 points and higher scores reveal greater neck-related disability [18].

207

208 Kinesiophobia, anxiety and self-efficacy

209 At baseline, participants were asked to complete three other questionnaires related to
210 kinesiophobia, anxiety and self-efficacy. Kinesiophobia was assessed using the validated

211 French version of the Tampa Scale of Kinesiophobia (TSK) a 17-items questionnaire which
212 allow to quantify fear of movement with higher score reflecting an increased level of
213 kinesiophobia [20]. To assess anxiety, the validated French-Canadian version of the state
214 trait anxiety inventory (STAI-Y) was used [21]. This 20-item questionnaire allows to
215 evaluate anxiety as a personality trait and anxiety as an emotional state related to a
216 particular situation [21]. Finally, self-efficacy was assessed by the validated French general
217 self-efficacy scale (GSE) which allows the evaluation of individuals perception to meet the
218 needs of tasks in different contexts [22, 23].

219

220 30-day data collection

221 On a daily basis, participants were asked for 30 days about the presence and intensity of
222 headaches within the last 24 hours. If any, they were invited to identify the type of headache
223 they had (TTH or migraine for participants known to have concomitant types of headache).
224 Based on participants' preferences, they were contacted by e-mails (5) or text messages
225 (43) and some people (11) preferred to complete a headache diary.


226

227 ***Physical Outcomes***

228 Maximum voluntary contraction (MVC) of neck-extensor muscles was tested in a prone
229 position on a table with the participant's head and the neck past the edge of the table. To
230 ensure minimal recruitment of thoracic and scapular muscles, the cervico-thoracic junction
231 was stabilized with a strap (see figure 2). To evaluate the strength of the neck extensor
232 muscles, another strap was disposed over the protuberancia occipitalis and was anchored
233 to the floor. The head strap was adjusted to ensure that participants' head was stabilized in

234 neutral horizontal position throughout testing. Participants were then asked to perform
235 three neck extensors MCV while keeping the neutral horizontal position of the neck and
236 the head. To perform the neck extensors MVC, participants were asked to progressively
237 increase muscle contraction until maximum, hold the maximum for 3 to 5 seconds and then
238 release. Maximum voluntary contractions were recorded using a force gauge (Model
239 IPM250; Futek Advanced Sensor Technology Inc, Irvine, CA, USA). The first trial was
240 performed to familiarized participants with the isometric extension contraction and a
241 further two trials were conducted after the familiarization task. Each trial was followed by
242 a period of rest of one minute.

243



244

245 Figure 2: Participants position during the evaluation of neck extensor maximal isometric strength

246

247 ***Data and statistical analyses***

248 Statistical analyses were performed using STATISTICA statistical package version 10
249 (Statsoft, Tulsa, OK), and the level of significance was set at $p < .05$. Because of the


250 asymmetrical data distribution (skewness) of several variables, data were normalized
251 whenever it was deemed necessary using log-transformation. Self-reported headache
252 frequency and intensity were compared to the data obtained from the 30-day daily
253 monitoring using dependent t-tests to assess participants' ability to estimate their headache
254 characteristics. Repeated ANOVA were performed for HIT-6 and NDI questionnaires to
255 assess the evolution of participants' headache and neck-related disabilities and Tukey post-
256 hoc tests were performed to identify significant differences between baseline, the 1-month
257 and the 3-month follow-up for HIT-6 and NDI. Correlations between headache frequency,
258 headache intensity, neck pain frequency, neck pain intensity, kinesiophobia, anxiety, self-
259 efficacy, neck-related disability and headache-related disability at the 3-month follow-up
260 were evaluated using the Pearson's correlation coefficient. Multiple regressions analysis
261 were conducted using the highest correlations value between all determinants and HIT-6
262 questionnaire at the 3-month follow-up to test if any variables predicted future headache-
263 related disability over a 3-month period. Significant correlated determinants were added
264 into the stepwise regression model and determinants that significantly contributed to
265 headache-related disability over a 3-month period were identify.

266

267 **Results**

268 Eighty-three participants were included at baseline. Twelve participants were lost at the 1-
269 month follow-up and 8 others were lost at the 3-month follow-up because participants did
270 not return the questionnaires (see figure 3). Before excluding a participant, a reminder was
271 sent on 3 different occasions to any participant who did not return their questionnaires. A

272 total of 59 participants were included in the analysis. Three participants had concomitant
273 infrequent migraine.



274

275 Figure 3: Flowchart of TTH participants enrollment and reasons for exclusion of the
276 analysis

277 Baseline demographics

278 Means scores and standard deviations were calculated for all clinical and physical baseline
279 outcomes (see table 2). Participants presented with low fear of movement (27.80±5.86) and
280 anxiety (35.64±9.47) mean scores and a high self-efficacy mean score (35.42±3.24). The
281 mean score for headache-related disability indicated moderate headache impact (score ≥50
282 points for the HIT-6 questionnaire).

283

284 Table 2: Participant's baseline results for clinical and physical outcomes

	<i>Variables</i>	<i>Mean</i>	<i>SD</i>
<i>Demographics</i>	Age (years)	27.88	9.41
	F:M	40:19	N/A
	Weight (kg)	67.72	14.12
	Height (m)	1.67	0.09
	BMI (kg/m ²)	24.18	4.10
	Years with headache	6.22	6.55
	Kinesiophobia (17-68)	27.80	5.86
	Self-efficacy scale (10-40)	35.42	3.24
	Anxiety (20-80)	35.64	9.47
<i>Headache</i>	Self-reported frequency of headache (previous month)	6.58	8.06
	Self-reported mean intensity of headache (/10)	3.93	1.77
	HIT-6 (36-78)	51.17	9.31
<i>Neck pain</i>	Self-reported frequency of neck pain (previous month)	5.97	8.15
	Self-reported mean intensity of neck pain (/10)	2.58	1.89
	NDI (/50)	5.52	4.99
<i>Strength</i>	MVC (N)	97.57	34.86

285 SD= standard deviation, F= female, M= male, BMI= body mass index, HIT-6= 6-item
286 headache impact test, NDI= neck disability index, MVC= maximum voluntary contraction.

287

288 Table 3 : Participant's results regarding frequency and intensity of headache for the 30-day
289 data collection

<i>Variables</i>	<i>Mean</i>	<i>SD</i>
<i>Frequency of headache (/30 days)</i>	7.75	6.84
<i>Intensity (/10)</i>	2.97	1.51

290

291

292 Frequency and intensity of headache

293 T-test for dependent samples comparing self-reported frequency (mean= 6.58; SD = 8.06)
294 and frequency assess at 30 days (7.75; SD= 6.69) revealed no significant difference
295 between self-reported headache frequency and 30 days data collection (p = 0.107).
296 However, there was a significant difference (p < 0.001) between self-reported headache
297 intensity (mean = 3.93; SD = 1.77) and headache intensity obtain from the 30-day data
298 collection (mean = 2.97; SD = 1.51) (see table 2 for self-reported data and table 3 for 30-
299 day data collection).


300

301 HIT-6 and NDI score evolution

302 The ANOVA indicated a significant effect of time on headache-related disability (F(2,
303 116)=4.53, p=0.013) and a significant effect of time on neck-related disability (F(2,

304 116)=4.89, $p=0.009$). Tukey post-hoc test showed a significant decrease in headache-
305 related disability between baseline and the 1-month follow-up ($p=0.036$) and between
306 baseline and the 3-month follow-up ($p=0.021$) (see figure 4) and showed a significant
307 diminution in neck-related disability between baseline and the 3-month follow-up
308 ($p=0.007$) (see figure 5).

309




310

311

312 Figure 4: HIT-6 score evolution from baseline to the 3-month follow-up in TTH participants.

313 * : $p= 0.025$, ** : $p= 0.017$

314



315

316 Figure 5 : NDI score evolution from baseline to the 3-month follow-up in TTH participants.

317 * : p=0.007

318

319 Table 4: Correlations between determinants and headache-related disability

	<i>Variables</i>	<i>HIT-6 (3-month)</i>	
		Correlation coefficient (r)	p
<i>Baseline</i>	MVC of neck extensors	-0.16	0.236
	Years with headache	0.53	<0.001
	Self-reported headache frequency	0.52	<0.001

	Self-reported headache intensity	0.61	<0.001
	Self-reported neck pain frequency	0.21	0.115
	Self reported neck pain intensity	0.29	0.025
	Kinesiophobia	-0.009	0.945
	Anxiety	0.28	0.031
	Self-efficacy	-0.10	0.437
	HIT-6	0.72	<0.001
	NDI	0.64	<0.001
30 days	Headache frequency	0.60	<0.001
data collection	Headache intensity	0.54	<0.001
1 month	HIT-6	0.86	<0.001
Follow-up	NDI	0.65	<0.001

320 MVC = Maximal Voluntary Contraction, HIT-6 = 6-item Headache Impact Test, NDI =

321 Neck Disability Index

322

323 Table 5: Predictors of headache-related disability

<i>Variables</i>	<i>β value</i>	<i>p</i>
------------------	----------------	----------

<i>Years with headaches</i>	0.087	0.460
<i>Self-reported neck pain intensity</i>	-0.073	0.508
<i>Headache frequency in 30 days</i>	0.284	0.032
<i>Headache intensity in 30 days</i>	0.253	0.024
<i>Anxiety</i>	0.100	0.304
<i>NDI (baseline)</i>	0.245	0.112
<i>Presence of concomitant migraine</i>	0.206	0.044
Multiple regression: $R^2 = 0.583$		

324

325 Predictors of headache-related disability at 3 months

326 High correlations were found between self-reported headache frequency, self-reported
327 headache intensity and HIT-6 score at the 3-month follow-up. However, self-reported
328 headache frequency and intensity were excluded from the multiple regression because
329 frequency and intensity were also prospectively collected for 30 days and were considered
330 more specific than self-reported data estimated at baseline considering the past month. Self-
331 reported neck pain frequency, kinesiophobia and self-efficacy were also excluded because
332 they did not correlate with HIT-6 at 3 months nor with any other outcome measures (see
333 table 5). Number of years with TTH ($r=.53$; $p<0.001$), self-reported neck pain intensity
334 ($r=.29$; $p=0.025$), headache frequency ($r=.60$; $p<0.001$), headache intensity ($r=.54$;
335 $p<0.001$), anxiety ($r=.28$; $p=0.031$) as well as neck-related disability at baseline ($r=.64$;
336 $p<0.001$) were included in the multiple regression model because of their high correlation
337 with the HIT-6 score at 3 months. Presence of concomitant infrequent migraine was added
338 as a covariable to the multiple regression model given that migraine is known to be a

339 predictor of poor outcome in TTH [12]. Number of years with TTH, self-reported neck
340 pain intensity, anxiety and neck-related disability at baseline were highly correlated with
341 headache-related disability at 3-month. However, in the stepwise regression model, none
342 of them remained significantly associated with headache-related disability at 3-month (p
343 values = 0.460, 0.508, 0.304 and 0.112 respectively). Results showed that headache
344 frequency, headache intensity and the presence of concomitant infrequent migraine were
345 good predictors of headache-related disability at 3 months. All together, predictors were
346 able to predict 58.3 % of the headache-related disability variance. Results of the regression
347 model are presented in table 5.

348

349 **Discussion**

350 The purpose of this study was to determine if headache profile can predict future disability
351 in participants with TTH. Participants' mean age was 27.88 ± 9.41 years which is consistent
352 with the age group with the highest TTH prevalence in the literature (25-30 years old). The
353 results showed that 30 days mean headache frequency and intensity as well as self-reported
354 mean intensity of neck pain, years with TTH, neck-related disability and presence of
355 concomitant infrequent migraine can predict future disability in patients with TTH.

356

357 **Headaches characteristics**

358 With regards to participants' headache characteristics, the results showed a high correlation
359 between the number of years with TTH and headache-related disability and between
360 number of TTH episodes in a month and headache-related disability as well as a moderate
361 correlation between TTH intensity and headache-related disability. Mild to moderate pain

362 intensity represents the clinical criteria required in the diagnosis of TTH and based on the
363 present results moderate intensity had a higher impact on headache-related disability. In
364 this study, multiple regressions showed that intensity is an important predictor of future
365 disability, but headache frequency was the most important predictive factor of headache-
366 related disability. These results are in accordance with a previous study which showed that
367 frequent headache categories (FETTH and CTTH) are associated with higher headache-
368 related impact than the infrequent headache category [24]. Kim et al. (2015) also reported
369 an increased burden of headache-related disability associated to the chronic form compared
370 to infrequent and frequent episodic forms [25]. TTH patients seem to be more affected by
371 the number of headache episodes than by their intensity. Suffering from TTH on a regular
372 basis seems to influence psychological well being. In fact, frequent headache and disability
373 seem to impair quality of life [26, 27]. Headache intensity contributes to occasional
374 disability in a month or in a year while a higher number of headaches in a month can
375 contribute to higher disability. A previous study showed the impact of ETTH and CTTH
376 and their contribution to absenteeism and presenteeism in TTH patients [28].

377

378 Neck pain and neck extensor muscle strength

379 In the current study, neck pain and neck extensor muscle MVC were assessed because neck
380 pain has been traditionally linked with tension type headache [29, 30] and neck pain has
381 been shown to be more prevalent in TTH patients than in individuals with no TTH [31].
382 Neck pain has previously been associated with decreased neck muscle strength [32, 33].
383 The present study did not reveal any correlation between neck extensor muscle strength
384 and headache-related disability. In addition, neck muscle strength was not retained among

385 predictive factors of future headache-related disability. However, results showed that
386 concomitant neck pain was present in TTH patients with a self-reported mean of 5.97 days
387 (SD = 8.15) which is consistent with the literature [31]. Results also showed that neck pain
388 intensity was correlated with a higher score in HIT-6 at the 3-month follow-up ($r=0.29$)
389 and that it can be considered as a predictive determinant of future disability. Ashina et al.
390 (2015) found a correlation between neck pain frequency and the frequency of TTH and
391 suggested a possible shared pathophysiological mechanism between neck pain and primary
392 headache including tension-type headache [31]. Regarding neck extensor muscle strength,
393 previous studies showed a decrease in neck extensor muscle force production in TTH
394 patients compared to healthy controls [33, 34]. However, our results showed that MVC is
395 not a physical determinant of future disability as evaluated by the HIT-6 questionnaire.

396

397 HIT-6 and NDI

398 Findings of the present study indicated statistically significant decreases in headache-
399 related disability and neck-related disability over the 3-month follow-up period but these
400 differences were not clinically significant. To be considered clinically significant, the
401 minimally clinical important change in TTH related disability should reach 8 points on the
402 HIT-6 score [17]. The results of the present study showed a decrease of only 2.07 points
403 on the HIT-6 mean score between baseline and the 3-month follow-up. Regarding the neck
404 disability, a difference of 3.5 points on the NDI scale represents the minimally clinical
405 important change [35] and the present results indicated a decrease of 1.21 points between
406 the baseline and 3-month scores therefore not reaching the threshold for neck-related
407 disability minimally clinical important change. The decrease for HIT-6 and NDI scores

408 found in the present study can be explained by the natural fluctuation of tension-type
409 headache over time. Indeed, prognostic factors of TTH recovery are less severe headache,
410 mild headache-related disability, not using medication, absence of anxiety, sleep problem,
411 depression or other pain [36].

412

413 Kinesiophobia, anxiety and self-efficacy

414 The results showed low levels of kinesiophobia and high self-efficacy scores and these
415 scores were not correlated to any of the clinical or physical outcomes. The current results
416 showed that anxiety was correlated with headache-related disability. Anxiety has been
417 extensively explored in TTH and some studies found that anxiety seemed to be more
418 dependent on the TTH frequency [37, 38] which means that CTTH patients would be more
419 anxious than ETTH patients. Results of the present study are consistent with these
420 previous findings.

421

422 Limitations

423 This study is not without limitations. Indeed, baseline neck pain frequency and intensity
424 were self-reported by participants based on episodes over the past month, which could have
425 been influenced by recollection bias. Although the comparison between retrospective and
426 prospective self-reported data showed that participants were able to correctly estimate
427 headache frequency, they overestimated headache intensity which could also have been
428 overestimated in self-reported neck pain intensity. In addition, in the present study, TTH
429 participants results were analyzed without considering TTH categories (infrequent
430 episodic, frequent episodic and chronic), and it should be kept in mind that results could

431 differ between the episodic forms and the chronic form. Another limitation is our small
432 number of participants, meaning that the results should be interpreted with caution.

433

434 Results of the present study showed that TTH frequency and intensity, and the presence of
435 concomitant infrequent migraine are predictors of future disability over a 3-month period.

436 Results also showed that neck extensor muscles strength was not correlated with headache-
437 related disability or with any other clinical outcomes and was not a good predictor of future

438 disability. Further studies are needed to evaluate the predictive value of other physical

439 outcomes on headache-related disability. Tension-type headache constitutes a major public

440 health problem and a better understanding of clinical and physical factors is needed. Health

441 professionals should consider clinical outcomes to evaluate and elaborate future treatment

442 strategies for patients with TTH.

443

444 **Declarations:**

445 **Ethics approval and consent to participate:** This study has been approved by the ethic

446 committee of human subjects of the Université du Québec à Trois-Rivières (CER-16-225-

447 07.15).

448 **Availability of data and material :** The datasets used and/or analysed during the current

449 study are available from the corresponding author on reasonable request.

450 **Competing interests :** None to declare.

451

- 453 1. Jensen, R.H., *Tension-Type Headache - The Normal and Most Prevalent Headache*.
454 Headache, 2018. **58**(2): p. 339-345.
- 455 2. Jensen, R. and L.J. Stovner, *Epidemiology and comorbidity of headache*. Lancet Neurol,
456 2008. **7**(4): p. 354-61.
- 457 3. *The International Classification of Headache Disorders, 3rd edition (beta version)*.
458 Cephalalgia, 2013. **33**(9): p. 629-808.
- 459 4. Rasmussen, B.K., et al., *Epidemiology of headache in a general population--a prevalence*
460 *study*. J Clin Epidemiol, 1991. **44**(11): p. 1147-57.
- 461 5. Crystal, S.C. and M.S. Robbins, *Epidemiology of tension-type headache*. Current pain and
462 headache reports, 2010. **14**(6): p. 449-454.
- 463 6. Schwaiger, J., et al., *Prevalence of primary headaches and cranial neuralgias in men and*
464 *women aged 55-94 years (Bruneck Study)*. Cephalalgia, 2009. **29**(2): p. 179-87.
- 465 7. Russell, M.B., *Tension-type headache in 40-year-olds: a Danish population-based sample*
466 *of 4000*. The journal of headache and pain, 2005. **6**(6): p. 441-447.
- 467 8. Ashina, S., L. Bendtsen, and M. Ashina, *Pathophysiology of tension-type headache*. Curr
468 Pain Headache Rep, 2005. **9**(6): p. 415-22.
- 469 9. Ashina, S., A. Lyngberg, and R. Jensen, *Headache characteristics and chronification of*
470 *migraine and tension-type headache: A population-based study*. Cephalalgia, 2010. **30**(8):
471 p. 943-52.
- 472 10. Hagen, K., et al., *Lifestyle factors and risk of migraine and tension-type headache. Follow-*
473 *up data from the Nord-Trondelag Health Surveys 1995-1997 and 2006-2008*. Cephalalgia,
474 2018: p. 333102418764888.
- 475 11. Juang, K.D., et al., *Comorbidity of depressive and anxiety disorders in chronic daily*
476 *headache and its subtypes*. Headache, 2000. **40**(10): p. 818-23.
- 477 12. Lyngberg, A.C., et al., *Prognosis of migraine and tension-type headache: a population-*
478 *based follow-up study*. Neurology, 2005. **65**(4): p. 580-5.
- 479 13. Sauro, K.M., et al., *HIT-6 and MIDAS as measures of headache disability in a headache*
480 *referral population*. Headache, 2010. **50**(3): p. 383-95.
- 481 14. Jensen, R., *Diagnosis, epidemiology, and impact of tension-type headache*. Current pain
482 and headache reports, 2003. **7**(6): p. 455-459.
- 483 15. Lyngberg, A.C., et al., *Secular changes in health care utilization and work absence for*
484 *migraine and tension-type headache: a population based study*. Eur J Epidemiol, 2005.
485 **20**(12): p. 1007-14.
- 486 16. Magnoux, E., M.A. Freeman, and G. Zlotnik, *MIDAS and HIT-6 French translation: reliability*
487 *and correlation between tests*. Cephalalgia, 2008. **28**(1): p. 26-34.
- 488 17. Castien, R.F., et al., *Minimal clinically important change on the Headache Impact Test-6*
489 *questionnaire in patients with chronic tension-type headache*. Cephalalgia, 2012. **32**(9): p.
490 710-4.
- 491 18. Wlodyka-Demaille, S., et al., *French translation and validation of 3 functional disability*
492 *scales for neck pain*. Arch Phys Med Rehabil, 2002. **83**(3): p. 376-82.
- 493 19. Macdermid, J.C., et al., *Measurement properties of the neck disability index: a systematic*
494 *review*. Journal of orthopaedic & sports physical therapy, 2009. **39**(5): p. 400-C12.
- 495 20. French, D.J., et al., *Fear of movement/(re)injury in chronic pain: a psychometric*
496 *assessment of the original English version of the Tampa scale for kinesiophobia (TSK)*. Pain,
497 2007. **127**(1-2): p. 42-51.

- 498 21. Gauthier J, B.S., *Adaptation canadienne française de la forme révisée du State Trait*
499 *Anxiety Inventory de Spielberger*. Rev Can Sci Comport 1993. **25**: p. 559–578.
- 500 22. Scholz, U., et al., *Is general self-efficacy a universal construct? Psychometric findings from*
501 *25 countries*. European journal of psychological assessment, 2002. **18**(3): p. 242.
- 502 23. Chen, G., S.M. Gully, and D. Eden, *Validation of a new general self-efficacy scale*.
503 *Organizational research methods*, 2001. **4**(1): p. 62-83.
- 504 24. Fumal, A. and J. Schoenen, *Tension-type headache: current research and clinical*
505 *management*. The Lancet Neurology, 2008. **7**(1): p. 70-83.
- 506 25. Kim, B.-S., et al., *Factors associated with disability and impact of tension-type headache:*
507 *findings of the Korean headache survey*. The Journal of Headache and Pain, 2015. **16**(1):
508 p. 40.
- 509 26. Cassidy, E.M., et al., *Factors associated with burden of primary headache in a specialty*
510 *clinic*. Headache, 2003. **43**(6): p. 638-44.
- 511 27. Marcus, D.A., *Identification of patients with headache at risk of psychological distress*.
512 *Headache: The Journal of Head and Face Pain*, 2000. **40**(5): p. 373-376.
- 513 28. Robbins, M.S. and R.B. Lipton. *The epidemiology of primary headache disorders*. in
514 *Seminars in neurology*. 2010. © Thieme Medical Publishers.
- 515 29. Marcus, D.A., et al., *Musculoskeletal abnormalities in chronic headache: a controlled*
516 *comparison of headache diagnostic groups*. Headache, 1999. **39**(1): p. 21-7.
- 517 30. Sohn, J., et al., *Differences in cervical musculoskeletal impairment between episodic and*
518 *chronic tension-type headache*. Cephalalgia, 2010. **30**(12): p. 1514-1523.
- 519 31. Ashina, S., et al., *Prevalence of neck pain in migraine and tension-type headache: a*
520 *population study*. Cephalalgia, 2015. **35**(3): p. 211-9.
- 521 32. Scheuer, R. and M. Friedrich, *Reliability of isometric strength measurements in trunk and*
522 *neck region: patients with chronic neck pain compared with pain-free persons*. Arch Phys
523 *Med Rehabil*, 2010. **91**(12): p. 1878-83.
- 524 33. Madsen, B.K., et al., *Neck and shoulder muscle strength in patients with tension-type*
525 *headache: A case-control study*. Cephalalgia, 2016. **36**(1): p. 29-36.
- 526 34. Madsen, B.K., et al., *Neck/shoulder function in tension-type headache patients and the*
527 *effect of strength training*. J Pain Res, 2018. **11**: p. 445-454.
- 528 35. Pool, J.J., et al., *Minimal clinically important change of the Neck Disability Index and the*
529 *Numerical Rating Scale for patients with neck pain*. Spine, 2007. **32**(26): p. 3047-3051.
- 530 36. Boardman, H., et al., *The natural history of headache: predictors of onset and recovery*.
531 *Cephalalgia*, 2006. **26**(9): p. 1080-1088.
- 532 37. Peñacoba-Puente, C., et al., *Interaction between anxiety, depression, quality of life and*
533 *clinical parameters in chronic tension-type headache*. European Journal of Pain, 2008.
534 **12**(7): p. 886-894.
- 535 38. Song, T.-J., et al., *Anxiety and depression in tension-type headache: a population-based*
536 *study*. PloS one, 2016. **11**(10): p. e0165316.

537

538

539