Short term effect of delayed-onset muscle soreness on trunk proprioception during force reproduction tasks in a healthy adult population: a crossover study

Authors informations:

Mariève Houle¹, Catherine Daneau², Arianne Lessard¹, Marie-Andrée Mercier¹, Martin Descarreaux¹, Jacques Abboud¹

¹ Department of Human Kinetics, Université du Québec à Trois-Rivières, 3351 Boul. des Forges, Trois-Rivières, Qc, G8Z 4M3, Canada

² Department of Anatomy, Université du Québec à Trois-Rivières, 3351 Boul. des Forges, Trois-Rivières, Qc, G8Z 4M3, Canada

Corresponding author: Jacques Abboud, 3351, boul. des Forges, C.P. 500, Trois-Rivières, Qc, Canada, G8Z 4M3. Telephone number: +1 (819) 376-5011 ext 3783. E-mail: <u>jacques.abboud@uqtr.ca</u>

Conflict of Interest Statement: There is no conflict of interest.

Funding: This research received funding from Natural Sciences and Engineering Research Council of

Canada (NSERC).

1 Abstract:

Purpose: The aim of this study was to evaluate the effects of lumbar muscle delayed-onset muscle soreness (DOMS) on the ability of the trunk muscles to reproduce different levels of force.

Methods: Twenty healthy adults (10 males and 10 females) were recruited for this study. Force reproduction in trunk extension and flexion was assessed at 50 and 75% of participants' maximal isometric voluntary contraction in flexion and extension before and after a lumbar muscle DOMS protocol. Trunk proprioception was evaluated and compared between these conditions using different variables such as constant errors (CE), absolute errors (AE), variable errors (VE) and time to peak force (TPF). For each variable, repeated measure ANOVAs were conducted.

Results: AE were higher when participants had to reach the target post-DOMS protocol in extension 11 compared to flexion and in presence of higher demand of force (p=0.02). For VE, results showed that 12 participants were more variable in extension than in flexion when the required force was higher (p=0.04). 13 CE variable was higher when participants had to reach the force target in extension compared to flexion 14 under the effect of DOMS (p=0.02). Results also showed that participants took less time to reach the force 15 target post-DOMS protocol in extension (0.62 ± 0.20 sec) and in flexion (0.53 ± 0.19 sec) than pre-DOMS 16 protocol in extension (0.55 ± 0.15) and in flexion (0.50 ± 0.20) (p<0.001).

17 Conclusion: Lumbar muscle DOMS affect trunk proprioception during force reproduction tasks
18 especially in trunk extension and at higher force.

19 Keys words: Delayed-onset muscle soreness, lumbar, sensorimotor control, pain, proprioception

 List of abbreviations DOMS: Delayed-onset muscle soreness MVC: Maximum voluntary contraction IPAQ: International physical activity questionnaire CE: Constant error AE: Absolute error VE: Variable error VE: Variable error TPF: Time to peak force 36 37 	
 27 End of abore viations 28 DOMS: Delayed-onset muscle soreness 29 MVC: Maximum voluntary contraction 30 IPAQ: International physical activity questionnaire 31 CE: Constant error 32 AE: Absolute error 33 VE: Variable error 34 TPF: Time to peak force 35 36 37 	
 ⁷ 20 DOMS: Delayed-onset induct solutions ⁸ 29 MVC: Maximum voluntary contraction ⁹ 30 IPAQ: International physical activity questionnaire ¹¹ 31 CE: Constant error ¹³ 32 AE: Absolute error ¹⁵ 33 VE: Variable error ¹⁶ 34 TPF: Time to peak force ¹⁹ 35 ²¹ 36 ²³ 37 	
 ⁹ 30 IPAQ: International physical activity questionnaire ¹¹ 31 CE: Constant error ¹³ 32 AE: Absolute error ¹⁵ 33 VE: Variable error ¹⁷ 34 TPF: Time to peak force ¹⁹ 35 ²¹ 36 ²³ 37 	
 So FrAQ: International physical activity questionnalite 31 CE: Constant error 32 AE: Absolute error 33 VE: Variable error 34 TPF: Time to peak force 35 36 37 	
12 31 CE: Constant error 13 32 AE: Absolute error 15 33 VE: Variable error 16 33 VE: Variable error 17 34 TPF: Time to peak force 19 35 21 36 23 37	
14 32 AE: Absolute error 15 33 VE: Variable error 17 34 TPF: Time to peak force 19 35 21 36 23 37	
16 33 VE: Variable error 17 34 TPF: Time to peak force 19 35 20 35 21 36 23 37	
18 34 TPF: Time to peak force 19 35 20 35 21 36 23 37	
20 35 21 36 23 37	
²¹ 36 ²³ 37	
²³ 37	
24	
25 38 26	
27 39 28	
29 40 30	
31 41 32	
33 42	
35 43	
36 37 44	
³⁸ 39 45	
⁴⁰ ₄₁ 46	
42 43 47	
44 45 48	
46 49	
47 48 50	
49 50 51	
51 52 52	
53 54 53	
55 56 54	
57 58 55	
59 60 56	
61	
62 63	

57 Introduction

Delayed-onset muscle soreness (DOMS) can be defined as musculoskeletal pain and soreness and as a sensation of discomfort (Cleak and Eston 1992) that lasts for several days (Weerakkody et al. 2001) and that is induced by unusual intense exercises and/or eccentric contractions (Coudreuse et al. 2004; MacIntyre et al. 1995; Proske et al. 2003). The DOMS effects peak between 24 to 72 hours following those exercises and disappear progressively in three to five days (Cheung et al. 2003; Coudreuse et al. 2004). DOMS is usually associated with inflammation and muscle damage and individuals presenting DOMS can experience muscle stiffness, pain and/or movement restrictions (Farias-Junior et al. 2019) and a decrease of maximal muscle strength (Abboud et al. 2019). Effects of DOMS on proprioception are characterized by a significant increase in errors in upper limb positioning and force reproduction tasks (Proske et al. 2003), suggesting that, when muscles become sore in the presence of DOMS or following paralysis, the motor command sent to muscles is not relevant to the desired outcome. Proprioception includes the sense of limb position and movement, and the sense of force and effort (Jerosch and Prymka 1996). Proprioception can be evaluated by repositioning tasks and force reproduction tasks, which mainly assess conscious proprioception (Hagert 2010). Trunk proprioception in lumbar muscles as well as abdominal muscles can be altered in patients with chronic or recurrent low back pain (Hodges and Richardson 1999; Rausch Osthoff et al. 2015; Tong et al. 2017). However, because of within- and between- patient variability in motor behaviour (van Dieen et al. 2017), drawing conclusion about the mechanism underlying proprioception alterations in this population is still challenging. Lumbar muscle DOMS is therefore a relevant experimental pain model

because of its ability to recreate altered motor functions, such as a decrease in lumbar muscles strength

- and an increase of fear of pain (Abboud et al. 2019; Bishop et al. 2011), which are also observed in lumbar
- 80 muscles of patients with chronic or recurrent non-specific low back pain (Hodges and Danneels 2019).

Furthermore, using DOMS as a pain model may help clarify the mechanism underlying
proprioception alterations in patients with low back pain. Therefore, the aim of this study was to evaluate,
in a healthy adult population, the effects of lumbar muscle DOMS during different force reproduction

conditions. We hypothesized that trunk proprioception in the direction of extension will be more altered
than in flexion and that this alteration will increase with higher force demand (della Volpe et al. 2006;
Proske et al. 2003).

Materials and Methods

Study design

We conducted a crossover study at the University [XXX] Laboratory. Recruitment and data
collection were completed from May to July 2018.

92 Participants

Twenty healthy adults, 10 females and 10 males, were recruited among the university community and employees and by social medias. To be included in the study, participants had to be back pain free. If they have experienced recurrent back pain or occasional pain in the last six months, they were not allowed to participate in this study. Other exclusion criteria were health conditions such as neuromuscular diseases, uncontrolled hypertension and heart disease, or cancer. Pregnant women were also excluded. The study was approved by the University humans research ethics board (CER-18-245-07.10) and written informed consent was obtained from each participant before the beginning of the experiment. Participants were advised that they had the possibility to withdraw from the study at any moment.

Experimental protocol

The experimental protocol was divided into two sessions separated by 24 to 36 hours. The period between the first and the second session was based on a previous study showing that pain and soreness following a lumbar muscle DOMS protocol peaks between 24- and 36 hours (Abboud et al. 2019). In the first session, participants were asked to fill in one questionnaire. Then, isometric muscles trunk extension and flexion maximum voluntary contraction (MVC) and different force reproduction tasks were evaluated for each participant. Finally, participants were asked to perform the lumbar muscle DOMS protocol. In

108	the second session, trunk extension and flexion MVC and force reproduction tasks were assessed again.
109	A timeline for clinical and physical outcome assessment is presented in Figure 1.
110	
111	[Insert figure 1. about here]
112	

Questionnaire

To assess level of physical activity, participants were invited to complete the short version of the International Physical Activity Questionnaire (IPAQ). Reliability and validity of the IPAQ short form have been tested in over twelve countries (Craig et al. 2003). This questionnaire is composed of 9 items assessing the intensity of physical activity habits in the past week (Lee et al. 2011).

118 Force reproduction tasks

All force reproduction conditions were performed on an isokinetic device (The LIDO Active Loredan Biomedical, West Sacramento, CA). Participants were semi-seated in a neutral position and they were attached to the device with four belts (Figure 2). Neutral position was defined as natural spine curve, hip angle was ~135° and knees were in full extension to minimize the contribution of lower limbs muscles and to better isolate trunk muscles during the force reproduction tasks. In fact, it has been shown that the pelvic stabilization increases the recruitment of low back muscles and decreases the contribution of hip extensors during dynamic lumbar extensions (da Silva et al. 2009). One belt was placed over the chest, another one was over the upper abdomen and the last two were over the hips and on the thighs. At first, three MVC were realized for both flexion and extension. The highest value of MVC for flexion and extension was used for the force reproduction tasks. Participants were free to experiment flexion and extension on the isokinetic device to familiarize with the equipment before performing MVC. For the flexion MVC, participants were told to push as hard as they can for 5 seconds against a resistance located at the sternum. For the extension MVC, the middle of the resistance was placed on the eighth thoracic vertebra. Then, trunk force reproduction was assessed in four conditions: trunk flexion and extension at 50% and 75% of MVC. Each condition of the force reproduction task was conducted both with and without visual feedback. The condition's order was randomized using computer random number generator (randomization.com) to minimize possible learning effects and residual muscle fatigue. Prior to recording the force reproduction task without visual feedback, practice trials within a 10% margin error of the target goal were allowed to each participant to get familiar with the task. Practice trials were stopped when 10 consecutive trials were performed within the margin of error. Participants were then asked to reproduce 10 trials of the same force level without visual feedback (Figure 3) and analyses were conducted considering these 10 repetitions without feedback. Participants were given a one-minute rest period between conditions to limit the occurrence of muscular fatigue. For all trials, participants were asked to provide a single impulse without correcting the force once the contraction was initiated. Participants were instructed to perform the task as quickly as possible. [Insert figures 2 and 3 about here] DOMS protocol First, participants completed three MVC in trunk extension on a 45-degrees Roman chair to evaluate lumbar extensors maximal strength (Figure 4; (Lariviere et al. 2011; Parreira et al. 2013). They had their trunk parallel to the floor in a prone position and were asked to push as hard as possible against a belt installed over the participant's shoulders. A load cell (Model IPM250; Futek Advanced Sensor Technology Inc, Irvine, CA, USA) was connected to the belt and gave indications about peak torque in trunk extension. The highest MVC value was used to determine a 10% external weight which was used for the entire endurance DOMS protocol. Then, participants were invited to complete the DOMS protocol that targeted low back muscles. The lumbar muscle endurance DOMS protocol was performed on the 45-degrees Roman chair in the same position used to establish MVC. In fact, participants initiated the DOMS protocol in a horizontal position with their trunk parallel to the ground. This protocol consisted of 5 sets of 20 repetitions of trunk flexion-extension with the 10% external weight in the hands and with two

minutes of rest between each set. A repetition consisted of (1) three seconds 30-degrees trunk flexion from horizontal (2) three seconds of isometric contraction and (3) one-second trunk extension starting from the flexion position to 30-degrees trunk extension from the horizontal (head, trunk and lower limbs needed to be in a neutral alignment). There were two indicators placed to help participants to complete the task adequately: one at 30-degrees trunk extension position and one at 30-degrees of trunk flexion. Participants hips and ankles were stabilized using straps to minimize pelvic tilt movements, which could limit the contribution of muscle groups other than paraspinal muscles during the DOMS protocol. A visual and auditory feedback was provided for participants during the protocol to help them following the tempo (3-3-1). Participants were motivated by verbal encouragements given by the assessors. The validation of the DOMS protocol was performed in a previous study (Abboud et al. 2019).

[Insert figures 4 about here]

172 Pain and Soreness Assessment

Lumbar muscle pain and soreness were assessed via text messages or emails sent to participants immediately following the first session (DOMS protocol). Text messages or emails were sent by one evaluator and this evaluator was not implicated neither in the lumbar muscle DOMS protocol and in force reproduction tasks. This evaluator was also naïve to expected results of the study. Data collection was completed over five consecutive days, three times a day. Participants received the message at 9 am, 3 pm and 9 pm (Figure 1). Participants were asked to rate the intensity of both lumbar muscle pain and lumbar muscle soreness using a 0-10-point scale. They were also asked to report any other side effects while answering daily text messages. During these five days, participants were asked to avoid any high intensity or unusual exercise or medication aiming to reduce pain or soreness. Based on the pain and soreness scores of each participant, the time it takes to higher level of pain and soreness were computed using the average time until the occurrence of the highest pain and soreness scores.

184 Dependent Variables

Constant error (CE), absolute error (AE), variable error (VE) and time to peak force (TPF) were calculated and compared between each condition (50% and 75% in extension and flexion) and each session (pre-DOMS and post-DOMS). These four variables are commonly used to assess trunk proprioception (Abboud et al. 2018; Boucher et al. 2015; Lee et al. 2010; McNair and Heine 1999). CE was the positive or the negative difference between the force value deployed by participants and the targeted force identified based on 50 or 75% of participants' MVC in extension or in flexion. AE was the absolute difference between the force value deployed by participants to reach the target and the force identified as the target. VE was defined by the peak force reach consistency compared with the average score of participants. TPF represented the time needed by participants to reach the force target.

194 Statistical Analysis

Analyses were performed using STATISTICA statistical package version 10 (Statsoft, Tulsa, OK), and the level of significance was set at $p \le 0.05$. Normality of distribution was assessed with the Kolmogorov–Smirnov test and by visual inspection. A mixed model three-way repeated measure ANOVAs were conducted to assess for each dependent variable: (1) the direction effect (flexion versus extension); (2) the force intensity effect (50 versus 75% MVC); (3) the DOMS effect (pre- versus post-DOMS); and (4) all the interaction effects. When necessary, the Tukey post-hoc test was performed as the post-hoc analysis for pairwise comparisons. Effects size of significant difference were calculated using

partial eta-squared (0.01 = small effect; 0.06 = medium effect; 0.14 = large effect).

Results

205 Baseline demographics

Twenty participants (10 females and 10 males) were included in the study and completed the protocol. Mean scores and standard deviation were calculated for all clinical and physical outcomes and are presented in Table 1. All participants experienced pain and/or soreness in the lumbar muscles. The

highest pain values were observed approximately 20 hours following the DOMS protocol, while the
highest soreness values occurred after 30 hours. Immediately after DOMS protocol, 2 participants
reported a light hyperalgesia in the thigh lasting 2 days due to the contact pressure point on the inclined
Roman bench.

[Insert table 1. about here]

Force reproduction task

Dependent variable means and standard deviations for each condition during pre-DOMS and post-DOMS protocol are presented in Table 2. ANOVAs results showed significantly higher values for all dependent variables (CE, AE, VE and TPF) in extension when compared to flexion (all $ps \le 0.01$; Table 3 and Figure 5). Moreover, a significant DOMS X Direction X Force interaction was found for AE (p=0.02) as illustrated in Figure 6. Results from the post-hoc test showed significantly higher AE value in extension post-DOMS in comparison to flexion post-DOMS protocol at 50% (p=0.046) and 75% (p=0.01). Results also showed a significant influence of force intensities (50% versus 75%), with higher value at 75% MVC for AE (p=0.03) and VE (p=0.04). A significant decrease was shown in TPF (p<0.001) between pre- and post-DOMS protocol. There was also a significant main effect of direction (extension versus flexion; p=0.01) showing that participants were poorer in extension than in flexion to reproduce the task. A significant Direction X Force intensity interaction effect for VE variable was also found (p=0.04) and Tukey post-hoc revealed that participants were more variable at 75% than at 50% in extension for VE $(p \le 0.003)$ but not in flexion (p = 0.74). Tukey post-hoc also showed that they were more variable in extension than in flexion for both forces (p=0.02 at 50% and $p\leq 0.001$ at 75%). Another significant Direction X DOMS interaction effect for the CE variable was observed (p=0.02). Tukey post-hoc showed that CE was higher in extension compared to flexion post-DOMS protocol ($p \le 0.001$). Post-hoc also showed that CE increased in extension post-DOMS protocol compared to flexion post-DOMS protocol (p=0.03). All other results were not statistically significant (Table 3).

[Insert tables 2 and 3 about here]

[Insert figures 5 and 6 about here]

240 Discussion

The objective of the present study was to evaluate the effect of lumbar muscle DOMS on trunk proprioception during different force reproduction tasks in a healthy adult population. Our hypothesis was that trunk proprioception in the direction of extension would be more altered than in flexion and that this alteration would increase with higher force demand. Results showed that participants (1) were more variable to reproduce forces (VE) in extension than in flexion regardless of the presence of lumbar muscle DOMS; (2) larger force production errors occurred for the higher level of force and more variability in the produced force was present in extension than in flexion; (3) under the influence of DOMS the performance to reach the force target in trunk extension was altered, while it remained unchanged in trunk flexion; (4) participants were faster in the force reproduction tasks under the influence of lumbar muscle DOMS.

251 Trunk Proprioception

Across all conditions, force production was observed to be more accurate in flexion compared to extension. Such difference between extension and flexion movement accuracy can be explained by the fact that participants generated higher MVC contractions in trunk extension than in flexion, leading to higher target forces in extension during the force reproduction protocol (more than 2 times higher). In line with this observation, results of the current study also showed differences between force accuracy at 50% and 75% of MVC for both trunk flexion and extension tasks. Participants were more accurate during the execution of force reproduction task at 50% of MVC than to during those performed at 75% of MVC. These results taken together suggest that force variability increases as the target force increases. It has been previously shown that, force variability increases linearly with force at moderate levels of force(Sherwood and A. Schmidt 1980).

262 DOMS and Trunk Proprioception

Trunk proprioception was altered under the influence of lumbar muscle DOMS, with the observation of higher AE and CE values in trunk extension in comparison to flexion. These findings suggest that DOMS had a direct impact on the proprioception of muscles that have undergone eccentric contractions, while the proprioception of the unaffected muscles (trunk flexors) remained unchanged. Even if not directly assessed in the current study, these observations support the recent views regarding the important contribution of peripheral sensory information in the production of force (Luu et al. 2011; Scotland et al. 2014) and expand it to axial muscles. A recent study showed that, in healthy individuals under the influence of experimental low back pain triggered by a combination of DOMS and hypertonic saline solution, the increase in trunk extensor muscle activity was not accompanied by an increase in trunk flexor muscle activity during postural perturbations (Larsen et al. 2017). These previous results along with those of the present study suggest that minimal or no change in the control of the trunk flexor muscles are necessary to achieve a desired motor outcome, such as trunk proprioception or postural control (Bartlett et al. 2007).

However, VE variable was not affected by lumbar muscle DOMS. Such lack of DOMS effect on VE could partially be explained by the participants' overall level of physical activity. Participants were considered moderately (at least 600 MET-min/week) to highly (3000 MET-min/week) active with a mean score of 2.7 on the short form of the IPAQ questionnaire. It has been proposed that a higher level of motor variability that is functionally related to the task is present in individuals that are physically active which could favour motor performance while limiting the occurrence of muscle fatigue (Bartlett et al. 2007; Robins et al. 2006). It can be hypothesized that under the influence of lumbar muscle DOMS, participants were able to find a new strategy, such as variation in muscle activity, to perform the desired task. This should be addressed in future research to better understand the effect of DOMS in the lumbar region. Future studies should also consider exploring the relation between trunk proprioception under DOMSeffects and physical activity by including individuals from each active group (sedentary to very active).

287 Difference and Similarities Between DOMS and Low Back Pain

As expected, the present results showed that the DOMS protocol induced experimental low back pain and soreness, which is consistent with previous studies (Abboud et al. 2019; Cheung et al. 2003). The presence of DOMS was confirmed by lumbar pain and soreness values of 2.8 and 3.8 respectively, which represent mild pain intensity and moderate soreness intensity.

DOMS has been used as a relevant pain model, which is able to reproduce alteration usually present in patients with chronic or recurrent low back pain (Abboud et al. 2019; Bishop et al. 2011). As shown in the present study, mild to moderate pain and soreness in addition to the decrease of proprioception are features of DOMS, which are similar to characteristics also found in chronic or recurrent patients with low back pain. DOMS has been associated with a decrease of muscle strength, muscle power and range of motion due to micro muscle damage (Cheung et al. 2003; Mizumura and Taguchi 2016). These muscle damages can create temporary muscles dysfunctions and perceptions (location in the space and/or strength) and, in the same way, affect performance (Larsen et al. 2017; Paschalis et al. 2007; Pearcey et al. 2015) such as precision of movement (decrease of joint range of motion) and proprioception (Vila-Chã et al. 2011), which can affect the recruitment patterns (Larsen et al. 2017; Pearcey et al. 2015; Vila-Chã et al. 2011). Alteration in trunk proprioception, such as increase of errors in reproduction force task, has been also observed in patients with chronic low back pain (Descarreaux et al. 2007).

In the present study, a significant difference in TPF for both flexion and extension tasks were observed between the pre- and post-DOMS protocol. It was recently reported that movements associated with pain are performed faster compared to movements without pain (Karos et al. 2017). Even if participants were statistically faster in pre-DOMS condition, it should be noted that differences between TPF pre- and post-DOMS protocol varied from 30 to 60 milliseconds in flexion and varied from 60 to 70

milliseconds in extension. In a previous study, participants had to reproduce 50% and 75% of their MVC in flexion and in extension in an isometric condition with their eyes closed. The authors showed that patients with chronic low back pain took ~ 120 milliseconds longer than the healthy group to reach the force target (Descarreaux et al. 2007). Therefore, it remains unknown if changes in TPF should be considered as relevant functional changes for patients with chronic or recurrent low back pain as healthy participants post-DOMS protocol. Theories of short-term pain adaptations propose that changes in the motor system are related to a protection mechanism, while in the long-term this adaptive behaviour may lead to further problems (Hodges and Tucker 2011; van Dieen et al. 2017). Another explanation for the difference between DOMS and clinical low back pain effects on TPF is the fact that participants in the current study were moderately to highly active, while the group of patients with chronic low back pain in Descarreaux et al., (2007) study were considered as moderately disabled.

Limitations and Future Recommendations

Participants were mostly young adults moderately to highly physically active, which could have minimized the effect of DOMS. However, they reported levels of back muscle pain and soreness similar to other studies using similar protocol to induce lumbar muscle DOMS (Abboud et al. 2019; Hjortskov et al. 2005), which suggest the occurrence of DOMS in the lumbar muscles. Having a small group of participants with similar characteristics can limit the generalization which may lead to an overestimation of the current results. However, most of the differences of the current results were highly significant (p=0.02 to p<0.001). Adaptations in the muscle recruitment strategy could have occurred under the influence of DOMS to perform the task, as observed in patients with chronic low back pain (Abboud et al. 2019; Falla et al. 2014). In addition, even if there is a rest time between force reproduction task and that force reproduction tasks were randomized, it was impossible to ensure that participants did not have residual fatigue during the experimentation. Future studies should assess lumbar muscle recruitment strategies under the influence of lumbar muscle DOMS to confirm this theory. Future studies also should assess sex-comparison to evaluate if there is differences in strategies used during force reproduction tasks.

1		
2		
4 5	335	
6 7	336	Conclusion
8 9	337	Lumbar muscle DOMS affect lumbar muscles proprioception during force reproduction tasks
10 11 12	338	especially in extension at higher level of force, while this performance was unchanged in trunk flexion.
13 14	339	This study suggests that lumbar muscles proprioception in lumbar muscles has been altered in muscles
15 16	340	that have been directly affected by the DOMS effects, supporting the important contribution of the
17 18 19	341	peripheral sensory systems in force reproduction. DOMS represent a relevant pain model to better
20 21	342	understand function alterations and pain mechanisms present in complex anatomical systems such as the
22 23	343	trunk in patients with chronic and recurrent low back pain.
24 25 26	344	
20 27 28	345	
29		
30 31		
32		
33		
34		
35 36		
37		
38		
39		
40		
41 42		
43		
44		
45		
46 47		
48		
49		
50		
51 52		
53		
54		
55		
56 57		
58		
59		
60		
61 62		
0∠ 63		15
64		
65		

1		
2		
3 4	246	Deferrences
5	346	References
6	347	Abboud J, Lessard A, Piché M, Descarreaux M (2019) Paraspinal muscle function and pain sensitivity
7	348	following exercise-induced delayed-onset muscle soreness European Journal of Applied
8 9	349	Physiology doi:10.1007/s00421-019-04117-6
10	350	Abboud J, Rousseau B, Descarreaux M (2018) Trunk proprioception adaptations to creep deformation
11	351	Eur J Appl Physiol 118:133-142 doi:10.1007/s00421-017-3754-2
12	352	Bartlett R, Wheat J, Robins M (2007) Is movement variability important for sports biomechanists?
13 17	353	Sports Biomech 6:224-243 doi:10.1080/14763140701322994
15	354	Bishop MD, Horn ME, George SZ, Robinson ME (2011) Self-reported pain and disability outcomes from
16	355	an endogenous model of muscular back pain BMC musculoskeletal disorders 12:35
17	356	Boucher JA, Abboud J, Nougarou F, Normand MC, Descarreaux M (2015) The Effects of Vibration and
18	357	Muscle Fatigue on Trunk Sensorimotor Control in Low Back Pain Patients PLoS One
19 20	358	10:e0135838 doi:10.1371/journal.pone.0135838
20	359	Cheung K, Hume P, Maxwell L (2003) Delayed onset muscle soreness : treatment strategies and
22	360	performance factors Sports Med 33:145-164
23	361	Cleak MJ, Eston RG (1992) Delayed onset muscle soreness: Mechanisms and management Journal of
24	362	Sports Sciences 10:325-341 doi:10.1080/02640419208729932
25 26	363	Coudreuse JM, Dupont P, Nicol C (2004) [Delayed post effort muscle soreness] Ann Readapt Med Phys
27	364	47:290-298 doi:10.1016/j.annrmp.2004.05.012
28	365	Craig CL et al. (2003) International physical activity questionnaire: 12-country reliability and validity
29	366	Med Sci Sports Exerc 35:1381-1395 doi:10.1249/01.Mss.0000078924.61453.Fb
30	367	da Silva RA, Lariviere C, Arsenault AB, Nadeau S, Plamondon A (2009) Pelvic stabilization and
31 32	368	semisitting position increase the specificity of back exercises Med Sci Sports Exerc 41:435-443
33	369	doi:10.1249/MSS.0b013e318188446a
34	370	della Volpe R, Popa T, Ginanneschi F, Spidalieri R, Mazzocchio R, Rossi A (2006) Changes in
35	3/1	coordination of postural control during dynamic stance in chronic low back pain patients Gait
36 37	372	Posture 24:349-355 doi:10.1016/J.gaitpost.2005.10.009
38	3/3	Descarreaux M, Lalonde C, Normand MC (2007) isometric force parameters and trunk muscle
39	374 275	recruitment strategies in a population with low back pain Journal of manipulative and
40	375	physiological inerapeutics 50:91-97
41	570 777	distribution of activity across back muscle regions in individuals with low back pain Dain
42 43	577 270	155-044 952 doi:10 1016/i poin 2014 01 027
44	270	Earias Junior LE at al. (2019) Bsychological responses, muscle damage, inflammation, and delayed
45	380	onset muscle soreness to high-intensity interval and moderate-intensity continuous exercise in
46	381	overweight men Physiology & hehavior 199-200-209
47	382	Hagert F (2010) Proprioception of the wrist joint: a review of current concepts and possible
40 49	383	implications on the rehabilitation of the wrist Journal of Hand Therapy 23:2-17
50	384	Hiortskov N. Essendron M. Skotte I. Fallentin N (2005) The effect of delayed-onset muscle soreness on
51	385	stretch reflexes in human low back muscles Scandinavian journal of medicine & science in
52	386	sports 15:409-415
53	387	Hodges PW, Danneels I. (2019) Changes in Structure and Function of the Back Muscles in Low Back
54 55	388	Pain: Different Time Points, Observations, and Mechanisms Orthon Sports Phys Ther 49:464-
56	389	476 doi:10.2519/iospt.2019.8827
57	390	Hodges PW, Richardson CA (1999) Altered trunk muscle recruitment in people with low back pain with
58	391	upper limb movement at different speeds Arch Phys Med Rehabil 80:1005-1012
59 60	392	doi:10.1016/s0003-9993(99)90052-7
61		
62		16
63		10
64		
65		

1 2		
3		
4 5 6	393 394	Hodges PW, Tucker K (2011) Moving differently in pain: a new theory to explain the adaptation to pain Pain 152:S90-98 doi:10.1016/j.pain.2010.10.020
ю 7 8	395 396	Jerosch J, Prymka M (1996) Proprioception and joint stability Knee Surg Sports Traumatol Arthrosc 4.171-179
9	397	Karos K. Meulders A. Gatzounis R. Seelen HAM. Geers RPG. Vlaeven JWS (2017) Fear of pain changes
10 11	398 200	movement: Motor behaviour following the acquisition of pain-related fear Eur J Pain 21:1432-
12	399	1442 doi:10.1002/ejp.1044
13 14	400	back muscle roman chair exercise in healthy and back pain subjects Med Sci Sports Exerc
15 16	402	43:157-164 doi:10.1249/MSS.0b013e3181e96388
17	403	Larsen LH, Hirata RP, Graven-Nielsen T (2017) Pain-evoked trunk muscle activity changes during fatigue
18	404	and DOMS Eur J Pain 21:907-917 doi:10.1002/ejp.993
19	405	Lee AS, Cholewicki J, Reeves NP, Zazulak BT, Mysliwiec LW (2010) Comparison of trunk proprioception
20	406	between patients with low back pain and healthy controls Archives of physical medicine and
21	407	rehabilitation 91:1327-1331
22	408	Lee PH, Macfarlane DJ, Lam T, Stewart SM (2011) Validity of the international physical activity
24	409	questionnaire short form (IPAQ-SF): A systematic review International Journal of Behavioral
25	410	Nutrition and Physical Activity 8:115
26	411	Luu BL, Day BL, Cole JD, Fitzpatrick RC (2011) The fusimotor and reafferent origin of the sense of force
27	412	and weight J Physiol 589:3135-3147 doi:10.1113/jphysiol.2011.208447
28	413	MacIntyre DL, Reid WD, McKenzie DC (1995) Delayed muscle soreness. The inflammatory response to
29 30	414	muscle injury and its clinical implications Sports Med 20:24-40
31	415	McNair PJ, Heine PJ (1999) Trunk proprioception: enhancement through lumbar bracing Arch Phys
32	416	
33	41/	Mizumura K, Taguchi T (2016) Delayed onset muscle soreness: Involvement of neurotrophic factors
34	418	The Journal of Physiological Sciences 66:43-52 doi:10.100//s125/6-015-039/-0
30 36	419	Parreira RB, Amorim CF, Gil AW, Teixeira DC, Bilodeau M, da Silva RA (2013) Effect of trunk extensor
37	420	fatigue on the postural balance of elderly and young adults during unipodal task Eur J Appl
38	421	Physiol 113:1989-1996 doi:10.100//s00421-013-2627-6
39	422	Paschalis V, Giakas G, Baltzopoulos V, Jamurtas AZ, Theoharis V, Kotzamanidis C, Koutedakis Y (2007)
40	423	The effects of muscle damage following eccentric exercise on gait biomechanics Gait & posture
41 42	424	
43	425	Pearcey GE, Bradbury-Squires DJ, Kawamoto J-E, Drinkwater EJ, Benm DG, Button DC (2015) Foam
44	426	rolling for delayed-onset muscle soreness and recovery of dynamic performance measures
45	427	Journal of athletic training 50:5-13
46	428	Proske U, Weerakkody NS, Percival P, Morgan DL, Gregory JE, Canny BJ (2003) Force-matching errors
4'/	429	after eccentric exercise attributed to muscle soreness Clin Exp Pharmacol Physiol 30:576-579
40 49	430	Rausch Osthoff AK, Ernst MJ, Rast FM, Mauz D, Graf ES, Kool J, Bauer CM (2015) Measuring lumbar
50	431	reposition accuracy in patients with unspecific low back pain: systematic review and meta-
51	432	analysis Spine (Phila Pa 1976) 40:E97-E111 doi:10.1097/brs.000000000000677
52	433	Robins R, Wheat J, Irwin G, Bartlett R (2006) The effect of shooting distance on movement variability in
53	434	basketball Journal of Human Movement Studies 50:217-238
54 55	435	Scotland S, Adamo DE, Martin BJ (2014) Sense of effort revisited: Relative contributions of sensory
56	436	Teedback and efferent copy Neuroscience letters 561:208-212
57	437	Snerwood D, A. Schmidt R (1980) The Relationship Between Force and Force Variability in Minimal and
58	438	Near-Maximal Static and Dynamic Contractions vol 12. doi:10.1080/00222895.1980.10735208
59		
60 61		
o⊥ 62		
63		17
64		
65		

1 2		
3 4 5 6 7 8 9 10 11	439 440 441 442 443 444 445	 Tong MH, Mousavi SJ, Kiers H, Ferreira P, Refshauge K, van Dieen J (2017) Is There a Relationship Between Lumbar Proprioception and Low Back Pain? A Systematic Review With Meta-Analysis Arch Phys Med Rehabil 98:120-136 e122 doi:10.1016/j.apmr.2016.05.016 van Dieen JH, Flor H, Hodges PW (2017) Low-Back Pain Patients Learn to Adapt Motor Behavior With Adverse Secondary Consequences Exerc Sport Sci Rev 45:223-229 doi:10.1249/jes.00000000000121 Vila-Chã C, Riis S, Lund D, Møller A, Farina D, Falla D (2011) Effect of unaccustomed eccentric exercise
12 13 14	446 447	on proprioception of the knee in weight and non-weight bearing tasks Journal of Electromyography and Kinesiology 21:141-147
15 16	448 449	Weerakkody NS, Whitehead NP, Canny BJ, Gregory JE, Proske U (2001) Large-fiber mechanoreceptors contribute to muscle soreness after eccentric exercise The journal of pain 2:209-219
17 18	450	
20 22 22 22 22 22 22 22 22 22 22 22 22 2		
56 57		
58 59 60		
61 62		18
ьз 64 65		

Table 1: Participant's results on clinical and physical outcomes

	Outcomes	Experimental Group
		(n=20)
		$Mean \pm SD$
Demographics	Age (years)	25.5 ± 5.2
<u> </u>	F: M	10:10
	Weight (kg)	69.6 ± 14.6
	Height (m)	1.7 ± 0.1
	BMI (kg/m^2)	23.3 ± 2.7
	IPAQ-SF	2.7 ± 0.5
Pain	Peak intensity (/10)	2.75 ± 2.27
	Days with pain	1.65 ± 1.27
Soreness	Peak intensity (/10)	3.80 ± 2.35
	Days with soreness	2.10 ± 0.91
MCV	Extension pre-DOMS (Nm)	174.89 ± 78.12
	Extension post-DOM (Nm)	178.53 ± 89.14
	Flexion pre-DOMS (Nm)	79.60 ± 34.03
	Flexion post-DOMS (Nm)	86.18 ± 33.88

F: female, M: Male, BMI: Body Masse Index, IPAQ-SF: International Physical Activity (short-form), MVC: Maximal Voluntary Contraction, DOMS: Delayed onset muscle soreness

Table 2: Means of errors	and time to peak for	pre- and post-DOMS	in flexion and extension
	*	* *	

	Flexion		Extension		
	Pre-DOMS	Post-DOMS	Pre-DOMS	Post-DOMS	
50% force					
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	
CE (Nm)	0.78 ± 11.09	-4.92 ± 5.78	4.80 ± 7.14	9.46 ± 15.28	
AE (Nm)	8.07 ± 7.82	6.63 ± 4.13	8.15 ± 5.67	13.95 ± 12.01	
VE (Nm)	4.04 ± 2.07	4.07 ± 2.25	6.81 ± 4.49	6.05 ± 2.86	
TPF (sec)	0.54 ± 0.16	0.48 ± 0.14	0.62 ± 0.15	0.56 ± 0.10	
75% force					
CE (Nm)	2.47 ± 6.50	-2.15 ± 11.15	6.05 ± 24.20	11.97 ± 20.13	
AE (Nm)	6.23 ± 4.53	9.71 ± 6.31	18.22 ± 17.07	18.35 ± 14.49	
VE (Nm)	4.42 ± 5.53	5.15 ± 2.86	9.80 ± 7.53	9.06 ± 6.83	
TPF (sec)	0.53 ± 0.19	0.50 ± 0.20	0.62 ± 0.20	0.55 ± 0.15	

CE: constant error, AE: absolute error, VE: variable error, TPF: time to peak force, SD: standard deviation, DOMS: Delayed onset muscle soreness

Table 3: Statistical analysis for each dependent variable

	Direction	Force (F)	DOMS (Do)	Di x F	Di x Do	F x Do	Di x F x Do
	(Di)						
CE	F=10.75	F=1.32	F=0.001	F=0.003	F=6.32	F=0.22	F=0.0005
	* <i>p</i> ≤0.001	<i>p</i> =0.26	<i>p</i> =0.97	<i>p</i> =0.95	*p=0.02	<i>p</i> =0.64	<i>p</i> =0.98
	$\eta p^2 = 0.36$				$\eta p^2 = 0.25$		
AE	F=26.96	F=5.61	F=3.17	F=2.76	F=0.57	F=0.02	F=6.18
	* <i>p</i> ≤0.001	*p=0.03	<i>p</i> =0.09	<i>p</i> =0.11	<i>p</i> =0.46	p=0.88	*p=0.02
	$\eta p^2 = 0.59$	$\eta p^2 = 0.23$	-	_	-	_	$\eta p^2 = 0.25$
VE	F=23.82	F=4.92	F=0.29	F=5.00	F=1.84	F=0.18	F=0.48
	* <i>p</i> ≤0.001	*p=0.04	<i>p</i> =0.59	*p=0.04	<i>p</i> =0.19	p=0.67	<i>p</i> =0.50
	$\eta p^2 = 0.56$	$\eta p^2 = 0.21$	-	$\eta p^2 = 0.21$	-	-	•
TPF	F=8.00	F=0.004	F=11.38	F=0.22	F=0.14	F=0.52	F=0.97
	* <i>p</i> =0.01	<i>p</i> =0.95	* <i>p</i> ≤0.001	<i>p</i> =0.64	<i>p</i> =0.71	<i>p</i> =0.48	<i>p</i> =0.34
	$np^2 = 0.30$	<u>^</u>	$np^2 = 0.37$	<u>^</u>	<u>^</u>	î	<u>~</u>

 $\mu p^2 = 0.57$ CE: constant error, AE: absolute error, VE: variable error, TPF: time to peak force, SD: standard deviation, *significant *p* values based on ANOVA

Figure captions

Fig. 1. Timeline for clinical and physical outcomes.

DOMS: delayed-onset muscle soreness, MVC: maximum voluntary contraction.

Fig. 2. Position of participants on lido for trunk strength reproduction task.

Fig. 3. Example of steps of the force reproduction task.

Fig. 4. Position of participants on the 45 degrees Romain chair during the lumbar muscle DOMS protocol.

Fig. 5. Direction effect (extension vs flexion) for each dependent variable (CE, AE, VE and TPF).

CE: constant error, AE: absolute error, VE: variable error, TP: time to peak, bars indicate standard deviation.

Fig. 6. ANOVA for interaction between direction, force and DOMS for the AE.

Bars indicate standard errors.

















Figure 5



Author contribution statement

All authors have contributed substantially to the manuscript. Study conception and design (MH, JA, MD), acquisition of data (MH, CD, AL, MAM), analysis and interpretation of data (all authors), drafting the manuscript (MH, CD, JA), revising it critically for important intellectual content (all authors), and final approval of the version to be published (all authors).