


RESEARCH

Open Access



Transition of care from post-acute services for the older adults in Quebec: a pilot impact evaluation

Shashank Ghai^{1,2*} , Kathleen Chass e³, Marie-Jeanne Renaud³, Lilian Guicherd-Callin³, Andr e Bussi eres^{4,5,6} and Diana Zidarov^{7,8}

Abstract

Background Early discharge of frail older adults from post-acute care service may result in individuals' reduced functional ability to carry out activities of daily living, and social, emotional, and psychological distress. To address these shortcomings, the Montreal West Island Integrated University Health and Social Services Centre in Quebec, Canada piloted a post-acute home physiotherapy program (PAHP) to facilitate the transition of older adults from the hospital to their home. This study aimed to evaluate: (1) the implementation fidelity of the PAHP program; (2) its impact on the functional independence, physical and mental health outcomes and quality of life of older adults who underwent this program (3) its potential adverse events, and (4) to identify the physical, psychological, and mental health care needs of older adults following their discharge at home.

Methods A quasi-experimental uncontrolled design with repeated measures was conducted between April 1st, 2021 and December 31st, 2021. Implementation fidelity was assessed using three process indicators: delay between referral to and receipt of the PAHP program, frequency of PAHP interventions per week and program duration in weeks. A battery of functional outcome measures, including the Functional Independence Measure (FIM) and the Patient-Reported Outcomes Measurement Information System (PROMIS) Global-10 scale, as well as fall incidence, emergency visits, and hospitalizations were used to assess program impact and adverse events. The Patient's Global Impression of Change (PGICS) was used to determine changes in participants' perceptions of their level of improvement/deterioration. In addition, the Camberwell Assessment of Need for the Elderly (CANE) questionnaire was administered to determine the met and unmet needs of older adults.

Results Twenty-four individuals (aged 60.8 to 94 years) participated in the PAHP program. Implementation fidelity was low in regards with delay between referral and receipt of the program, intensity of interventions, and total program duration. Repeated measures ANOVA revealed significant improvement in FIM scores between admission and discharge from the PAHP program and between admission and the 3-month follow-up. Participants also reported meaningful improvements in PGICS scores. However, no significant differences were observed on the physical or mental health T-scores of the PROMIS Global-10 scale, in adverse events related to the PAHP program, or in the overall unmet needs.

*Correspondence:

Shashank Ghai
shashank.ghai@kau.se

Full list of author information is available at the end of the article



  The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Conclusion Findings from an initial sample undergoing a PAHP program suggest that despite a low implementation fidelity of the program, functional independence outcomes and patients' global impression of change have improved. Results will help develop a stakeholder-driven action plan to improve this program. A future study with a larger sample size is currently being planned to evaluate the overall impact of this program.

Clinical trial registration Retrospectively registered NCT05915156 (22/06/2023).

Keywords Post-acute care, Multimorbidity, Geriatric care, Rehabilitation

Background

The Canadian healthcare system is facing tremendous challenges as a result of an aging population [1]. While approximately 0.5% of Canadians are between the age of 50 to 74 years, and 2.6% are 75 years or older, these individuals account for nearly half of all hospital days (45.6% and 56.1%, respectively) [2]. In addition, people with higher health care utilization tend to be hospitalized for longer periods of time (mean of two months) [2]. Hospitalization and prolonged acute care stays can lead to a functional decline, one of the most common adverse outcomes of hospitalization [3, 4]. Furthermore, patients often experience prolonged non-medical stays where they occupy beds while waiting for an alternative level of care, such as home care, rehabilitation, or other services [5, 6]. Importantly, post-hospitalization functional decline may be sustained up to one year following discharge, and failure to return to baseline functional status has been associated with increased risk of institutionalization [7], prolonged disability [8], and death [9, 10]. Post-acute care services are often used to address this functional decline post-hospitalization. Instead of being discharged from acute care to home or directly to a nursing home, many seniors first transition to post-acute care to receive specialized inpatient and/or outpatient rehabilitation to regain functional autonomy [11].

Older adults with multimorbidity have substantial health and social needs after discharge from post-acute care facilities [12], such as maintaining or regaining the ability to perform activities of daily living [13]. Patients have also emphasized a strong desire to remain at home with appropriate support rather than being institutionalized [12, 14]. Similarly, social isolation and emotional distress are also widely reported upon discharge from post-acute care, highlighting the need for social support interventions [12, 15]. Support is also needed for personal care tasks, which include assistance with activities of daily living (e.g. bathing or showering) and with instrumental activities of daily living (e.g. light housework), assistance with medication management, or use of assistive devices [16]. To better meet these needs, health care professionals and caregivers highlighted the importance of effective coordination of services and supports (e.g., a case coordinator) and clear care pathways [17].

In Quebec's conventional care model, particularly within the Montreal West Island Integrated University Health and Social Services Center (MWIIUHSSC), the patient trajectory includes admission to an acute care hospital, then depending on the patient's needs, the person may be discharged home or can be transferred to internal post-acute rehabilitation services or intensive functional rehabilitation (RFI). After discharge from these services (acute or post-acute), patients may be referred to external outpatient rehabilitation services to improve functional ambulation and community integration. These publicly funded outpatient services include adult ambulatory rehabilitation (SARCA), home care (SAD), day-hospital care, and community reintegration programs. Despite the availability of these models of care, access to these rehabilitation services in Canada remains a challenge [18]. For example, at Local Community Service Centers (LCSCs) in the MWIIUHSSC's territory, individuals often face long waits of up to 119 days to access outpatient physiotherapy services after discharge from post-acute care. Similarly, a study by Deslauriers, Raymond [19] found a mean waiting time of 7.6 months (range from 0 to 77 months) for outpatient physiotherapy services in publicly funded hospitals in the province of Quebec (Canada). This delay in access to external rehabilitation services is exacerbated by a persistent shortage of healthcare professionals and an increasing demand for services for an aging population [20, 21]. With the goal of optimizing post-acute care physical functioning and functional independence among older adults admitted to acute care or their rehabilitation facilities, the MWIIUHSSC piloted the Post-Acute Home Physiotherapy Program (PAHP). This program was designed as a new clinical care pathway in accordance with the *Cadre de référence Montréalais* [22], and adapted from the Institut national d'excellence en santé et services sociaux (INESSS) guidelines to address the rehabilitation needs of clients and their families by providing center-based or home-based outpatient rehabilitation services and early supported discharge to improve access and continuity of rehabilitation care [23]. The PAHP program was also intended to accelerate the turnover of acute care and rehabilitation beds.

The PAHP program was implemented in November 2018 in the four LCSCs in the MWIIUHSSC's territory.

Although promising, the impact of the PAHP program and whether it addresses the healthcare needs of older people remained unknown for three years post-implementation. Therefore, the current study aimed to assess: 1) the implementation fidelity of the program, which refers to the degree to which the program was delivered as intended [24]; the impact of the PAHP program on the functional independence, physical and mental health outcomes and quality of life of older adults who underwent this program 3) its potential adverse events (incidence of falls, rehospitalizations, and medical consultations); and 4) to identify the physical, psychological, and mental health care needs of older adults after their discharge to home.

Methods

Design

This quasi-experimental uncontrolled study with three measurement periods was carried out between April 1st, 2021 and December 31, 2021. This study was approved by the Research Ethics Committee of the MWI-IUHSSC (IRB number: SMHC-20-21). The study was retrospectively registered on June 22, 2023 on clinicaltrials.gov (registration number: NCT05915156). All patients signed an informed consent form. A Transparent Reporting of Evaluations with Nonrandomized Designs (TREND) checklist and a Template for Intervention Description and Replication (TIDieR) checklist are provided as Additional Files 1 and 2, respectively. The purpose of using the TREND checklist was to ensure transparent reporting of the current study to improve research synthesis and facilitate evidence-based recommendations [25]. Similarly, the TIDieR checklist was used to provide a structured account of the PAHP intervention evaluated in this study [26].

Participants and recruitment

Potential participants were first approached by a member of their inpatient rehabilitation team to seek their interest in participating in the study, before being contacted by one of the project leaders (KC, MJR). Patients were eligible if they were discharged home with or without home-care services and could understand French or English. Patients were excluded if they had an unstable medical, psychiatric, and postoperative condition or a severe cognitive impairment (subjective assessment of the rehabilitation team).

PAHP program (intervention)

The PAHP program was designed to provide physical rehabilitation services to patients recently discharged from acute or post-acute care. In this study, the term “discharged from acute or post-acute care”, specifically refers to individuals who received in-patient rehabilitation

services. As part of the PAHP program planning, each patient was to be treated by one of the four physiotherapy technologists assigned to the PAHP program working at the four LCSCs in the MWIIUHSSC’s territory within 48 h of discharge from an acute or post-acute care facility, and to receive home physiotherapy at least three days a week for up to six weeks. Each session lasted between 45 and 60 min, and the same physiotherapy technologist provided care to the patients in their caseload from the admission to the discharge of the patient from the program. The profession of a physiotherapy technologist is specific to the province of Quebec [27]. These professionals become involved in the patient’s care after the patient’s initial assessment by a physiotherapist or a physician. Their responsibilities include analyzing information from the patient’s medical record and using this data to develop, implement, and monitor individualized treatment plans. Physiotherapy interventions provided through the PAHP were tailored by the physiotherapy technologists based to the reasons for the referral and the patient’s needs. For instance, if a patient was unable to climb or descend stairs independently after discharge from acute or post-acute care, the PAHP therapist focused on building lower extremity strength and providing home stair training. In addition, when patients were referred to the PAHP program, the referring therapist sent goals to work on, and the physiotherapy technologist continued home rehabilitation to achieve those goals. The physiotherapy technologist was also independent in evaluating and determining the specific exercises/treatments needed to achieve these goals. In addition, because of their caseloads or other organizational issues (e.g., staff shortages, other responsibilities, etc.) the physiotherapy technologists were not necessarily able to provide the 3 weekly sessions required by the PAHP to all the older adults and had to extend the duration of the program beyond the required 8 weeks in order to meet the patient’s needs and achieve the established functional goals.

Measuring instruments

Sociodemographics of all study participants enrolled in the PAHP were collected from the participant’s medical record and interview, including age, gender, education, marital status, principal comorbidity, admission diagnosis to post-acute care, type of living arrangement, presence of caregiver/spouse at home, and length of time in the PAHP program. The primary outcome measures in this study are consistent with the goals of the PAHP program and the recommendations of the International Consortium for Health Outcomes Measurement (ICHOM) for core outcome measures to be used in older adults [28]. The two physiotherapists who performed the

outcome assessments were different from the physiotherapy technologists who administered the PAHP program.

Implementation fidelity

Implementation fidelity of the intervention was assessed using three process indicators: 1) delay between referral and the receipt of the PAHP program (24 to 48 h was targeted); 2) frequency of PAHP visits per week (a minimum of 3 interventions/week was targeted) and 3) duration of the PAHP program in weeks (4 to 6 weeks was targeted).

Impact of PAHP program

1. The Functional Independence Measure (FIM) assessed the individual's functional ability. The FIM measures independent performance in the areas of self-care, sphincter control, transfers, locomotion, communication, and social cognition. The FIM instrument consists of 18 items, and each item score ranges from 1 to 7 (a score of 7 is categorized as "complete independence", while a score of 1 represents "total assistance"). The total score ranges from 18 (the lowest) to 126 (higher scores on the FIM indicate patients with a greater level of independence and less need for assistance). Thirteen of the 18 FIM items are from the motor subscale, and the remaining five items are from the cognitive subscale [29]. It has been shown that the total FIM scores can be treated as interval scores [30].
2. The Patient-Reported Outcomes Measurement Information System (PROMIS) Global-10 (10-item) survey assesses the self-reported perceptions of health, physical, mental and social health, and the quality of life using a 5-point Likert scale. The scale is reliable, precise and comparable to legacy instruments [31]. The PROMIS-10 physical and mental health scores are converted into standardized T-scores [32]. A T-score of 50 is identified as the mean score for the general population, with a higher or a lower score indicating better or worse physical and mental health, respectively.
3. Patient Global Impression of Change Scale (PGICS) (1 item) assessed participants' perceptions of their level of improvement or deterioration in their physical function at discharge from the PAHP program. An 11-point Likert scale was used with options ranging from "very much worse" to "completely recovered". The minimum clinically important change is 1.35 points on this 11-point scale [33]. The scale has been previously validated in the context of chronic pain rehabilitation [34, 35].

Potential adverse events of PAHP program

Adverse events related to the PAHP program were assessed using fall events at two time points: first, from the time participants were discharged from acute or post-acute care until the end of the PAHP program, and second, at the 3-month follow-up after discharge from the PAHP program. To further evaluate the adverse events associated with the program, patient self-reports of the number of emergency visits and hospital readmissions during these time periods were also analyzed.

Needs assessment

The Camberwell Assessment for the Need of the Elderly (CANE) was used to assess the met and unmet physical, psychological, social and environmental care needs (24 domains) from the perspective of the older adults [36]. The CANE has previously been used and validated with older adults living in the community [37–41], and with individuals discharged from acute psychiatric care [42]. For each domain, there is a question about a specific need. Responses are scored on a three-point scale, with 0 indicating no need, 1 indicating met need (problem received appropriate intervention), 2 indicating unmet need (problem left without optimal intervention), and 9 indicating unknown if the patient did not know the type of assistance received.

Data collection procedure

Figure 1 shows the data collection procedure for all the outcomes. The two project leaders, physical therapists by training, collected all outcome measures through face-to-face or telephone interview with participants. They were all trained to administer the FIM and the CANE using the CANE manual [43].

Data analysis

Descriptive statistics were calculated for participants' demographic and clinical characteristics, and questionnaire scores using means and standard deviations (continuous variables) and frequencies and percentages (categorical variables). Descriptive statistics were also used to compare implementation fidelity and outcomes related to adverse events (i.e., falls and medical visits related to the primary diagnosis) between two time points: events occurring from acute/post-acute care discharge to PAHP discharge and after PAHP discharge at the 3-month follow-up. We evaluated the impact of the PAHP program on FIM scores and PROMIS-10 (i.e., separately on T-score physical, T-score mental, general health, social activities and role) over time and whether the changes persisted during the 3-month follow-up, using a one-way repeated measures ANOVA with the three time points (i.e., hospital discharge, PAHP

	Acute/post-acute care discharge	PAHP admission	PAHP discharge	3 months PAHP post-discharge
Primary outcome measure				
Functional Independence Measure (FIM)	X		X	X
Secondary clinical outcomes measures				
PROMIS Global-10		X	X	X
Patient Global Impression of Change Scale (PGICS)			X	
Adverse events				
- Fall				
- Emergency visit			X	
- Re-hospitalizations				X
Need assessment tool				
Camberwell Assessment for the Need of the Elderly (CANE)		X	X	

Fig. 1 Time periods of data collections

discharge, three-month follow-up) as the within-subject factor. When significant effects were observed, post hoc comparisons with Bonferroni correction were performed to determine whether group differences emerged over time. Independent samples t-tests were conducted to evaluate the influence of the PAHP program on needs assessment, including the met and unmet needs as assessed by the CANE questionnaire, across two time points (i.e., PAHP admission and PAHP discharge). The statistical significance level was set at 5%, and all analyses were performed using IBM SPSS Version 28.0 (Armonk, NY: IBM Corp).

Results

Demographic and clinical variables

During the study period, 54 patients were referred to the PAHP program of whom 12 declined to participate, nine were ineligible due to cognitive impairment or language barriers, and nine were lost due to administrative issues, leaving 24 (44.4%) study participants (Fig. 2). Data for the three evaluation times were available for 24 participants (50% female, mean age was 80.6 ± 7.9 years). The primary reason for the admission to acute or post-acute care for the patients was musculoskeletal disorders (66.7%), followed by infectious diseases (12.5%), cardio-pulmonary disorders (12.5%), neurological disorders (4.2%), and other (4.2%) (Table 1).

Implementation fidelity

Out of the 664 post-acute care users, 54 older adults (8.1%) were referred to the PAHP program. Among these, 24 individuals (44.4%) actively participated in the intervention. The intervention was delivered a mean

of 12.5 ± 10 days (ranging from 1 to 48 days) after referral. The PAHP intervention consisted of an average of 1.6 ± 0.72 sessions per week, lasting for a total of 7.85 ± 5 weeks. Patient medical information was available for all participants prior to the intervention. The physiotherapy technologist planned a total of 10.2 ± 2.6 sessions for the participants, while the actual sessions delivered were 10.8 ± 6.6 .

Functional independence measure

The mean (SD) FIM scores at hospital discharge, PAHP discharge, and follow-up were 105.8 (17.5), 111.2 (13.6) and 111.3 (15.5), respectively (Table 2). The results indicate a significant effect of time of measurement on the FIM scores ($F_{1,4,30.8}=7.58$, $p=0.005$). Post-hoc comparison with Bonferroni corrections suggests that FIM had improved between PAHP discharge and hospital discharge ($p=0.005$) and between the follow-up and hospital discharge ($p=0.04$).

PROMIS-10 global health

The mean PROMIS-10 Global Health T-scores for physical health at admission, discharge, and follow-up were close to normal 42.6, 43.3, and 44, respectively (Table 2). Similarly, the T-scores for mental health at admission, discharge, and follow-up were also close to normal at 47.6, 46.9, and 48, respectively. There were no significant differences between the admission, discharge, and the follow-up T-scores for mental ($F_{2,44}=0.09$, $p=0.9$) and physical ($F_{2,44}=0.48$, $p=0.61$) health. For the two subscales that were not included in the T-scores, no significant differences were observed for either the general health subscale ($F_{2,44}=0.52$, $p=0.60$) or the social activities and roles subscale ($F_{2,44}=0.13$, $p=0.87$).




Fig. 2 Schematic representation of the included participants

Patient Global Impression of Change score.

The average change in the PGICS for older adults participating in the PAHP program was 1.95 ± 1.3 .

Adverse events.

The number of falls decreased from 19 at post-acute care discharge to nine at the 3-month follow up (Table 2). Similarly, the number of individuals who experienced falls decreased from 11 at PAHP discharge to five at the 3-month follow-up. There was also a reduction in the number of emergency care visits related to principal diagnosis. Here, the number of individuals who had emergency care visits at PAHP discharge was five, which was reduced to three at the 3-month follow up. However, the

number of individuals who had a hospitalization related to the primary diagnosis increased slightly at 3 months after PAHP discharge compared to PAHP discharge (2 to 3 individuals).

Camberwell assessment for needs in elderly

The mean met and unmet needs at the time of PAHP admission and discharge were similar (Table 3). Independent samples t-tests revealed no significant effect of time of measurement on the overall unmet ($t_{46} = -0.27$, $p = 0.78$) and met ($t_{46} = 0.37$, $p = 0.70$) needs. Regarding specific needs related to the environment, older adults undergoing PAHP had a decrease in unmet needs in areas such

Table 1 Demographic and clinical characteristics of participants

Characteristics	Mean (standard deviation), or N (%)
Total participants	24
Gender	
Female	12 (50%)
Male	12 (50%)
Age (years)	80.6 ± 7.9 (range: 60.8–94)
Marital status	
Single	11 (45.8%)
Married	13 (54.1%)
Admission diagnosis	
Musculoskeletal disorder	16 (66.7%)
Infectious disease	3 (12.5%)
Cardiopulmonary disorder	3 (12.5%)
Neurological disorder	1 (4.2%)
Other	1 (4.2%)
Principal comorbidity	
Cardiopulmonary disorder	13 (54.2%)
Musculoskeletal disorder	5 (20.8%)
Infectious disease	1 (4.2%)
Neurologic disorder	1 (4.2%)
Other	4 (16.7%)
Homecare services after discharge	
With services (home care)	12 (50%)
Without services	12 (50%)
Caregiver at home	
Yes	15 (62.5%)
No	9 (37.5%)
Length of PAHP program (weeks)	7.8 ± 4.9 (range: 0.57–21.4 weeks)

as accommodation (% admission vs. discharge: 8.3% vs. 4.2%), looking after home (12.5% vs. 8.3%), and caring for someone (8.3% vs. 4.2%). In addition, decreases in unmet social needs related to daytime activities (20.8% vs. 16.7%) and company (12.5% vs. 8.3%) and physical needs related to self-care (12.5% vs. 0%) were also reported by older adults in the PAHP program. In contrast, increases in unmet physical needs were observed for mobility (16.7% vs. 20.8%), physical health (4.2% vs. 12.5%), eyesight/hearing (0% vs. 8.3%) and medication management (0% vs. 8.3%). Other areas where unmet needs have increased during the PAHP program were food-related needs (environment, 8.3% vs. 12.5%) and memory (0% vs. 8.3%) and psychological distress (psychological needs, 8.3% vs. 12.5%).

Discussion

The MWIIUHSSC piloted the PAHP program which reduced waiting times for home physiotherapy rehabilitation to improve the autonomy of older adults transitioning from acute or post-acute care to home. Individuals in the PAHP program showed a significant improvement in functional independence and gains in functional

Table 2 Descriptive and statistical analysis of outcome measures

Outcome	Hospital-Discharge	PAHP-Admission	PAHP-Discharge	PAHP-follow up	p-value
FIM (±SD)	105.8 ± 17.5	-	111.2 ± 13.6	111.3 ± 15.5	0.005
PROMIS-10 Global Health					
Physical health					
T-score (±SE)	-	42.6 ± 5.8	43.3 ± 5.1	44 ± 7.5	0.61
Mental health					
T-score (±SE)	-	47.6 ± 9.9	46.9 ± 10.0	48.0 ± 8.7	0.91
General health (±SD)	-	3.1 ± 0.8	2.8 ± 0.7	3.1 ± 0.8	0.59
Social activities & role (±SD)	-	3.1 ± 1.1	3.0 ± 1.1	3.0 ± 1.1	0.87
PGICS (±SD)	-	-	1.95 ± 1.3	-	-
Adverse events					
Falls (N)	-	-	19	9	
Number of individuals	-	-	11	5	
Emergency visits (N)**	-	-	5	3	
Number of individuals	-	-	5	3	
Hospitalizations (N)**	-	-	2	3	
Number of individuals	-	-	2	3	

PGICS: Patient global impression of change scale, ADL: Activities of daily living, CANE: Camberwell Assessment for Needs in Elderly, PROMIS: Patient Reported Outcome Measurement Information System, p-value: associated with analysis of variance, any significant ($p < 0.05$) results indicate differences in scores across the three evaluation periods, **visits in line with the principal diagnosis

independence were maintained at the 3-month follow-up. Meaningful gains in the PGICS score were also reported by patients following the PAHP program. We observed no statistical differences in the physical and mental health outcomes with the implementation of the PAHP program, nor in the overall met and unmet needs of older adults undergoing the PAHP program.

Implementation fidelity

Assessing intervention fidelity is critical for understanding how well an intervention was implemented and whether it followed its intended design [44]. In our study, the implementation of the PAHP program was not executed as planned, with the intended 48-hour visit by a physiotherapy technologist delayed by an average of 12 days. The frequency of physiotherapy technologist visits was also lower than intended, with an average of two visits per week compared to the intended three visits per

Table 3 Distribution of met and unmet needs according to Camberwell Assessment of Need for the Elderly (CANE)

CANE sections	PAHP-Admission				PAHP-Discharge			
	Met need		Unmet need		Met need		Unmet need	
	N	%	N	%	N	%	N	%
Environmental needs								
Accommodation	5	20.8	2	8.3	2	8.3	1	4.2
Looking after the home	20	83.3	3	12.5	21	87.5	2	8.3
Food	15	62.5	2	8.3	14	58.3	3	12.5
Caring for someone	3	12.5	2	8.3	2	8.3	1	4.2
Money/budgeting	5	20.8	0	0.0	5	20.8	0	0.0
Benefits	0	0.0	0	0.0	0	0.0	0	0.0
Physical needs								
Self-care	10	41.7	3	12.5	9	37.5	0	0.0
Eyesight/hearing	5	20.8	0	0.0	4	16.7	2	8.3
Mobility/falls	19	79.2	4	16.7	17	70.8	5	20.8
Incontinence	4	16.7	1	4.2	3	12.5	1	4.2
Physical health	17	70.8	1	4.2	15	62.5	3	12.5
Drugs	9	37.5	0	0.0	6	25.0	2	8.3
Psychological needs								
Psychotic symptoms	1	4.2	0	0.0	1	4.2	0	0.0
Memory	5	20.8	0	0.0	5	20.8	2	8.3
Psychological distress	0	0.0	2	8.3	1	4.2	3	12.5
Safety to self	0	0.0	0	0.0	0	0.0	0	0.0
Inadvertent self-harm	0	0.0	0	0.0	0	0.0	0	0.0
Behavior	0	0.0	0	0.0	0	0.0	0	0.0
Alcohol	2	8.3	0	0.0	1	4.2	0	0.0
Social needs								
Information	5	20.8	4	16.7	2	8.3	4	16.7
Abuse/neglect	0	0.0	0	0.0	0	0.0	0	0.0
Company	6	25.0	3	12.5	4	16.7	2	8.3
Intimate relationships	0	0.0	0	0.0	1	4.2	0	0.0
Daytime activities	6	25.0	5	20.8	8	33.3	4	16.7
Overall Mean (%)	5.7	23.8	1.3	5.6	5	21	1.4	6.1
Standard deviation	6.2		1.6		6		1.5	

week. Moreover, while the duration of the PAHP program for each participant was close to the intended eight weeks, the duration of the intervention varied between participants, ranging from half a week to 22 weeks. These deviations from the planned frequency of interventions and duration of the program could be due to the varying rehabilitation needs of the patients, which prompted the physiotherapy technologists to adjust the intensity and duration of the interventions offered according to these needs. For example, some patients may have only needed interventions to increase their independence in outdoor locomotor activities, such as endurance walking outdoors or walking on uneven terrain. Others, more deconditioned upon discharge from acute or post-acute care, may have required longer interventions to increase their functional autonomy.

However, despite the deviations from the intended fidelity, it's important to note that the PAHP program has been sustained for more than 3 years, indicating recognition of its value within the health system. Perhaps,

instead of viewing low fidelity solely as a limitation, it can be viewed as an opportunity to adapt the intervention and processes to better meet the needs and realities of the local context, i.e., the LCSCs in the MWIIUHSSC's territory. For instance, several strategies can be explored to improve fidelity and enhance the effectiveness of the PAHP program. It is critical to assess and address the factors that contribute to delays in referral and delivery of the program. This may include streamlining referral processes, improving communication among multidisciplinary staff, educating clinicians, improving scheduling efficiency, and monitoring quality of care [45, 46].

Ongoing monitoring and evaluation of fidelity, including regular feedback from clinicians and decision makers, could be used to identify areas for improvement and track the impact of changes implemented. This iterative process of adaptation and refinement could not only enhance fidelity but also contribute to a deeper understanding of the effectiveness of the program and its integration into routine practice [47, 48].

Functional independence

Limited access to outpatient rehabilitation interventions for patients admitted to the acute care setting negatively impacts length of stay and affects continuity of care [49, 50]. This study found an increase in functional independence using FIM after the PAHP program and at 3-month follow-up, which has been associated with improved quality of life [51], and decreased mortality [52]. Small changes in FIM (i.e., as little as one point) are associated with an 8% improved odds of better quality of life [51]. Therefore, these improvements in functional independence support the clinical relevance of the PAHP program, given the improved performance in motor and cognitive subsets associated with functional independence in older adults with multimorbidity. Other studies have also found an association between changes in the FIM and improved balance and mobility outcomes in older adults [53, 54].

Global impression of change

We also evaluated older adults' perceptions of improvement/deterioration in physical function following the PAHP program. We observed a clinically meaningful change in perceived improvement in physical function [55, 56]. Improvements in PGICS have been associated with reductions in pain intensity and improvements in treatment effectiveness, mood, and quality of life [57].

Met and unmet needs

Assessing met and unmet needs from the patient's perspective is considered essential to better understand whether additional components related to mobility and functional independence need to be added to the PAHP [58, 59], and to allow for effective troubleshooting to make this pilot program more appropriate to the needs of our specific population. We found no significant difference between the overall unmet needs and met needs before and after the PAHP intervention. Further exploration of our findings provide additional insights into the development of effective post-discharge services. First, with regard to environmental needs, we found that older adults had fewer unmet needs related to accommodation, looking after home, and caring for someone after the PAHP program. We also found reduced unmet needs related to daytime activities, company, and self-care. Based on the improved functional independence, it could be interpreted that independence in performing activities of daily living, including self-care, have improved as a result of PAHP and helped this population to better adapt to their accommodations.

However, we also observed an increase in unmet needs for food-related needs, mobility, physical health, eye-sight/hearing, medication management, memory, and psychological distress. To address these unmet needs, a

standardized assessment of patient needs at discharge could be incorporated into the PAHP program. This strategy could be implemented in an integrated care model with the PAHP program, which could address the unmet needs by preventing and managing declines in the intrinsic capacity and functional ability rather than treating them [60–62]. This is in line with the World Health Organization's World Report on Ageing and Health and consistent with emerging evidence on the provision of integrated care for older people with complex health needs [61, 63]. The integrated and coordinated services of the PAHP program may involve the integration of different health care professionals, such as nurses, occupational therapists, general practitioners, and social workers, in a synchronized approach to care. This could facilitate the enhancement of intrinsic capacity and address the needs that extend beyond the expertise of a physiotherapist alone. In this way, through collaborative efforts, health care providers can provide personalized and comprehensive care, prioritizing the needs of patients and moving the health care system toward the goal of integrated care for older adults [64].

Future implications

The PAHP pilot program has the potential to benefit older adults transitioning from acute or post-acute care to their homes and could be implemented in clinical practice. By integrating the program into discharge planning and rehabilitation interventions, older adults could be supported to maintain their functional independence. Likewise, policymakers could consider implementing similar programs to help alleviate the ever-increasing burden on the healthcare system due to an ageing population. Future research could explore the potential impact of such a program on reducing healthcare costs and improving patient outcomes.

The older adults in our study had a range of unmet needs that could not be addressed by physiotherapy alone, which is the only service provided by the PAHP program. Previous reviews have emphasized the importance and benefits of multidisciplinary rehabilitation teams that include nurses, physicians, physiotherapists, occupational therapists, and dietitians. Such an approach has been shown to improve mobility [65], be cost-effective [66], and reduce the risk of mortality and hospital readmissions [67, 68]. Moreover, the presence of comorbidities in this population highlights the need for coordinated care, which could be addressed through the use of a case manager [69, 70]. A post-discharge case manager can help patients achieve autonomy by helping them navigate the healthcare system and access the resources they need to manage their condition and recover at home. A case manager can act as a liaison between the patient and their healthcare team, connecting them with

appropriate specialists and community resources such as home health services or support groups. They can also provide guidance on medication management help coordinate follow-up appointments, increase the use of community services, and help patients understand their care plan and what to expect during their recovery. In the case of our study, for example, social services could be directed to individuals with identified needs related to access to food, while a team of physicians and nurses could be assigned to those with needs related to medication management and sensory impairments. Pharmacists could also play a more prominent role in post-discharge services by reviewing medication regimens or addressing issues that may affect patients' ability to obtain medications [70]. Occupational therapists could play an important role by conducting assessments of individuals and their home environments, arranging or identifying a process for obtaining and installing equipment or modifying homes to prevent falls, and then instructing the patient on its use [71]. In addition, mental health services can be targeted to those with increasing unmet needs related to psychological distress. Previous studies have shown that post-discharge case management is effective in improving clinical outcomes and reducing hospital utilization and re-admissions by improving access to needed services [70, 72].

In addition more participants need to be recruited to assess the overall impact of the program, and future research could examine the effectiveness of physiotherapist-led and interdisciplinary team case management for older adults after hospital discharge [17]. Although not evaluated in the present study, the use of physiotherapy technologist may have provided an alternative model of care that, in addition to reducing costs, could alleviate challenges such as staffing shortages and increased workload for physiotherapists due to growing service demands [73, 74]. Future studies should evaluate the efficacy and cost-effectiveness of integrating physiotherapy technologists into clinical care models to address staffing shortages and rising demands for physiotherapy services. In addition, qualitative research should explore patient and provider perspectives on this alternative care model.

Limitations

To our knowledge, this is the first study to evaluate the impact of the PAHP pilot program in Quebec. Nevertheless, our study has several limitations. First, the sample was size small which may have resulted in type I or type II errors [75]. Second, due to the COVID-19 pandemic and the highly stressed healthcare system, we faced significant challenges in recruiting and retaining participants. Thirdly, our study used a single-group design without a comparator arm. This lack of a comparator group poses challenges in attributing observed changes

solely to the PAHP intervention, as external factors may contribute to the outcomes. However, the decision to use a single-group design was guided by the pilot nature of the PAHP program, which was implemented on a smaller scale to identify potential issues and refine the program prior to broader implementation. Ultimately, our goal is to use these findings to inform the development of a stakeholder-driven action plan aimed to improve the program. Subsequently, we plan to reassess the effectiveness of the program in a larger- trial, this time with a comparator arm for a more comprehensive evaluation. Fourth, our results indicate a mixed level of fidelity in the implementation of the PAHP program. For example, we observed that delays in the program initiation exceeded the intended period, and the frequency of sessions fell short of the intended minimum frequency. These issues, in addition to unmet participant needs will be used to inform a stakeholder-driven action plan to improve this program.

Conclusion

Findings from this pilot study evaluating the impact of a new accelerated access to care program in Quebec showed improved functional independence and a positive global impression of change among older adults participating in the PAHP program. Our findings provide valuable insights for the development of a stakeholder-driven action plan to improve the program by promoting a multidisciplinary approach and the inclusion of a post-discharge case manager to coordinate healthcare services. Robust research is needed to evaluate the overall impact of this program on healthcare costs and patient outcomes.

Abbreviations

PAHP	Post-Acute Home Physiotherapy program
FIM	Functional Independence Measure
PROMIS	Patient-Reported Outcomes Measurement Information System
PGICS	Patient's Global Impression of Change
CANE	Camberwell Assessment of Need for the Elderly
ANOVA	Analysis of Variance
MWIIUHSSC	Montréal West Island Integrated University Health and Social Services Center
INESSS	Institut national d'excellence en santé et services sociaux
LCSC	Local Community Service Centers
ICHOM	International Consortium for Health Outcomes Measurement

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12913-024-10818-2>.

Supplementary Material 1

Supplementary Material 2

Acknowledgements

Not applicable.

Author contributions

DZ, KC, MJR, LGC, AB conceptualized the study and were responsible for acquiring the funding, designing, and implementing the research. SG, DZ wrote the initial version of the manuscript and performed the data analysis. All authors read and approved the final version of the manuscript.

Funding

Open access funding provided by Karlstad University. This study was funded by Edith Strauss Rehabilitation Research Projects of McGill University. Open access funding provided by Karlstad University.

Data availability

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations**Competing interests**

The authors declare no competing interests.

Ethics approval and consent to participate

All the methods in this study had been performed in accordance with the Declaration of Helsinki and had been approved by the research ethics committee of the Montréal West Island Integrated University Health and Social Services Centre (IRB number: SMHC-20-21). Informed consent was taken from participants to participate in the study.

Consent for publication

Not applicable.

Author details

¹Department of Political, Historical, Religious and Cultural Studies, Karlstad University, Karlstad, Sweden

²Centre for Societal Risk Research, Karlstad University, Karlstad, Sweden

³Montréal West Island Integrated University Health and Social Services Centre, Montreal, Québec, Canada

⁴School of Physical and Occupational Therapy, McGill University, Montreal, Québec, Canada

⁵Centre de Recherche Interdisciplinaire en Réadaptation du Montréal Métropolitain, Montréal, Québec, Canada

⁶Département Chiropratique, Université du Québec à Trois-Rivières, Trois-Rivières, Québec, Canada

⁷Faculté de Médecine, Université de Montréal, Montréal, Québec, Canada

⁸Centre de Recherche Interdisciplinaire en Réadaptation (CRIR), Institut universitaire sur la réadaptation en déficience physique de Montréal (IURDPM), Centre intégré universitaire de santé et de services sociaux du Centre-Sud-de-l'Île-de-Montréal, Montréal, Québec, Canada

Received: 11 July 2023 / Accepted: 3 March 2024

Published online: 03 April 2024

References

- Wister AV, Speechley M. Inherent tensions between population aging and health care systems: what might the Canadian health care system look like in twenty years? *J Popul Ageing*. 2015;8:227–43.
- Statistics Canada. High use of acute care hospital services at age 50 or older. In: Catalogue no. 82-003-X HR, editor. Sept 2017. p. 3–16.
- King BD. Functional decline in hospitalized elders. *Medsurg Nurs*. 2006;15(5):265–71.
- Sager MA, Rudberg MA. Functional decline Associated with hospitalization for Acute Illness. *Clin Geriatr Med*. 1998;14(4):669–80.
- Sutherland JM, Crump RT. Alternative level of care: Canada's hospital beds, the evidence and options. *Healthc Policy*. 2013;9(1):26.
- Mathews KS, Long EF. A conceptual framework for improving critical care patient flow and bed use. *Annals Am Thorac Soc*. 2015;12(6):886–94.
- McPeake J, Bateson M, Christie F, Robinson C, Cannon P, Mikkelsen M, et al. Hospital re-admission after critical care survival: a systematic review and meta-analysis. *Anaesthesia*. 2022;77(4):475–85.
- Volpato S, Onder G, Cavalieri M, Guerra G, Sioulis F, Maraldi C, et al. Characteristics of nondisabled older patients developing New Disability Associated with Medical illnesses and hospitalization. *J Gen Intern Med*. 2007;22(5):668–74.
- Boyd CM, Landefeld CS, Counsell SR, Palmer RM, Fortinsky RH, Kresvick D, et al. Recovery of activities of daily living in older adults after hospitalization for acute medical illness. *J Am Geriatr Soc*. 2008;56(12):2171–9.
- Gallego-González E, Mayordomo-Cava J, Vidán MT, Valadés-Malagón MI, Serra-Rexach JA, Ortiz-Alonso J. Functional trajectories associated with acute illness and hospitalization in oldest old patients: impact on mortality. *Front Physiol*. 2022;13.
- Jenq GY, Tinetti ME. Post-acute care: who belongs where? *JAMA Intern Med*. 2015;175(2):296–7.
- McGilton KS, Vellani S, Yeung L, Chishtie J, Comisso E, Ploeg J, et al. Identifying and understanding the health and social care needs of older adults with multiple chronic conditions and their caregivers: a scoping review. *BMC Geriatr*. 2018;18(1):231.
- Mistiaen P, Duijnhouwer E, Wijkel D, Bont Md, Veeger A. The problems of elderly people at home one week after discharge from an acute care setting. *J Adv Nurs*. 1997;25(6):1233–40.
- Webkamigad S, Rowe R, Peltier S, Froehlich Chow A, McGilton KS, Walker JD. Identifying and understanding the health and social care needs of indigenous older adults with multiple chronic conditions and their caregivers: a scoping review. *BMC Geriatr*. 2020;20:1–19.
- Abdi S, Spann A, Borilovic J, de Witte L, Hawley M. Understanding the care and support needs of older people: a scoping review and categorisation using the WHO international classification of functioning, disability and health framework (ICF). *BMC Geriatr*. 2019;19(1):1–15.
- McBride SE, Beer JM, Mitzner TL, Rogers WA. Challenges for Home Health Care providers: a needs Assessment. *Phys Occup Ther Geriatr*. 2011;29(1):5–22.
- Yu Z, Gallant AJ, Cassidy CE, Boulos L, Macdonald M, Stevens S. Case Management models and Continuing Care: A literature review across nations, settings, approaches, and assessments. *Home Health Care Manage Pract*. 2020;33(2):96–107.
- Passalent LA, Landry MD, Cott CA. Exploring wait list prioritization and management strategies for publicly funded ambulatory rehabilitation services in Ontario, Canada: further evidence of barriers to access for people with chronic disease. *Healthc Policy*. 2010;5(4):e139.
- Deslauriers S, Raymond M-H, Laliberté M, Lavoie A, Desmeules F, Feldman DE, et al. Access to publicly funded outpatient physiotherapy services in Quebec: waiting lists and management strategies. *Disabil Rehabil*. 2017;39(26):2648–56.
- Deslauriers S, Raymond M-H, Laliberté M, Lavoie A, Desmeules F, Feldman DE, et al. Variations in demand and provision for publicly funded outpatient musculoskeletal physiotherapy services across Quebec, Canada. *J Eval Clin Pract*. 2017;23(6):1489–97.
- Raymond M-H, Demers L, Feldman DE. Waiting list management practices for home-care occupational therapy in the province of Quebec. *Can Health Social Care Community*. 2016;24(2):154–64.
- montreal Adlsses. Le Cadre De référence, services posthospitaliers en réadaptation fonctionnelle intensive en interne et soins subaigus pour la région de Montréal est une production. de l'Agence de la santé et des services sociaux de Montréal; 2010.
- Lafamme H, Valade S. Institut national d'excellence en santé et en services sociaux (INESSS). État Des pratiques: Pratiques organisationnelles en réadaptation à l'échelle Nationale et internationale pour la clientèle adulte présentant une incapacité motrice permanente ou temporaire. Québec, QC: INESSS; 2018. p. 104.
- Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, et al. Outcomes for implementation research: conceptual distinctions, measurement challenges, and research agenda. *Adm Policy Mental Health Mental Health Serv Res*. 2011;38:65–76.
- Des Jarlais DC, Lyles C, Crepaz N, Trend G. Improving the reporting quality of nonrandomized evaluations of behavioral and public health interventions: the TREND statement. *Am J Public Health*. 2004;94(3):361–6.
- Hoffmann TC, Glasziou PP, Boutron I, Milne R, Perera R, Moher D, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ: Br Med J*. 2014;348:g1687.
- (OPPQ) Opdpdq. Les deux professions Québec, Canada 2024 [Available from: <https://oppq.qc.ca/la-physiotherapie/les-deux-professions/>].

28. Akpan A, Roberts C, Bandeen-Roche K, Batty B, Bausewein C, Bell D et al. Standard set of health outcome measures for older persons. *BMC Geriatr*. 2018;18(1).
29. Ottenbacher KJ, Hsu Y, Granger CV, Fiedler RC. The reliability of the functional independence measure: a quantitative review. *Arch Phys Med Rehabil*. 1996;77(12):1226–32.
30. Maritz R, Tennant A, Fellinghauer C, Stucki G, Prodinge B. The functional independence measure 18-item version can be reported as a unidimensional interval-scaled metric: internal construct validity revisited. *J Rehabil Med*. 2019;51(3):193–200.
31. Cella D, Riley W, Stone A. The patient-reported outcomes Measurement Information System (PROMIS) developed and tested its first wave of adult self-reported health outcome item banks: 2005–2008. *J Clin Epidemiol*. 2010;63:1179–94.
32. HealthMeasures. PROMIS Scale v1.2 - Global Health 2021 [Available from: https://www.healthmeasures.net/index.php?option=com_instruments&view=measure&id=778&Itemid=992].
33. Kamper S, Maher C, Mackay G. Global rating of Change scales: a review of strengths and weaknesses and considerations for Design. *J Man Manipulative Therapy*. 2009;17(3):163–70.
34. Scott W, McCracken LM. Patients' impression of Change following treatment for Chronic Pain: global, specific, a single dimension, or many? *J Pain*. 2015;16(6):518–26.
35. Ferguson L, Scheman J. Patient global impression of change scores within the context of a chronic pain rehabilitation program. *J Pain*. 2009;10(4):573.
36. Reynolds T, Thornicroft G, Abas M, Woods B, Hoe J, Leese M, et al. Camberwell Assessment of need for the Elderly (CANE): development, validity and reliability. *Br J Psychiatry*. 2000;176(5):444–52.
37. Smith F, Orrell M. Does the patient-centred approach help identify the needs of older people attending primary care? *Age Ageing*. 2007;36(6):628–31.
38. Walters K, Iliffe S, Tai SS, Orrell M. Assessing needs from patient, carer and professional perspectives: the Camberwell Assessment of need for Elderly people in primary care. *Age Ageing*. 2000;29(6):505–10.
39. Wiczorowska-Tobis K, Talaraska D, Kropińska S, Jaracz K, Tobis S, Suwalska A, et al. The Camberwell Assessment of need for the Elderly questionnaire as a tool for the assessment of needs in elderly individuals living in long-term care institutions. *Arch Gerontol Geriatr*. 2016;62:163–8.
40. Carvacho R, Carrasco M, Lorca MBF, Miranda-Castillo C. Met and unmet needs of dependent older people according to the Camberwell Assessment of Need for the Elderly (CANE): a scoping review. *Rev Esp Geriatr Gerontol*. 2021;56(4):225–35.
41. van der Ploeg ES, Bax D, Boersma M, Nijpels G, van Hout HPJ. A cross-sectional study to compare care needs of individuals with and without dementia in residential homes in the Netherlands. *BMC Geriatr*. 2013;13(1):1–8.
42. Simons L, Petch A. Needs assessment and discharge: a Scottish perspective. *J Psychiatr Ment Health Nurs*. 2002;9(4):435–45.
43. Orrell M, Hancock G. CANE-Camberwell assessment of need for the elderly: a needs assessment for older mental health service users. Gaskell London; 2004.
44. Breitenstein SM, Gross D, Garvey CA, Hill C, Fogg L, Resnick B. Implementation fidelity in community-based interventions. *Res Nurs Health*. 2010;33(2):164–73.
45. Seyed-Nezhad M, Ahmadi B, Akbari-Sari A. Factors affecting the successful implementation of the referral system: a scoping review. *J Family Med Prim Care*. 2021;10(12):4364–75.
46. Greenwood-Lee J, Jewett L, Woodhouse L, Marshall DA. A categorisation of problems and solutions to improve patient referrals from primary to specialty care. *BMC Health Serv Res*. 2018;18(1):986.
47. Borrelli B. The Assessment, Monitoring, and Enhancement of Treatment Fidelity in Public Health Clinical Trials. *J Public Health Dent*. 2011;71(s1):S52–63.
48. Carroll C, Patterson M, Wood S, Booth A, Rick J, Balain S. A conceptual framework for implementation fidelity. *Implement Sci*. 2007;2(1):40.
49. Roberts DE, Holloway RG, George BP. Post-acute care discharge delays for neurology inpatients. *Neurology: Clin Pract*. 2018;8(4):302.
50. Bryan K. Policies for reducing delayed discharge from hospital. *Br Med Bull*. 2010;95(1):33–46.
51. Ostir GV, Berges I-M, Smith PM, Smith D, Rice JL, Ottenbacher KJ. Does change in functional performance affect quality of life in persons with orthopaedic impairment? *Soc Indic Res*. 2006;77(1):79–93.
52. D'Andrea A, Le Peillet D, Fassier T, Prendki V, Trombert V, Remy JL, et al. Functional independence measure score is associated with mortality in critically ill elderly patients admitted to an intermediate care unit. *BMC Geriatr*. 2020;20(1):334.
53. Jung HY, Kim TH, Park JH. Relationship between Berg Balance Scale and Functional Independence measure in Stroke patients. *J Korean Acad Rehabilitation Med*. 2005;29(2):167–70.
54. Lee D-J, Kim S-Y, Song C-H. The correlations between the balance test, functional movement, visual perception test and functional independent measure in stroke patients. *J Korean Phys Therapy*. 2009;21(2):39–45.
55. Hui D, Bruera E. What is clinically meaningful? The Use of Patient's Global Impression and Personalized Symptom goals. *Cancer*. 2016;122(1):159.
56. Kamper SJ, Maher CG, Mackay G. Global rating of change scales: a review of strengths and weaknesses and considerations for design. *J Man Manipulative Therapy*. 2009;17(3):163–70.
57. Perrot S, Lantéri-Minet M. Patients' global impression of change in the management of peripheral neuropathic pain: clinical relevance and correlations in daily practice. *Eur J Pain*. 2019;23(6):1117–28.
58. Wright J, Williams R, Wilkinson JR. Development and importance of health needs assessment. *BMJ*. 1998;316(7140):1310–3.
59. Stevens A, Gillam S. Needs assessment: from theory to practice. *BMJ*. 1998;316(7142):1448–52.
60. Araujo de Carvalho I, Epping-Jordan J, Pot AM, Kelley E, Toro N, Thiyagarajan JA, et al. Organizing integrated health-care services to meet older people's needs. *Bull World Health Organ*. 2017;95(11):756–63.
61. Briggs AM, Jordan JE, Jennings M, Speerin R, Bragge P, Chua J, et al. Supporting the evaluation and implementation of musculoskeletal models of care: a globally informed framework for judging readiness and success. *Arthritis Care Res*. 2017;69(4):567–77.
62. Briggs AM, Valentijn PP, Thiyagarajan JA, Araujo de Carvalho I. Elements of integrated care approaches for older people: a review of reviews. *BMJ Open*. 2018;8(4):e021194.
63. World Health O. World report on ageing and health. World Health Organization; 2015.
64. Soo WK, Yin V, Crowe J, Lane H, Steer CB, Därziņš P, et al. Integrated care for older people with cancer: a primary care focus. *Lancet Healthy Longev*. 2023;4(6):e243–5.
65. Verweij L, van de Korput E, Daams JG, Ter Riet G, Peters RJG, Engelbert RHH, et al. Effects of postacute multidisciplinary rehabilitation including exercise in out-of-hospital settings in the aged: systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2019;100(3):530–50.
66. Preitschopf A, Holstege M, Ligthart A, Groen W, Burchell G, Pol M, et al. Effectiveness of outpatient geriatric rehabilitation after inpatient geriatric rehabilitation or hospitalisation: a systematic review and meta-analysis. *Age Ageing*. 2023;52(1):afac300.
67. Handoll HHG, Cameron ID, Mak JCS, Panagoda CE, Finnegan TP. Multidisciplinary rehabilitation for older people with hip fractures. *Cochrane Database Syst Reviews*. 2021(11).
68. Wilson MG. Rapid Synthesis: Examining the Effectiveness and Cost-effectiveness of Rehabilitation-care Models for Frail Seniors. 2013.
69. Lin C-J, Cheng S-J, Shih S-C, Chu C-H, Tjung J-J. Discharge planning. *Int J Gerontol*. 2012;6(4):237–40.
70. Hunter T, Birmingham J. Preventing readmissions through comprehensive discharge planning. *Prof case Manage*. 2013;18(2):56–63.
71. Gitlin LN. Environmental adaptations for older adults and their families in the home and community. *Int Handb Occup Therapy Interventions*. 2009:53–62.
72. Low L-F, Yap M, Brodaty H. A systematic review of different models of home and community care services for older persons. *BMC Health Serv Res*. 2011;11(1):93.
73. Ellis B, Connell NAD. Factors determining the current use of Physiotherapy assistants: views on their future role in the South and West UK Region. *Physiotherapy*. 2001;87(2):73–82.
74. Jensen M, Junod C, Fatemi N, Liew K, Ulaj E, Bean CC, et al. Navigating the Grey Zone of Physiotherapy Assistant Autonomy in Home Care: perspectives of physiotherapists and Physiotherapy assistants. *Physiother Can*. 2022;74(3):249–56.
75. Akobeng AK. Understanding type I and type II errors, statistical power and sample size. *Acta Paediatr*. 2016;105(6):605–9.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.