

Protocol

Acceptability and Feasibility of a Return-to-Work Intervention for Posttreatment Breast Cancer Survivors: Protocol for a Co-design and Development Study

Karine Bilodeau^{1,2}, PhD; Marie-Michelle Gouin³, PhD; Alexandra Lecours⁴, PhD; Valérie Lederer⁵, PhD; Marie-José Durand³, PhD; Kelley Kilpatrick^{2,6}, PhD; David Lepage⁷, MSc; Lauriane Ladouceur-Deslauriers⁷, MSc; Tomas Dorta⁸, PhD

¹Faculty of Nursing, University of Montreal, Montreal, QC, Canada

²Centre de recherche Hopital Maisonneuve Rosemont, Montreal, QC, Canada

³Faculté de médecine et des sciences de la santé, University of Sherbrooke, Longueuil, QC, Canada

⁴Département de relations industrielles, Université du Québec à Trois-Rivières, Trois-Rivières, QC, Canada

⁵Département de relations industrielles, Université du Québec en Outaouais, Gatineau, QC, Canada

⁶Ingram School of Nursing, McGill University, Montreal, QC, Canada

⁷Centre intégré universitaire de santé et de services sociaux de l'Est de l'île de Montréal, Montréal, QC, Canada

⁸Faculté de l'aménagement, École de Design, Université de Montréal, Montreal, QC, Canada

Corresponding Author:

Karine Bilodeau, PhD

Faculty of Nursing

University of Montreal

PO Box 6128

Station Centre-ville

Montreal, QC, H3C 3J7

Canada

Phone: 1 514 343 6111 ext 43254

Fax: 1 514 343 2306

Email: karine.bilodeau.2@umontreal.ca

Related Article:

This is a corrected version. See correction statement in: <https://www.researchprotocols.org/2023/1/e50784>

Abstract

Background: The mortality rate from breast cancer has been declining for many years, and the population size of working-age survivors is steadily increasing. However, the recurrent side effects of cancer and its treatment can result in multiple disabilities and disruptions to day-to-day life, including work disruptions. Despite the existing knowledge of best practices regarding return to work (RTW) for breast cancer survivors, only a few interdisciplinary interventions have been developed to address the individualized needs and multiple challenges of breast cancer survivors, health care professionals, and employer and insurer representatives. Thus, it seems appropriate to develop RTW interventions collaboratively by using a co-design approach with these specific stakeholders.

Objective: This paper presents a protocol for developing and testing an innovative, interdisciplinary pilot intervention based on a co-design approach to better support RTW and job retention after breast cancer treatment.

Methods: First, a participatory research approach will be used to develop the intervention in a co-design workshop with 12 to 20 participants, including people affected by cancer, employer and insurer representatives, and health care professionals. Next, a pilot intervention will be tested in a primary care setting with 6 to 8 women affected by breast cancer. The acceptability and feasibility of the pilot intervention will be pretested through semistructured interviews with participants, health care professionals, and involved patient partners. The transcribed data will undergo an iterative content analysis.

Results: The first phase of the project—the co-design workshop—was completed in June 2021. The pilot test of the intervention will begin in spring 2022. The results from the test will be available in late 2022.

Conclusions: The project will offer novel data regarding the use of the co-design approach for the development of innovative, co-designed interventions. In addition, it will be possible to document the acceptability and feasibility of the pilot intervention with a primary care team. Depending on the results obtained, the intervention could be implemented on a larger scale.

International Registered Report Identifier (IRRID): DERR1-10.2196/37009

(*JMIR Res Protoc* 2022;11(4):e37009) doi: [10.2196/37009](https://doi.org/10.2196/37009)

KEYWORDS

co-design; breast cancer; intervention; return-to-work; primary care; qualitative

Introduction

It has been reported that 1 in 8 Canadian women will develop breast cancer in their lifetime [1,2]. Of these, 88% will survive for more than 5 years, and of these survivors, 50% are of working age. In the current context of labor shortages [3], the contribution of those with a cancer diagnosis who want to return to work (RTW) is valuable. Indeed, many survivors desire to RTW because it is a sign of a return to a “normal” life [2,4-7]. On the other hand, 1 in 5 women will leave their jobs due to an inability to perform their tasks resulting from recurrent side effects of cancer or its treatments [8]. There are a variety of reasons for the development of these limiting side effects. Breast cancer treatments, such as mastectomy or chemotherapy, can cause significant physical symptoms [9] (eg, pain, the loss of arm mobility, and lymphedema) that may limit the act of lifting or work involving repetitive motions [10]. These symptoms and physical limitations can severely complicate and even hinder daily work [7,10]. Fatigue, as well as memory loss and difficulty concentrating, can also impact RTW [11]. Furthermore, breast cancer survivors often live more precariously than the general population due to taking repeated sick leaves from work or having only part-time employment after cancer treatments [12-16]. Despite these challenges, both working and having good working conditions are related to better health [17], as is allowing for the maintenance of social interactions, self-esteem, psychological well-being, and financial security [2,18,19].

Interdisciplinary interventions that support RTW and are specifically tailored to the issues experienced by breast cancer survivors remain rare in Canada and do not exist in Quebec, and this scarcity of tailored interventions seems likely to continue [20-22]. Indeed, a Cochrane review about RTW interventions for patients who have completed cancer treatments revealed that interventions that targeted multiple modalities (physical, psychological, and vocational) and were delivered by an interdisciplinary team seemed more appropriate for addressing the needs of cancer survivors [23]. This multimodal type of intervention appears promising, although the effect size remained tenuous and suggested the need for further study [23]. Other studies suggest using more comprehensive approaches, such as discussing work-related issues during psycho-oncological care [24,25], involving employers [26], and performing early intervention [27]. It has even been suggested that work-related issues should be discussed as early as the active treatment period to maintain social roles, including those of workers [25,28]. According to a scoping review [27], RTW interventions that have been identified to support breast cancer survivors are offered in ad hoc interventions by different health

care professionals and are highly variable in terms of information booklets, physical activities, and deployment times (eg, at the end of treatment) [27]. More structured interdisciplinary interventions remain to be developed to meet the unique needs of this clientele regarding RTW.

It seems desirable and realistic that interdisciplinary interventions be offered by a primary care team [20]. By definition, primary care can include health promotion, disease prevention, the monitoring of chronic and episodic diseases, and rehabilitation [29]. It is also possible that such a team would be able to address the RTW-related concerns of women affected by breast cancer, which include symptom management, RTW-related decision-making, resource navigation, and the reintegration of daily activities [4,30]. Primary care teams promote interdisciplinary work and provide services as close to a given population as possible [31]. Furthermore, given the scope of primary care, it is strongly recommended that those affected by cancer be managed by such teams during the recovery period [32,33]. It is also suggested that intervention components should be deployed during key points in the experience of cancer survivorship at 1, 3, and 6 months after cancer treatment and encourage the self-management of side effects (eg, cognitive difficulties and fatigue), RTW-related decision-making, resource navigation, and the reintegration of daily activities [4,21]. As for the intensity of interventions that should be offered, it is mentioned that nearly 50% of affected individuals require more assiduous support due to the presence of persistent side effects [34], such as severe fatigue or physical limitations induced by lymphedema [35,36]. Moreover, studies have shown that women affected by breast cancer who have received chemotherapy treatments experience more difficulty during RTW due to persistent side effects [37-40]. This clientele therefore should receive personalized support for coping with the challenges of completing cancer treatments, including RTW.

RTW after cancer treatment is complex because it involves multiple stakeholders [41], including breast cancer survivors, health care professionals, employers, and insurer representatives, who come from multiple settings and have divergent concerns, constraints, and resources. The available interventions that support RTW after cancer treatment have proven to be insufficient in connecting workplace stakeholders and addressing RTW-related coordination challenges [20,27]. Our project aims to develop a supportive RTW intervention to address the current lack of coordinated, interdisciplinary interventions offered in primary care. Based on current evidence, it is imperative to develop new interventions to support the RTW of breast cancer survivors. These interventions should have several characteristics that align with the Medical Research Council

criteria [42]. As such, these interventions will be complex, especially given the variety of the intervention components needed, the interactions between these components, the number of people or organizational levels involved, the degree of flexibility required, and the need for the ability to adapt to specific contexts. The development of a complex intervention requires several phases that do not follow a linear sequence [42]. To address this complexity, it is advisable to combine theoretical, empirical, and experiential approaches when designing interventions [43]. However, these approaches are time-consuming, and it is difficult to address all of the issues related to implementing interventions [44]. To engage stakeholders and achieve contextually appropriate intervention components, our project will involve a coideation process for developing an RTW intervention and its components through a co-design approach. This novel methodological approach to developing RTW interventions brings together cross-sectoral stakeholders, including health care professionals, employer and insurer representatives, and breast cancer survivors. The components of interventions can be difficult for stakeholders to understand, and such components can be difficult to contextualize during the development process. The co-design approach is therefore appropriate, as it was developed so that stakeholders can actively participate in a process of coideation for the interventions that are dedicated to them. The proposed approach differs from others because it offers the possibility for all participants to cocreate and negotiate ideas simultaneously, starting from the beginning (conceptual phase) of the co-design process [45]. Another characteristic of the


approach is the presence of design professionals who facilitate the coideation process [46] by making graphic representations that are adapted to the participants to support the creative process. Finally, coideation has proven to be an innovative approach that has several potential benefits. First, coideation allows for the creation of an intervention without preconceived ideas. Second, the creation of an intervention can be faster, allowing for quicker testing in a pilot study context. This provides better direction for subsequent developments.

Given the need for support among breast cancer survivors, the inherent challenges of RTW, and the complexity of intervening to support RTW, there is a need to coconstruct a coordinated, interdisciplinary intervention with RTW stakeholders that is offered by a primary care team. To achieve this, we propose a co-design approach. The goal of the study is to develop and test a pilot intervention for supporting RTW after breast cancer treatment. The intervention will be offered by a primary care team. The objectives are to (1) develop an innovative intervention in collaboration with key stakeholders, (2) test the intervention with a primary care team, and (3) determine the acceptability and feasibility of the intervention in a primary care setting.

Methods

The project is divided into the following two phases: the development of the pilot intervention and the feasibility assessment (Figure 1).

Figure 1. Study design. RTW: return to work.



Phase 1: Pilot Intervention Development

Study Design

A participatory action research design [47] and a co-design approach [48] will be used. A participatory action design engages researchers and participants in a reflective process when they seek solutions [47,49]. In addition, it encourages the partnership between researchers and stakeholders throughout the research process. Participatory research makes it possible to produce useful and precise knowledge that is consistent with

the realities of key RTW stakeholders. The synergy created in partnerships allows for studies that are both culturally appropriate for target audiences and logistically realistic [50]. The development of the pilot intervention will be inspired by the reflexive cycle—a key part of the participatory approach that encourages describing and defining observations, analyzing and interpreting ideas, and finding solutions [49,51]. In our case, the solutions will be creative and innovative. This process will take place during a co-design workshop with key stakeholders. Co-design allows people to actively participate in a process of coideation for finding innovative solutions,

during which they can simultaneously participate in the cocreation process at the beginning of a project by following a democratic and multidisciplinary perspective [48].

Population and Recruitment

A variety of participants (n=12-20) representing key RTW stakeholders will participate in the codevelopment of the pilot intervention. For this phase, snowball sampling will be conducted [52]. All participants must speak French and be aged ≥ 18 years. To be selected for the codevelopment phase, breast cancer survivors must have experienced RTW after cancer treatments for at least 1 year, employer or insurer representatives must have at least coordinated or accompanied the RTW of a woman affected by breast cancer, and health care professionals must have previously provided support to a woman with breast cancer during either oncology or primary care. Participants will be approached through the research team's professional networks. This strategy is appropriate, given the nature of the interactive activity and the need for participants to be engaged and involved in the RTW journey. These are features of some participatory and co-design approaches [53].

Sequence of the Co-design Workshop

In preparation for the workshop, a 10-minute informational video will be sent to participants. This video will present the current knowledge regarding RTW for cancer survivors as well as recommendations from research. By using the Zoom platform (Zoom Video Communications Inc), a co-design workshop will be held over 4 hours. Participants will be informed that the purpose of the workshop is to co-design components of a pilot intervention that will be performed by a primary care team to address the issues faced by survivors who RTW after breast cancer treatment. Further, 3 case scenarios that represent the potential issues of RTW after breast cancer treatment will be presented to participants. Because the workshop will include key RTW stakeholders, the scenarios will serve as a "representative artifact," that is, a deliberation strategy involving interactions between public participants and health professionals that encourage exchanges via a common and appropriate language [54]. Participants will also be guided through the following two dimensions of intervention: the assessment of cancer and treatment side effects and RTW discussion and planning. Instead, participants will be invited to think of novel solutions.

During the workshop, participants will interact in plenary and breakout sessions. Further, 2 to 4 diverse subgroups will be created, depending on the number of participants and the representativeness of the RTW stakeholders, for activities in a breakout room. Each subgroup will be accompanied by a

facilitator and a design professional who will help to frame the discussions. The facilitator will be a postdoctoral fellow or a graduate student from an undergraduate design program (eg, industrial design). The involvement of design professionals is recommended for health science initiatives [46]. As presented in Table 1, the workshop will be divided into the following five steps: (1) the reframing of the problem, (2) immature coideation, (3) mature coideation, (4) the presentation of subgroup solutions, and (5) the debriefing of the workshop experience. In step 1, participants will be invited to discuss the problem of RTW after cancer treatment. The facilitator can refer to the case scenarios and frame the discussions to raise priority issues. In step 2, participants will be asked to develop solutions. The facilitator will assist the participants in developing solutions that are relevant to issues that were identified earlier and representing them graphically (eg, diagrams and drawings). At this stage, the ideas are to be formulated but not yet fully developed. In step 3, participants will be invited to fully develop the solutions that they find the most important, in some cases by making more detailed graphic representations. At this stage, specific details will be offered by participants in each subgroup with regard to the implementation or components of the selected solutions. In step 4, a plenary discussion will be held to summarize the work of each subgroup. Finally, in step 5, the workshop will end with a debriefing on the participants' perceptions of the positive and negative aspects to be retained or considered for the repetition of the workshop. Of note, throughout the workshop, participants will make use of a graphical representational ecosystem (eg, drawings and digital sketches) [55], along with the verbal exchanges, to externalize their ideas and discuss proposals more clearly. In addition, subgroup discussions can be structured by using the following key principles of design conversation [48,56]: (1) naming the problem, (2) constraining the ideas, (3) proposing the ideas (the key element), (4) negotiating the ideas through questioning and explaining, (5) making decisions, and (6) making design advances with graphical representations. This approach keeps discussions focused on a common goal and ensures that stakeholders collaborate within a limited time frame during all of the workshop stages. The activity will take place on the internet due to the current pandemic context and will be conducted by using the Miro platform (Participatory Culture Foundation; Figures 2-4). A pretest of the workshop will be conducted with students before the activity.

The workshop will be recorded, and explicit notes will be written verbatim. The deliverable at this stage will be the general guidelines for an intervention that supports RTW after breast cancer treatment. The intervention will be designed so that it is fit for use in primary care settings.

Table 1. Sequence of the co-design workshop.

| Activities and steps | Length (total: 235 minutes) |
|---|---|
| Welcome and introduction | <ul style="list-style-type: none"> • 10 minutes (plenary session) |
| Step 1: reframing the problem | <ul style="list-style-type: none"> • 30 minutes (subgroup session) • 15 minutes (plenary session) |
| Transition and breakout room integration | <ul style="list-style-type: none"> • 5 minutes |
| Step 2: immature coideation | <ul style="list-style-type: none"> • 30 minutes (subgroup session) • 15 minutes (plenary session) |
| Break | <ul style="list-style-type: none"> • 15 minutes |
| Step 3: mature coideation | <ul style="list-style-type: none"> • 30 minutes (subgroup session) • 15 minutes (plenary session) |
| Transition and breakout room integration | <ul style="list-style-type: none"> • 5 minutes |
| Step 4: presentation of subgroup solutions | <ul style="list-style-type: none"> • 30 minutes (plenary session) |
| Transition and breakout room integration | <ul style="list-style-type: none"> • 5 minutes |
| Step 5: debriefing the workshop experience | <ul style="list-style-type: none"> • 30 minutes (plenary session) |

Figure 2. Miro interface for “Reframing the problem.”

CLÉ ANIMATION - RECADRAGE DU PROBLÈME

1. Quels sont les défis qui semblent le plus prioritaires vis-à-vis du retour au travail ? Du point de vue du professionnel de la santé ? Du point de vue de l'assureur ?
2. Quel devrait être l'attitude des professionnels de la santé ? Celle de l'informaticien, celle du travailleur social, celle du professionnel en réadaptation, celle du médecin ?
3. Quelles informations devraient être communiquées ? Par les professionnels ? Par les personnes touchées par le cancer ? Par les assureurs ? Par les employeurs ?
4. Quels sont les défis liés à la discussion du retour au travail suite à un cancer du sein ?
5. Quels sont les défis liés à la planification de la reprise du travail ? (p. ex., difficultés en lien avec les tâches, l'horaire de travail, l'embarquement, la persistance des effets secondaires du travail, la référence aux experts, etc.)
6. Quels se qui semble le plus prioritaire dans chacune des vignettes ?

ÉQUIPE 1

PREMIÈRE SESSION : RECADRER LE PROBLÈME

Acteurs / Personnes

Lieux

Symptômes

Outils

Type de travail

Les enjeux et les problèmes clés les plus forts

Scénario 1 :

Sylvia a 44 ans. Elle habite avec son conjoint et elle est la mère de 3 enfants. Elle travaille comme prépariste au service à la clientèle dans une épicerie. Il y a 2 ans, Sylvia a été diagnostiquée d'un cancer du sein. Après la fin des traitements, Sylvia se trouve affaiblie par les traitements qu'elle a subis (chimiothérapie, radiothérapie, tamoxifène sans succès). Sylvia ne se sent pas prête pour retourner au travail immédiatement parce qu'elle a peur qu'on lui demande d'accomplir la même quantité de travail qu'avant, sans tenir compte de sa fatigue et de son inconfort au bras gauche qui semble persister après la fin de ses traitements. D'un autre côté, elle se dit qu'elle doit retourner travailler, car la famille s'en entend pendant son congé maladie.

Scénario 2 :

Cécile a 40 ans et elle est mère d'un jeune fils de 8 ans qu'elle a élevée seule. Elle occupait un emploi de technicienne en informatique dans un établissement d'enseignement. Il y a un an et demi, Cécile a subi une tumeur cérébrale pour un cancer du sein, de la radiothérapie et de la chimiothérapie. Elle se sent fatiguée et se sent comme qu'elle oublie beaucoup de choses. Elle doute par moments de sa capacité à travailler devant son écran toute la journée. Elle doit faire deux semaines par jour pour pouvoir faire ses tâches à la maison. Cécile se questionne quant aux meilleures conditions pour se réintégrer du travail en termes d'horaires et de tâches. Cécile se sent malgri tout chanceuse, car elle ne sent pas de pression de retourner en emploi, car elle a une bonne assurance santé.

Scénario 3 :

Linda a 58 ans et habite avec son conjoint. Elle est mère de deux jeunes adultes qui ont quitté la maison. Elle occupait un poste de cadre supérieure dans une institution bancaire. Il y a 11 mois, elle a eu une amputation de son bras et a eu des épisodes de dépression et de douleur. Linda a fini ses traitements il y a deux semaines et se questionne déjà quant à son retour au travail. Linda se sent avachi. Elle ne voit pas comment elle pourra gérer sa tâche de travail et rester concentrée sur tous les devoirs qui doivent l'attendre. Elle a aussi des douleurs sous l'aisselle droite et se sent inconfortable assise devant un ordinateur. Linda se remet en question quant à son éventuel retour au travail. Elle se sent seule et se demande si d'autres femmes vivent la même chose.

Y-a-il des problèmes inter-liés ?

<https://www.researchprotocols.org/2022/4/e37009>

JMIR Res Protoc 2022 | vol. 11 | iss. 4 | e37009 | p. 6
(page number not for citation purposes)

XSL-FO
RenderX

Figure 3. Miro interface for “Immature coideation.”




Figure 4. Miro interface for “Mature coideation.”



Phase 2: Pilot Test and Evaluation of Acceptability and Feasibility

The pilot intervention will be delivered by a primary care team and patient partners who will be trained in the intervention. A resource person will be available to answer their questions or provide coaching during the project.

Population and Recruitment

By using purposive sampling [57], 6 to 8 breast cancer survivors will be recruited to test the pilot intervention. The selection criteria will include women aged 18 to 60 years; those who have completed breast cancer treatments, such as surgery, chemotherapy (doxorubicin, cyclophosphamide, and paclitaxel), and radiation therapy; and those who are planning to RTW within the next year. Surgeon oncologists will help with

targeting potential participants who are nearing the completion of their cancer treatments.

Evaluation of Acceptability and Feasibility

At the end of the project, the acceptability and feasibility of the pilot intervention will be measured by conducting semistructured individual interviews with participants (n=6-8) and health care professionals, as well as patient partners who participated in the intervention (n=4-8). The use of qualitative interviews is recommended during a feasibility study [58]. The interview guide and analysis will be based on the Theoretical Framework of Acceptability [58], which explains that acceptability is a multidimensional construct with 7 components. Specifically, the interview questions will address (1) affective attitudes toward the intervention, (2) the effort required to complete the intervention, (3) consistency with the individuals' values, (4) individuals' understanding of the intervention and how it works, (5) the perceived benefits provided by the intervention, (6) perceived efficacy, and (7) the perceived ability to complete the intervention. The interview guide will be pretested with 2 people who have the same characteristics as those of the participants.

Data Analysis

The transcribed interview will undergo an iterative content analysis, which will include the following activities: condensation, data presentation, and the development and verification of findings [59]. Further, 2 team members will conduct the coding process. The data from the breast cancer survivors will be contrasted to highlight similarities and differences in experiences with the pilot intervention. The same exercise will be conducted for the health care professionals. In addition, to validate the emerging findings, a discussion with the coresearchers will be conducted throughout the research process. This discussion will consider the data transcripts and field notes. NVivo software (QSR International) will be used for qualitative data management. To ensure the quality of the study, techniques such as data triangulation will be used to ensure internal credibility and validity. Further, we will validate some of the participants' data to ensure their reliability, assess the procedural documentation of the research process to determine accountability, and provide a detailed description of the context to ensure external transferability and validity [59].

Ethics Approval

Ethics approval was granted in May 2021 for the first component of the project (Centre intégré universitaire de santé et de services sociaux de l'Est-de-l'Île-de-Montréal; project number: #2022-2610).

Results

The project received funding in March 2021. The co-design workshop took place on June 16, 2021. A total of 11 people participated in the activity. The transcripts of the discussions were analyzed and helped to target the following intervention themes for intervention development: (1) mitigating the assessment and self-management of side effects, (2) assessing RTW needs and abilities, and (3) communicating with the employer. The clinical tools for the intervention (questionnaire

and decision support tree) and the intervention logic model remain to be finalized. The preliminary results suggest that the pilot intervention should take place in a primary care setting in the Montreal area (Quebec, Canada). It is anticipated that 3 meetings will occur at 1, 4, and 6 months after treatment is completed [21]. During these meetings, a health professional will analyze RTW challenges and may propose solutions that are tailored to women's needs. The intervention will include pairing participants with patient partners who will be able to answer the participants' questions and share their RTW experiences [60]. Depending on the pandemic context of COVID-19 in Canada, recruitment will begin in spring 2022.

Discussion

Principal Results

In brief, the project will allow us to document the relevance of an RTW intervention that is delivered by a primary care team. The shared views of patients and health care professionals will help us determine the feasibility and acceptability of the intervention. In addition, the project will offer methodological recommendations for the use of a co-design approach in intervention development. More specifically, the project will have clinical, methodological, and organizational benefits.

On the clinical level, the primary outcome of the project is the development of a program that supports RTW and is based on the perspectives of RTW stakeholders, including breast cancer survivors [41]. Only a few European studies have mobilized these actors for the development of RTW interventions for patients who have completed cancer treatments [61,62]. The participatory approach that will be used in the project properly contextualizes the intervention to a Canadian context. This should facilitate implementation in the second phase of the study. The results of the pilot project will also provide information for improving the intervention before large-scale implementation, which will make it possible to evaluate its effects. The project will also offer an initial solution for women affected by breast cancer who need support [63] but are not covered by current health services [21]. Indeed, it is more urgent than ever to address the issues of RTW after cancer treatment to limit the development of disabilities, particularly as work is a known determinant of health and social participation is beneficial to individuals [17,63]. Additionally, and perhaps more broadly, RTW-related disabilities result in considerable consequences that also affect society as a whole [64-66], especially when absences are prolonged over time [64]. Fostering RTW is therefore essential, especially in the context of an aging population and labor shortages [67].

In terms of methodology, the project will explore an innovative and participatory approach that is perfectly suited to the trend of including patients and those who are involved in creating interventions (eg, a patient-centered research strategy [68]). To engage patients and partners, the use of approaches that have been adapted from design-related approaches [46] (eg, experience-based design [69] and Hacking Health hackathons [70]) has become increasingly common in health care. These approaches are appreciated and are essential, given the complexity of the interventions that are to be developed in health

care [42]. That said, many deplore the time and investment required to complete such processes [71,72]. Our project addresses this concern via the use of a structured co-design ideation approach to develop innovative and robust solutions within a reasonable time frame. Through this approach, we will be able to decrease intervention development time, increase cross-sectoral stakeholder participation, and make the intervention creation process more efficient. The coideation process will promote the creation of robust solutions for effective implementation and result in a decrease in the time required for the intervention creation cycle (ie, from ideation to implementation). The new data provided by our project will facilitate the development of new interdisciplinary interventions that benefit clientele in various health care settings.

This protocol is the first to propose a pilot RTW intervention for patients who have completed cancer treatments that will be delivered by a primary care team. It is widely documented that the supply of care after cancer treatment is inadequate [21,73,74]. It remains difficult to provide services to people affected by cancer beyond specialized oncology services, despite numerous international recommendations [75]. It is hoped that our RTW-themed project will encourage health care professionals and employers to commit to its goals and principles by demonstrating that the intervention will be acceptable to participants and consistent with the mission of primary care teams (ie, promoting health and preventing disease). The project will thus offer new data on the feasibility of offering this follow-up intervention in a primary care setting.

Anticipated Challenges and Limitations

This protocol is proposed in a Canadian context (province of Quebec). In Canada, health care is free and primary care services are readily accessible to the general population. It should be

noted that the majority of Canadians affected by cancer have paid sick leave included in their private insurance contracts [16]. Many people, including those without private insurance, only have access to 15 weeks of compensation from the Canadian government. Moreover, with regard to RTW after cancer treatment, no agreement has been reached with employers, and no legislation has been created, as is the case in France. This protocol must therefore be interpreted in this context. Furthermore, the evaluation of feasibility and acceptability will be based on a few qualitative interviews. The number of interviews should be sufficient, that is, from a scientific point of view [76], for documenting whether the intervention is feasible in this context. Finally, the pandemic context of COVID-19 may limit the testing of the project. Primary care teams are facing the off-loading of their professionals to other areas, including those that require vaccination efforts. To mitigate these effects, we will rely on the support of primary care physicians; oncology specialists; and members of the research team (DL and LLD), who are also clinicians. Despite the unfavorable context, health care professionals have reiterated their willingness to move forward, which demonstrates the relevance of the project and the interest of the health care organizations.

Conclusions

This protocol proposes to develop and test an intervention that supports RTW after breast cancer treatment and is delivered by a primary care team. The project will provide novel data on the use of a co-design method for the development of complex interventions. In addition, the results of the project will allow us to better document the acceptability and feasibility of the intervention, which will be delivered by a primary care team. Depending on the results obtained, the project could be tested on a larger scale.

Acknowledgments

We would like to thank the Quebec Rehabilitation Research Network for its financial support of the project. We would like to extend our thanks to Sana Boudhraa for her help with co-design workshop organization. Finally, we would like to thank Asma Fadhlou and Cynthia Henriksen.

Conflicts of Interest

None declared.

Multimedia Appendix 1

Peer-review reports.

[PDF File (Adobe PDF File), 626 KB-Multimedia Appendix 1]

References

1. Canadian cancer statistics 2021. Canadian Cancer Society. 2021 Nov. URL: <https://tinyurl.com/mryvph5f> [accessed 2022-04-12]
2. Wells M, Williams B, Firmigl D, Lang H, Coyle J, Kroll T, et al. Supporting 'work-related goals' rather than 'return to work' after cancer? A systematic review and meta-synthesis of 25 qualitative studies. *Psychooncology* 2013 Jun;22(6):1208-1219 [doi: [10.1002/pon.3148](https://doi.org/10.1002/pon.3148)] [Medline: [22888070](https://pubmed.ncbi.nlm.nih.gov/22888070/)]
3. Grenier A. Les pénuries de main-d'œuvre guettent-elles le marché du travail québécois? *Regards sur le travail* 2011;7(2):1-12 [FREE Full text]
4. Bilodeau K, Tremblay D, Durand MJ. Return to work after breast cancer treatments: Rebuilding everything despite feeling "in-between". *Eur J Oncol Nurs* 2019 Aug;41:165-172 [doi: [10.1016/j.ejon.2019.06.004](https://doi.org/10.1016/j.ejon.2019.06.004)] [Medline: [31358250](https://pubmed.ncbi.nlm.nih.gov/31358250/)]

5. Brusletto B, Torp S, Ihlebæk CM, Vinje HF. A five-phase process model describing the return to sustainable work of persons who survived cancer: A qualitative study. *Eur J Oncol Nurs* 2018 Jun;34:21-27 [doi: [10.1016/j.ejon.2018.03.003](https://doi.org/10.1016/j.ejon.2018.03.003)] [Medline: [29784134](https://pubmed.ncbi.nlm.nih.gov/29784134/)]
6. Drageset S, Lindstrøm TC, Underlid K. "I just have to move on": Women's coping experiences and reflections following their first year after primary breast cancer surgery. *Eur J Oncol Nurs* 2016 Apr;21:205-211 [doi: [10.1016/j.ejon.2015.10.005](https://doi.org/10.1016/j.ejon.2015.10.005)] [Medline: [26521054](https://pubmed.ncbi.nlm.nih.gov/26521054/)]
7. Sun W, Chen K, Terhaar A, Wiegmann DA, Heidrich SM, Tevaarwerk AJ, et al. Work-related barriers, facilitators, and strategies of breast cancer survivors working during curative treatment. *Work* 2016;55(4):783-795 [FREE Full text] [doi: [10.3233/WOR-162449](https://doi.org/10.3233/WOR-162449)] [Medline: [28059814](https://pubmed.ncbi.nlm.nih.gov/28059814/)]
8. Return to work concerns faced by people dealing with cancer and caregivers: Literature review and consultation. Canadian Partnership Against Cancer. 2012 Apr. URL: <https://s22457.pcdn.co/wp-content/uploads/2018/12/Return-To-Work-Concerns-EN.pdf> [accessed 2022-04-12]
9. Stubblefield MD, Keole N. Upper body pain and functional disorders in patients with breast cancer. *PM R* 2014 Feb;6(2):170-183 [doi: [10.1016/j.pmrj.2013.08.605](https://doi.org/10.1016/j.pmrj.2013.08.605)] [Medline: [24360839](https://pubmed.ncbi.nlm.nih.gov/24360839/)]
10. Zomkowski K, de Souza BC, da Silva FP, Moreira GM, de Souza Cunha N, Sperandio FF. Physical symptoms and working performance in female breast cancer survivors: a systematic review. *Disabil Rehabil* 2018 Jun;40(13):1485-1493 [doi: [10.1080/09638288.2017.1300950](https://doi.org/10.1080/09638288.2017.1300950)] [Medline: [28325132](https://pubmed.ncbi.nlm.nih.gov/28325132/)]
11. Porro B, Michel A, Zinzindohoué C, Bertrand P, Monrignal E, Trentini F, et al. Quality of life, fatigue and changes therein as predictors of return to work during breast cancer treatment. *Scand J Caring Sci* 2019 Jun;33(2):467-477 [doi: [10.1111/scs.12646](https://doi.org/10.1111/scs.12646)] [Medline: [30664270](https://pubmed.ncbi.nlm.nih.gov/30664270/)]
12. Jeon SH. Les effets du cancer sur l'emploi et les gains des survivants du cancer. *Statistique Canada*. 2014 Sep. URL: <https://www150.statcan.gc.ca/n1/pub/11f0019m/11f0019m2014362-fra.htm> [accessed 2022-04-12]
13. Fenn KM, Evans SB, McCorkle R, DiGiovanna MP, Puzstai L, Sanft T, et al. Impact of financial burden of cancer on survivors' quality of life. *J Oncol Pract* 2014 Sep;10(5):332-338 [doi: [10.1200/JOP.2013.001322](https://doi.org/10.1200/JOP.2013.001322)] [Medline: [24865220](https://pubmed.ncbi.nlm.nih.gov/24865220/)]
14. O'Connor JM, Kircher SM, de Souza JA. Financial toxicity in cancer care. *J Community Support Oncol* 2016 Mar;14(3):101-106 [doi: [10.12788/jcso.0239](https://doi.org/10.12788/jcso.0239)] [Medline: [27058866](https://pubmed.ncbi.nlm.nih.gov/27058866/)]
15. Lorgelly PK, Neri M. Survivorship burden for individuals, households and society: Estimates and methodology. *J Cancer Policy* 2018 May;15:113-117 [FREE Full text] [doi: [10.1016/j.jcpo.2018.02.005](https://doi.org/10.1016/j.jcpo.2018.02.005)]
16. Longo CJ, Fitch MI, Loree JM, Carlson LE, Turner D, Cheung WY, et al. Patient and family financial burden associated with cancer treatment in Canada: a national study. *Support Care Cancer* 2021 Jun;29(6):3377-3386 [FREE Full text] [doi: [10.1007/s00520-020-05907-x](https://doi.org/10.1007/s00520-020-05907-x)] [Medline: [33403399](https://pubmed.ncbi.nlm.nih.gov/33403399/)]
17. Barnay T. Health, work and working conditions: a review of the European economic literature. *Eur J Health Econ* 2016 Jul;17(6):693-709 [doi: [10.1007/s10198-015-0715-8](https://doi.org/10.1007/s10198-015-0715-8)] [Medline: [26280132](https://pubmed.ncbi.nlm.nih.gov/26280132/)]
18. Nachreiner NM, Dagher RK, McGovern PM, Baker BA, Alexander BH, Gerberich SG. Successful return to work for cancer survivors. *AAOHN J* 2007 Jul;55(7):290-295 [doi: [10.1177/216507990705500705](https://doi.org/10.1177/216507990705500705)] [Medline: [17665826](https://pubmed.ncbi.nlm.nih.gov/17665826/)]
19. Rasmussen DM, Elverdam B. The meaning of work and working life after cancer: an interview study. *Psychooncology* 2008 Dec;17(12):1232-1238 [doi: [10.1002/pon.1354](https://doi.org/10.1002/pon.1354)] [Medline: [18623607](https://pubmed.ncbi.nlm.nih.gov/18623607/)]
20. Bilodeau K, Tremblay D, Durand MJ. Exploration of the contexts surrounding the implementation of an intervention supporting return-to-work after breast cancer in a primary care setting: starting point for an intervention development. *J Multidiscip Healthc* 2018 Feb 01;11:75-83 [FREE Full text] [doi: [10.2147/JMDH.S152947](https://doi.org/10.2147/JMDH.S152947)] [Medline: [29440910](https://pubmed.ncbi.nlm.nih.gov/29440910/)]
21. Bilodeau K, Tremblay D, Durand MJ. Gaps and delays in survivorship care in the return-to-work pathway for survivors of breast cancer-a qualitative study. *Curr Oncol* 2019 Jun;26(3):e414-e417 [FREE Full text] [doi: [10.3747/co.26.4787](https://doi.org/10.3747/co.26.4787)] [Medline: [31285687](https://pubmed.ncbi.nlm.nih.gov/31285687/)]
22. Cocchiara RA, Sciarra I, D'Egidio V, Sestili C, Mancino M, Backhaus I, et al. Returning to work after breast cancer: A systematic review of reviews. *Work* 2018;61(3):463-476 [doi: [10.3233/WOR-182810](https://doi.org/10.3233/WOR-182810)] [Medline: [30400123](https://pubmed.ncbi.nlm.nih.gov/30400123/)]
23. de Boer AGEM, Taskila TK, Tamminga SJ, Feuerstein M, Frings-Dresen MHW, Verbeek J. Interventions to enhance return-to-work for cancer patients. *Cochrane Database Syst Rev* 2015 Sep 25;2015(9):CD007569 [FREE Full text] [doi: [10.1002/14651858.CD007569.pub3](https://doi.org/10.1002/14651858.CD007569.pub3)] [Medline: [26405010](https://pubmed.ncbi.nlm.nih.gov/26405010/)]
24. Knott V, Zrim S, Shanahan EM, Anastassiadis P, Lawn S, Kichenadasse G, et al. Returning to work following curative chemotherapy: a qualitative study of return to work barriers and preferences for intervention. *Support Care Cancer* 2014 Dec;22(12):3263-3273 [doi: [10.1007/s00520-014-2324-y](https://doi.org/10.1007/s00520-014-2324-y)] [Medline: [25066834](https://pubmed.ncbi.nlm.nih.gov/25066834/)]
25. Tamminga SJ, Verbeek JHAM, Bos MMEM, Fons G, Kitzen JJEM, Plaisier PW, et al. Effectiveness of a hospital-based work support intervention for female cancer patients - a multi-centre randomised controlled trial. *PLoS One* 2013 May 22;8(5):e63271 [doi: [10.1371/journal.pone.0063271](https://doi.org/10.1371/journal.pone.0063271)] [Medline: [23717406](https://pubmed.ncbi.nlm.nih.gov/23717406/)]
26. Tamminga SJ, Wolvers MDJ, Greidanus MA, Zaman AGNM, Braspenning AM, Frings-Dresen MHW, et al. Employees diagnosed with cancer: Current perspectives and future directions from an employer's point of view. *J Occup Rehabil* 2019 Jun;29(2):472-474 [FREE Full text] [doi: [10.1007/s10926-018-9802-x](https://doi.org/10.1007/s10926-018-9802-x)] [Medline: [30039313](https://pubmed.ncbi.nlm.nih.gov/30039313/)]
27. Bilodeau K, Tremblay D, Durand MJ. Exploration of return-to-work interventions for breast cancer patients: a scoping review. *Support Care Cancer* 2017 Jun;25(6):1993-2007 [doi: [10.1007/s00520-016-3526-2](https://doi.org/10.1007/s00520-016-3526-2)] [Medline: [28054145](https://pubmed.ncbi.nlm.nih.gov/28054145/)]

28. Campagna M, Loscerbo R, Pilia I, Meloni F. Return to work of breast cancer survivors: Perspectives and challenges for occupational physicians. *Cancers (Basel)* 2020 Feb 04;12(2):355 [FREE Full text] [doi: [10.3390/cancers12020355](https://doi.org/10.3390/cancers12020355)] [Medline: [32033165](https://pubmed.ncbi.nlm.nih.gov/32033165/)]
29. Soins de santé communautaires de première ligne. Instituts de recherche en santé du Canada. URL: <https://cihr-irsc.gc.ca/f/43626.html> [accessed 2022-04-12]
30. Fadhlaoui A, Mrad H, Vinette B, Bilodeau K. L'expérience de transition vers le retour au travail de survivantes du cancer du sein âgées de moins de 50 ans après la fin des traitements. *Can Oncol Nurs J* 2021 Nov 01;31(4):386-392 [FREE Full text] [doi: [10.5737/23688076314386392](https://doi.org/10.5737/23688076314386392)] [Medline: [34786456](https://pubmed.ncbi.nlm.nih.gov/34786456/)]
31. Groupe de médecine de famille (GMF), GMF-U et super-clinique (GMF accès-réseau). Gouvernement du Québec. URL: <https://www.msss.gouv.qc.ca/professionnels/soins-et-services/groupe-de-medecine-de-famille-gmf-et-super-cliniques-gmf-reseau/> [accessed 2022-04-12]
32. Ensemble, en réseau, pour vaincre le cancer: Plan directeur en cancérologie. Direction québécoise de cancérologie. 2013. URL: <http://publications.msss.gouv.qc.ca/acrobat/f/documentation/2013/13-902-02W.pdf> [accessed 2022-04-12]
33. McCabe MS, Bhatia S, Oeffinger KC, Reaman GH, Tyne C, Wollins DS, et al. American Society of Clinical Oncology statement: achieving high-quality cancer survivorship care. *J Clin Oncol* 2013 Feb 10;31(5):631-640 [doi: [10.1200/JCO.2012.46.6854](https://doi.org/10.1200/JCO.2012.46.6854)] [Medline: [23295805](https://pubmed.ncbi.nlm.nih.gov/23295805/)]
34. Fitch MI. Supportive care framework. *Can Oncol Nurs J* 2008;18(1):6-24 [doi: [10.5737/1181912x181614](https://doi.org/10.5737/1181912x181614)] [Medline: [18512565](https://pubmed.ncbi.nlm.nih.gov/18512565/)]
35. Boyages J, Kalfa S, Xu Y, Koelmeyer L, Mackie H, Viveros H, et al. Worse and worse off: the impact of lymphedema on work and career after breast cancer. *Springerplus* 2016 May 17;5:657 [FREE Full text] [doi: [10.1186/s40064-016-2300-8](https://doi.org/10.1186/s40064-016-2300-8)] [Medline: [27330922](https://pubmed.ncbi.nlm.nih.gov/27330922/)]
36. Alfano CM, Kent EE, Padgett LS, Grimes M, de Moor JS. Making cancer rehabilitation services work for cancer patients: Recommendations for research and practice to improve employment outcomes. *PM R* 2017 Sep;9(9S2):S398-S406 [FREE Full text] [doi: [10.1016/j.pmrj.2017.06.019](https://doi.org/10.1016/j.pmrj.2017.06.019)] [Medline: [28942911](https://pubmed.ncbi.nlm.nih.gov/28942911/)]
37. Jagsi R, Hawley ST, Abrahamse P, Li Y, Janz NK, Griggs JJ, et al. Impact of adjuvant chemotherapy on long-term employment of survivors of early-stage breast cancer. *Cancer* 2014 Jun 15;120(12):1854-1862 [FREE Full text] [doi: [10.1002/cncr.28607](https://doi.org/10.1002/cncr.28607)] [Medline: [24777606](https://pubmed.ncbi.nlm.nih.gov/24777606/)]
38. Johnsson A, Fornander T, Rutqvist LE, Vaez M, Alexanderson K, Olsson M. Predictors of return to work ten months after primary breast cancer surgery. *Acta Oncol* 2009;48(1):93-98 [doi: [10.1080/02841860802477899](https://doi.org/10.1080/02841860802477899)] [Medline: [18937082](https://pubmed.ncbi.nlm.nih.gov/18937082/)]
39. Peugniez C, Fantoni S, Leroyer A, Skrzypczak J, Duprey M, Bonnetterre J. Return to work after treatment for breast cancer: single center experience in a cohort of 273 patients. *Bull Cancer* 2011 Jul;98(7):E69-E79 [FREE Full text] [doi: [10.1684/bdc.2011.1401](https://doi.org/10.1684/bdc.2011.1401)] [Medline: [21684837](https://pubmed.ncbi.nlm.nih.gov/21684837/)]
40. Lindbohm M, Taskila T, Kuosma E, Hietanen P, Carlsen K, Gudbergsson S, et al. Work ability of survivors of breast, prostate, and testicular cancer in Nordic countries: a NOCWO study. *J Cancer Surviv* 2012 Mar;6(1):72-81 [doi: [10.1007/s11764-011-0200-z](https://doi.org/10.1007/s11764-011-0200-z)] [Medline: [22042662](https://pubmed.ncbi.nlm.nih.gov/22042662/)]
41. Loisel P. Work disability: It is not just the "lesion". In: Feuerstein M, editor. *Work and Cancer Survivors*. New York, NY: Springer; 2009:93-103
42. Craig P, Dieppe P, Macintyre S, Michie S, Nazareth I, Petticrew M, Medical Research Council Guidance. Developing and evaluating complex interventions: the new Medical Research Council guidance. *BMJ* 2008 Sep 29;337:a1655 [FREE Full text] [doi: [10.1136/bmj.a1655](https://doi.org/10.1136/bmj.a1655)] [Medline: [18824488](https://pubmed.ncbi.nlm.nih.gov/18824488/)]
43. Sidani S, Braden CJ. *Design, Evaluation, and Translation of Nursing Interventions*. Hoboken, New Jersey, United States: John Wiley & Sons; 2011.
44. Vennik FD, van de Bovenkamp HM, Putters K, Grit KJ. Co-production in healthcare: rhetoric and practice. *International Review of Administrative Sciences* 2015 Jun 29;82(1):150-168 [FREE Full text] [doi: [10.1177/0020852315570553](https://doi.org/10.1177/0020852315570553)]
45. Safin S, Dorta T. Unfolding laypersons creativity through social VR: A case study. 2020 Presented at: 38th eCAADe Conference; September 16-18, 2020; Berlin, Germany p. 355-364
46. Côté V, Bélanger L, Gagnon C. Le design au service de l'expérience patient. *Sciences du Design* 2017;2(6):54-64 [FREE Full text] [doi: [10.3917/sdd.006.0054](https://doi.org/10.3917/sdd.006.0054)]
47. Reason P, Bradbury H. *The SAGE Handbook of Action Research: Participative Inquiry and Practice*. Newcastle upon Tyne, United Kingdom: SAGE; 2007.
48. Dorta T, Safin S, Boudhraâ S, Marchand EB. Co-designing in social VR: Process awareness and suitable representations to empower user participation. 2019 Presented at: 24th International Conference of the Association for Computer-Aided Architectural Design Research in Asia (CAADRIA) 2019; April 15-18, 2019; Wellington, New Zealand p. 141-150 URL: http://papers.cumincad.org/data/works/att/caadria2019_240.pdf
49. Stringer ET. *Action Research*. Thousand Oaks, California, United States: SAGE Publications; 2013.
50. Jagosh J, Macaulay AC, Pluye P, Salsberg J, Bush PL, Henderson J, et al. Uncovering the benefits of participatory research: implications of a realist review for health research and practice. *Milbank Q* 2012 Jun;90(2):311-346 [FREE Full text] [doi: [10.1111/j.1468-0009.2012.00665.x](https://doi.org/10.1111/j.1468-0009.2012.00665.x)] [Medline: [22709390](https://pubmed.ncbi.nlm.nih.gov/22709390/)]

51. Koch T, Mann S, Kralik D, van Loon AM. Reflection: Look, think and act cycles in participatory action research. *J Res Nurs* 2005 May 01;10(3):261-278 [doi: [10.1177/174498710501000304](https://doi.org/10.1177/174498710501000304)]
52. Patton MQ. *Qualitative Research & Evaluation Methods: Integrating Theory and Practice*. Thousand Oaks, California, United States: SAGE Publications; 2014.
53. Leask CF, Sandlund M, Skelton DA, Altenburg TM, Cardon G, Chinapaw MJM, GrandStand, Safe Step and Teenage Girls on the Move Research Groups. Framework, principles and recommendations for utilising participatory methodologies in the co-creation and evaluation of public health interventions. *Res Involv Engagem* 2019 Jan 09;5:2 [FREE Full text] [doi: [10.1186/s40900-018-0136-9](https://doi.org/10.1186/s40900-018-0136-9)] [Medline: [30652027](https://pubmed.ncbi.nlm.nih.gov/30652027/)]
54. Morrison C, Dearden A. Beyond tokenistic participation: using representational artefacts to enable meaningful public participation in health service design. *Health Policy* 2013 Oct;112(3):179-186 [FREE Full text] [doi: [10.1016/j.healthpol.2013.05.008](https://doi.org/10.1016/j.healthpol.2013.05.008)] [Medline: [23786993](https://pubmed.ncbi.nlm.nih.gov/23786993/)]
55. Dorta T, Kinayoglu G, Hoffmann M. Hyve-3D and the 3D cursor: Architectural co-design with freedom in virtual reality. *International Journal of Architectural Computing* 2016 Apr 21;14(2):87-102 [doi: [10.1177/1478077116638921](https://doi.org/10.1177/1478077116638921)]
56. Dorta T, Kalay YE, Lesage A, Pérez E. Elements of design conversation in the interconnected HIS. *International Journal of Design Sciences and Technology* 2011 Jan;18(2):65-80
57. Patton MQ. *Qualitative research*. In: *Encyclopedia of Statistics in Behavioral Science*. Hoboken, New Jersey, United States: John Wiley & Sons; 2005.
58. Sekhon M, Cartwright M, Francis JJ. Acceptability of healthcare interventions: an overview of reviews and development of a theoretical framework. *BMC Health Serv Res* 2017 Jan 26;17(1):88 [FREE Full text] [doi: [10.1186/s12913-017-2031-8](https://doi.org/10.1186/s12913-017-2031-8)] [Medline: [28126032](https://pubmed.ncbi.nlm.nih.gov/28126032/)]
59. Miles MB, Huberman AM, Saldaña J. *Qualitative Data Analysis: A Methods Sourcebook*, 3rd Edition. Thousand Oaks, California, United States: SAGE Publications; 2014.
60. Caron M. Modélisation d'une intervention visant à favoriser le processus de réappropriation du travail des survivantes du cancer du sein. Université de Sherbrooke. 2020 Jan. URL: https://savoirs.usherbrooke.ca/bitstream/handle/11143/16400/Caron_Maryse_PhD_2020.pdf?sequence=1&isAllowed=y [accessed 2022-04-12]
61. Fassier JB, Lamort-Bouché M, Broc G, Guittard L, Péron J, Rouat S, et al. Developing a return to work intervention for breast cancer survivors with the Intervention Mapping Protocol: Challenges and opportunities of the needs assessment. *Front Public Health* 2018 Feb 23;6:35 [FREE Full text] [doi: [10.3389/fpubh.2018.00035](https://doi.org/10.3389/fpubh.2018.00035)] [Medline: [29527521](https://pubmed.ncbi.nlm.nih.gov/29527521/)]
62. Greidanus MA, de Boer AGEM, Tiedtke CM, Frings-Dresen MHW, de Rijk AE, Tamminga SJ. Supporting employers to enhance the return to work of cancer survivors: development of a web-based intervention (MiLES intervention). *J Cancer Surviv* 2020 Apr;14(2):200-210 [FREE Full text] [doi: [10.1007/s11764-019-00844-z](https://doi.org/10.1007/s11764-019-00844-z)] [Medline: [31938966](https://pubmed.ncbi.nlm.nih.gov/31938966/)]
63. Paltrinieri S, Fugazzaro S, Bertozzi L, Bassi MC, Pellegrini M, Vicentini M, et al. Return to work in European cancer survivors: a systematic review. *Support Care Cancer* 2018 Sep;26(9):2983-2994 [doi: [10.1007/s00520-018-4270-6](https://doi.org/10.1007/s00520-018-4270-6)] [Medline: [29845421](https://pubmed.ncbi.nlm.nih.gov/29845421/)]
64. Nastasia I, Durand MJ, Coutu MF, Collinge C, Cibotaru A. Pratiques des milieux de travail pour assurer un retour en emploi sain et durable. Institut de recherche Robert-Sauvé en santé et en sécurité du travail. 2017. URL: <http://www.irsst.qc.ca/media/documents/PubIRSST/R-983.pdf> [accessed 2022-04-12]
65. Murray CJL, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, et al. Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012 Dec 15;380(9859):2197-2223 [doi: [10.1016/S0140-6736\(12\)61689-4](https://doi.org/10.1016/S0140-6736(12)61689-4)] [Medline: [23245608](https://pubmed.ncbi.nlm.nih.gov/23245608/)]
66. OECD. *Sickness, Disability and Work: Breaking the Barriers: A Synthesis of Findings across OECD Countries*. Paris, France: Organisation for Economic Co-operation and Development; 2010.
67. Côté M, Boyle É. Pour contrer la pénurie de main-d'œuvre : pourquoi pas les personnes handicapées? *Ordre des conseillers en ressources humaines agréés*. 2011 Nov 07. URL: <https://ordrecrha.org/ressources/revue-rh/archives/pour-contrer-la-penurie-de-main-d-oeuvre-pourquoi-pas-les-personnes-handicapees> [accessed 2022-04-12]
68. Stratégie de recherche axée sur le patient. Instituts de recherche en santé du Canada. URL: <http://www.cihr-irsc.gc.ca/f/41204.html> [accessed 2022-04-12]
69. Bate P, Robert G. Experience-based design: from redesigning the system around the patient to co-designing services with the patient. *Qual Saf Health Care* 2006 Oct;15(5):307-310 [FREE Full text] [doi: [10.1136/qshc.2005.016527](https://doi.org/10.1136/qshc.2005.016527)] [Medline: [17074863](https://pubmed.ncbi.nlm.nih.gov/17074863/)]
70. Chowdhury J. Hacking Health: Bottom-up innovation for healthcare. *Technology Innovation Management Review* 2012 Jul;2(7):31-35 [FREE Full text] [doi: [10.22215/timreview/579](https://doi.org/10.22215/timreview/579)]
71. Locock L, Robert G, Boaz A, Vougioukalou S, Shuldham C, Fielden J, et al. Testing accelerated experience-based co-design: a qualitative study of using a national archive of patient experience narrative interviews to promote rapid patient-centred service improvement. *Health Services and Delivery Research* 2014 Feb;2(4):1-150 [doi: [10.3310/hsdr02040](https://doi.org/10.3310/hsdr02040)] [Medline: [25642558](https://pubmed.ncbi.nlm.nih.gov/25642558/)]
72. Donetto S, Tsianakas V, Robert G. Using experience-based Co-design (EBCD) to improve the quality of healthcare: mapping where we are now and establishing future directions. ResearchGate. 2014. URL: <https://www.researchgate.net/publication/>

- [262198403 Using Experience-based Co-design EBCD to improve the quality of healthcare mapping where we are now and establishing future directions](#) [accessed 2022-04-12]
73. Mattison CA, Moat KA, Waddell K, Lavis JN. Optimizing patient and family transitions from cancer treatment to primary- and community-care supports in Canada. McMaster Health Forum. 2018 Mar 27. URL: https://www.mcmasterforum.org/docs/default-source/product-documents/evidence-briefs/cancer-care-transitions-eb.pdf?sfvrsn=844354d5_2 [accessed 2022-04-12]
74. Tremblay D, Prady C, Bilodeau K, Touati N, Chouinard MC, Fortin M, et al. Optimizing clinical and organizational practice in cancer survivor transitions between specialized oncology and primary care teams: a realist evaluation of multiple case studies. BMC Health Serv Res 2017 Dec 16;17(1):834 [FREE Full text] [doi: [10.1186/s12913-017-2785-z](https://doi.org/10.1186/s12913-017-2785-z)] [Medline: [29246224](https://pubmed.ncbi.nlm.nih.gov/29246224/)]
75. Recklitis CJ, Syrjala KL. Provision of integrated psychosocial services for cancer survivors post-treatment. Lancet Oncol 2017 Jan;18(1):e39-e50 [FREE Full text] [doi: [10.1016/S1470-2045\(16\)30659-3](https://doi.org/10.1016/S1470-2045(16)30659-3)] [Medline: [28049576](https://pubmed.ncbi.nlm.nih.gov/28049576/)]
76. Guest G, Bunce A, Johnson L. How many interviews are enough?: An experiment with data saturation and variability. Field methods 2006 Feb 01;18(1):59-82 [doi: [10.1177/1525822x05279903](https://doi.org/10.1177/1525822x05279903)]

Abbreviations

RTW: return to work

Edited by T Leung: This paper was externally peer-reviewed by the Quebec Rehabilitation Research Network (Réseau Provincial de Recherche en Adaptation-Réadaptation, REPAR) - Réseau thématique soutenu par le FRQS - Programme 1.1 Recherche clinique 2020-2021 (Montréal, Canada). See the Multimedia Appendix for the peer-review report; Submitted 25.02.22; accepted 29.03.22; published 22.04.22.

Please cite as:

Bilodeau K, Gouin MM, Lecours A, Lederer V, Durand MJ, Kilpatrick K, Lepage D, Ladouceur-Deslauriers L, Dorta T
Acceptability and Feasibility of a Return-to-Work Intervention for Posttreatment Breast Cancer Survivors: Protocol for a Co-design and Development Study

JMIR Res Protoc 2022;11(4):e37009

URL: <https://www.researchprotocols.org/2022/4/e37009>

doi: [10.2196/37009](https://doi.org/10.2196/37009)

PMID: [35451972](https://pubmed.ncbi.nlm.nih.gov/35451972/)

©Karine Bilodeau, Marie-Michelle Gouin, Alexandra Lecours, Valérie Lederer, Marie-José Durand, Kelley Kilpatrick, David Lepage, Lauriane Ladouceur-Deslauriers, Tomas Dorta. Originally published in JMIR Research Protocols (<https://www.researchprotocols.org>), 22.04.2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Research Protocols, is properly cited. The complete bibliographic information, a link to the original publication on <https://www.researchprotocols.org>, as well as this copyright and license information must be included.