

From pathway to products: Canadian achievements in plant specialized metabolism

Karen Cristine Gonçalves dos Santos^{a,b}, Fatima Awwad^{a,b}, Natacha Mérindol^{a,b}, and Isabel Desgagné-Penix^{a,b}

^aDepartment of Chemistry, Biochemistry and Physics, Université du Québec à Trois-Rivières, Trois-Rivières, QC, Canada; ^bPlant Biology Research Group, Université du Québec à Trois-Rivières, Trois-Rivières, QC, Canada

Corresponding author: Isabel Desgagné-Penix (email: Isabel.Desgagne-Penix@uqtr.ca)

Abstract

Canada has made significant contributions to the field of plant biochemistry, with numerous researchers actively focusing on elucidating the biosynthetic pathways of plant specialized metabolites and producing these compounds in heterologous systems, such as bacteria, yeast, or other plant species. The review aims to highlight the strengths of Canadian research in this domain over the last three decades. It will describe advances in pathway elucidation, enzyme characterization, and production of enzymes and metabolites in heterologous systems, particularly in the areas of alkaloids, terpenoids, and phenolic compounds. Canadian researchers have not only made pivotal scientific discoveries but have also ensured the continuity of scientific excellence by mentoring new generations of principal investigators in plant specialized metabolites. These advances warrant recognition and financial support to retain future talent and to maintain Canada's leadership in scientific progress on the global stage.

Key words: biochemical pathways, scientific innovation, biotechnology, metabolic engineering, alkaloids


1. Introduction

Plants synthesize a wide arsenal of metabolites to protect themselves against herbivores, pathogens, and parasites, and abiotic stresses, or to attract pollinators. Humans use plants as food and fuel, and exploit plants specialized metabolites for medicines, cosmetics, and nutritional supplements. One notable example of a phytometabolite with pharmaceutical properties is TaxolTM (generic name paclitaxel), a diterpene that was discovered in 1967 from bark extracts of the North American tree *Taxus brevifolia* (Pacific yew) (Weaver 2014). Paclitaxel was first approved by the USA's Food and Drug Administration (FDA) for the treatment of ovarian cancer in 1992, and is now also approved for the treatment of breast and non-small cell lung cancer, as well as Kaposi's sarcoma (Weaver 2014). Another example is capsaicin (*trans*-8-methyl-N-vanillyl-6-nonenamide), a phenolic alkaloid from *Capsicum* spp. (e.g., chili peppers) native to the American continent (Costa et al. 2022) that brings spiciness to food by causing a burning sensation in the mucous membranes of the mouth. Besides its role in gastronomy, the compound is used in medicine and has been approved for pain relief by both the FDA and the European Medicines Agency (Sharma et al. 2013; Weaver 2014; Chung and Campbell 2016). Both metabolites have biological relevance: paclitaxel acts against parasites and fungal attacks in *T. brevifolia* (Elmer et al. 1994), and capsaicin repels mammal herbivores (Tewksbury and Nabhan 2001).

In Canada, several research groups are dedicated to elucidating how plants produce specialized metabolites of interest. For instance, in 2013, a multi-laboratory effort led to sequencing, de novo assembly, and annotation of the transcriptomes from 20 plant species of medicinal importance, enabling the identification of genes in biosynthetic pathways of specialized metabolites, including valerenic acid, benzyloquinoline alkaloids (BIAs), and the polyketide hypericin (Xiao et al. 2013). A quick search on Scopus shows that, between 2019 and the first half of 2024, Canada was among the top 10 countries publishing on plant metabolism elucidation and plant metabolic engineering, with Canadian researchers contributing 142 research and review articles among 4538 published worldwide.

In this review, we provide an overview of the advances led by principal investigators (PIs) active in the domain of plant specialized metabolism across Canada. Due to space constraints, we focused on PIs whose research centres on pathway elucidation and/or bioengineering of plant alkaloids, terpenoids, and phenolic compounds, excluding phytohormones. We apologize if we have inadvertently omitted any PIs; our intention was not to offend anyone but to highlight the significant contribution of Canadian researchers. We have done our best to include everyone and would appreciate it if you could inform us of any omissions so we can include them in a future revision of this review. Additionally, we listed researchers with at least three publications in the

Fig. 1. Alkaloids and their plants of origin investigated in Canada, along with their main precursor: aromatic amino acids. Numbers in parenthesis indicate the researchers that investigate the topic: (1.1) Vincenzo De Luca; (1.2) Peter J. Facchini; (1.3) Thu-Thuy T. Dang; (1.4) Yang Qu; (1.5) Vincent JJ Martin; (1.6) Isabel Desgagné-Penix. *Narcissus pseudonarcissus*' drawing was done with Krita 5.2.9. The following images were obtained from Wikimedia Commons: *Camptotheca acuminata*: Auckland Museum (CC BY 4.0) <https://commons.wikimedia.org/w/index.php?curid=65765508>; *Ephedra sinica*: By alexlomas (Flickr, CC BY 2.0) <https://commons.wikimedia.org/w/index.php?curid=6343381>; *Papaver somniferum*: By Rachelhs (CC0) <https://commons.wikimedia.org/w/index.php?curid=34838409>; *Rauwolfia serpentina*: By MAPeFY (CC BY-SA 4.0) <https://commons.wikimedia.org/w/index.php?curid=80304394>; *Vinca minor*: By Ryan Kaldari, <https://commons.wikimedia.org/w/index.php?curid=6369683>.



filed within the last 5 years. Major breakthroughs in the field over the past 30 years, as well as synthetic biology innovations for producing plant specialized metabolites in heterologous systems, will be described.

2. Investigating alkaloids

Alkaloids, derived from amino acids such as phenylalanine, tryptophan, and tyrosine, form the one of the largest groups of specialized metabolites. This category includes the neuroactive caffeine (McLellan et al. 2016), the antitumoral vinblastine and vincristine (Dhyani et al. 2022), the anti-gout agent colchicine (Dhyani et al. 2022; Stamp et al. 2024), and the acetylcholinesterase inhibitor galanthamine (Kaur et al. 2022).

Canadian research teams have significantly advanced the understanding of monoterpene indole alkaloids (MIAs) and BIAs. Their work highlights the enzymes involved in biosynthesis, plant defense roles, biological activities, medicinal potential, and heterologous production meth-

ods. International efforts have contributed to the elucidation of the MIA pathway, with several reviews providing comprehensive insights (O'Connor and Maresh 2006; St-Pierre et al. 2013; De Luca et al. 2014; Pan et al. 2015; Salim et al. 2023).

Meanwhile, in Canada, BIA research has primarily focused on poppy species and Amaryllidaceae. Global studies predominantly investigate alkaloids from *Papaver somniferum* (opium poppy), but species of other orders have also been investigated, with advances in gene discovery, expression, and protein characterization (Menendez-Perdomo and Facchini 2018; Singh et al. 2019; Georgiev et al. 2020; Desgagné-Penix 2021; Aghaali and Naghavi 2024; Aghaali et al. 2024; Liyanage et al. 2024).

This section reviews advancements in MIA metabolism led by Professors Vincenzo De Luca and Yang Qu, contributions by Professors Peter Facchini and Vincent J.J. Martin on poppy alkaloids, Pr. Isabel Desgagné-Penix on Amaryllidaceae alkaloids, and Pr. Thu-Thuy T. Dang's research on cytochrome P450s in alkaloid biosynthesis pathways (Fig. 1).

2.1. Vincenzo de Luca

The legacy of Pr. Vincenzo de Luca (Brock University) includes his team's major discoveries on alkaloids biosynthesis and training of the next generation of Canadian plant biochemists. His group focused on the elucidation of the biosynthesis pathway of the MIA vindoline and vinblastine, derived from tryptophan and secologanin (Salim et al. 2023), mainly in *Catharanthus roseus* (Madagascar periwinkle). This massive complex pathway of 31 steps was recently elucidated in its entirety, with 17 enzymes being discovered and characterized by Pr. De Luca's team and other three being further characterized by them (Fig. 2; Kulagina et al. 2022).

Pr. de Luca's team also researched MIA biosynthesis in Indian snakeroot (*Rauvolfia serpentina*), characterizing the role of γ -tocopherol-like *N*-methyltransferases (Cazares-Flores et al. 2016; Levac et al. 2016) in ajmaline biosynthesis, and uncovered the transporters involved in the export of vincamine in *Vinca minor* (periwinkle) (Demessie et al. 2017). His group has also reported on grapes and their glucosyltransferases, flavonols, and esters (Wang and De Luca 2005; Hall and De Luca 2007; Hall et al. 2011a, 2012).

In addition to the research achievements describe above, the training of a new generation of accomplished researchers in the area is also part of Pr. De Luca's legacy. Several big and emerging names in the plant specialized metabolites research in Canada passed by his laboratory, including Pr. Peter Facchini, Pr. Yang Qu, and Pr. Mehran Dastmalchi, whose research works are discussed below.

2.2. Peter J. Facchini

Pr. Peter J. Facchini's (Calgary University) research on the BIA pathway in *Papaver somniferum* (opium poppy) started with the characterization of tyrosine/dihydroxyphenylalanine decarboxylases (TYDCs) and the profiling of their expression in different tissues (Facchini and De Luca 1994, 1995a, 1995b; Facchini et al. 1996a). In the following years, his group tackled the identification and characterization of enzymes in this pathway, including transporters, as well as the expression profiles of the genes encoding them, including berberine bridge enzyme 1 (Facchini et al. 1996b), TYDCs (Facchini et al. 1998), tyramine *N*-(hydroxycinnamoyl)transferase (Yu and Facchini 1999), norcoclaurine synthase (Samanani and Facchini 2001, 2002; Samanani et al. 2004; Luk et al. 2007; Lee and Facchini 2010), and many more (Bird et al. 2003; Liscombe and Facchini 2007; Hagel and Facchini 2010; Lee and Facchini 2011; Dang and Facchini 2012; Hagel et al. 2012; Beaudoin and Facchini 2013; Dang et al. 2015; Chen et al. 2018; Dastmalchi et al. 2019a; Li et al. 2020).

Pr. Facchini's team investigated the biosynthesis of BIA in other species as well, including *Thalictrum flavum*, *Eschscholzia californica*, *Papaver bracteatum*, and *Glaucium flavum* (yellow meadow rue, California poppy, Iranian poppy, and yellow horned poppy, respectively), which allowed the study of the evolution of norcoclaurine synthase, berberine-bridge enzyme, and *O*-methyltransferases (Liscombe et al. 2005), and the identification of new *N*-methyltransferases, including the

first isolated enzyme in the pavine alkaloid branch pathway (Liscombe et al. 2009; Samanani et al. 2005).

His group also performed a transcriptomic study of *Lophophora williamsii* and elucidated the biosynthetic pathway of tetrahydroisoquinoline and phenethylamine alkaloids, including mescaline and pellotine (Watkins et al. 2023). Facchini's team also characterized aromatic aminotransferase implicated in ephedrine and associated alkaloids from *Ephedra sinica* and *Catha edulis* (ephedra and khat) (Kilpatrick et al. 2016; Morris et al. 2018). Through Pr. Facchini's collaborations with the structural biologist Pr. Kenneth Ng, the crystal structures of *P. somniferum*'s codeine reductase (Carr et al. 2021), *T. flavum*'s pavine *N*-methyltransferase (Torres et al. 2016), and *G. flavum*'s tetrahydroprotoberberine *N*-methyltransferase (Lang et al. 2019) were resolved, contributing to our overall understanding of substrate recognition by methyltransferases and reductases.

Finally, Pr. Facchini also trained recently established researchers in the plant alkaloid research in Canada. For instance, Pr. David Liscombe's (adjunct assistant professor at Brock University, Ontario, and researcher at Vineland Research and Innovation Centre) recent study of the invasive species *Vincetoxicum rossicum* led to an update on the biosynthesis pathway of phenanthroindolizidine alkaloid (Kempthorne et al. 2024). Pr. Mehran Dastmalchi, Pr. Thu-Thuy T. Dang, and Pr. Isabel Desgagné-Penix, who are discussed below, are also among the specialized metabolism experts that were trained by Pr. Facchini.


2.3. Thu-Thuy T. Dang

Pr. Thu-Thuy T. Dang recently established her laboratory at University of British Columbia (UBC), where her team uses a combination of genomics and biochemistry to understand plant alkaloid biosynthesis. The group published abundantly on the topic (Bui et al. 2023; Kwan et al. 2023; Nguyen and Dang 2021, 2022; Nunes et al. 2022), as well as with Canadian and international collaborators (Dang et al. 2012, 2017; Tatsis et al. 2017; Stavrinides et al. 2018; Carqueijeiro et al. 2018; Bui et al. 2023). Additionally, Pr. Thu-Thuy T. Dang's team used gene expression analysis to identify CYP encoding genes in *Camptotheca acuminata* (Happy tree), and enzymatic assays to show that the corresponding enzymes were able to oxidize camptothecin, a quinoline alkaloid (Nguyen et al. 2021), resulting in a patent (Dang et al. 2021). The patented innovation highlights isolated cytochrome P450 sequences, as well as synthetic biology application in plants, highlighting the role of specific cytochrome P450s in plant defense and isoprenoid compound synthesis. The group was also the first to identify and characterize CYP responsible for the formation of two spirooxindole alkaloids using comparative transcriptomics and enzymatic assays, namely MsCYP72056 from *Mitragyna speciosa* (kratom) (Nguyen et al. 2023).

2.4. Yang Qu

Pr. Yang Qu previously worked with Pr. Dae-Kyun Ro (University of Calgary) on the biosynthesis of natural rubber in lettuce (Chakrabarty et al. 2015; Qu et al. 2015b; Barnes et al. 2021), and with Pr. Vincenzo de Luca on MIA metabolism

Fig. 2. Enzymes from *Catharanthus roseus* monoterpene indole alkaloid pathway. Filled maple leaves indicate steps characterized by Pr. De Luca's research teams; hollow maple leaves indicate steps characterized by teams outside Canada, but that were further characterized by Canadian teams. Numbers inside maple leaves indicate studies that characterized the enzymes. (1) De Luca et al. (1989); (2) Salim et al. (2014); (3) Asada et al. (2013); (4) Murata and De Luca (2005); (5) Qu et al. (2018a, 2018b); (6) Qu et al. (2019); (7) St-Pierre and De Luca (1995); (8) Levac et al. (2008); (9) Qu et al. (2015a); (10) De Carolis and De Luca (1993, 1994); De Carolis et al. (1990); Vazquez-Flota et al. (1997); (11) Power et al. (1990). MVA: mevalonic acid; GPP: geraniol pyrophosphate; GES: geraniol synthase; TDC: tryptophan decarboxylase; G8H: geraniol 8-hydroxylase; CPR: cytochrome P450 reductase; CYB5: cytochrome B5; CYPADH: cytochrome P450 alcohol dehydrogenase; 7DLGT: 7-deoxyloganetic acid transferase; 7DLH: 7-deoxyloganic acid hydroxylase; LAMT: loganic acid methyltransferase; SLS: secologanin synthase; STR: strictosidine synthase; SGD: strictosidine 8HGO: 8-hydroxygeraniol oxidoreductase; ISY: iridoid synthase; IO: iridoid oxidase; SGD: strictosidine β -D-glucosidase; GS: geissoschizine synthase; GO: geissoschizine oxidase; Redox1/2: protein redox 1/2; SAT: stemmadenine-O-acetyltransferase; PAS: precondylocarpine acetate synthase; DPAS: dehydroprecondylocarpine acetate synthase; CS/TS: catharanthine/tabersonine synthase; T16H: tabersonine 16-hydroxylase; 16OMT: 16-hydroxytabersonine O-methyltransferase; T3O: tabersonine 3-oxygenase; T3R: tabersonine 3-reductase; NMT: 3-hydroxy-16-methoxy-2,3-dihydroxytabersonine-N-methyltransferase; D4H: desacetoxyvindoline-4-hydroxylase; DAT: deacetylvindoline-4-O-acetyltransferase; PRX1: class III peroxidase. The colored boxes lines indicate strictosidine (blue), tabersonine/catharanthine (yellow) and vindoline (red) modules. Adapted from Zhang et al. (2022).



(Demessie et al. 2017; Kidd et al. 2019; Qu et al. 2018b, 2019; Williams et al. 2019). In De Luca's lab, Pr. Qu bioengineered yeast to produce vindoline (Qu et al. 2015a). He established his laboratory at the University of New Brunswick (New Brunswick) focusing on natural medicine biosynthesis, including alkaloids and bibenzyls. Among his works on the

elucidation of biosynthetic pathways of specialized metabolites is the characterization of vindolinine synthase, which diverts tabersonine from vinblastine metabolism to the formation of 19S-vindolinine, 19R-vindolinine, and venalstonine (Eng et al. 2022). His group also elucidated the biosynthesis pathway of the analgesic mitragynine, from kratom, through

the transcriptomic study of the latter and other species of *Rubiaceae*, leading to the identification of reductases and enol methyltransferases that completed the pathway (Kim et al. 2023).

Finally, his team recently identified and characterized the enzymes catalyzing the last two unknown steps of the antiarrhythmic MIA ajmaline biosynthesis in Indian snakeroot, namely vomilenine 1,2(R)-reductase and 1,2-dihydrovomilenine 19,20(S)-reductase, as well as a new isoform of the sarpagan bridge enzyme and two new isoforms of the 17-O-acetylnorajmaline acetyl esterase (Guo et al. 2024). The group succeeded in applying these findings for the de novo biosynthesis of ajmaline in yeast. They also recently bioengineered yeast to produce multiple terpenoids and alkaloids, mainly vinblastine and other *Catharanthus* metabolites (Shahsavarani et al. 2023; Guo et al. 2024).

2.5. Vincent JJ Martin

Pr. Vincent JJ Martin is interested in bioengineering microorganisms to produce compounds of interest such as BIAS at Concordia University. He collaborated with Pr. Facchini on opioids and with Pr. Jay Keasling (University of California) on cannabinoid pathway elucidation and heterologous production. Pr. Martin's recent work includes synthetic biology approaches to optimize microbial platforms, mainly yeast (Pyne et al. 2020, 2023b; Dykstra et al. 2023), as well as pathway elucidation efforts for multiple BIA compounds. For instance, his team screened plant O-methyltransferases, enabling de novo synthesis of liensinine (Pyne et al. 2023a), and designed an enzyme-coupled biosensor for reticuline upstream intermediate L-3,4-dihydroxyphenylalanine (Fossati et al. 2014). Pr. Martin is also a co-director of Concordia university's Centre for Applied Synthetic Biology, also known as the Concordia Genome Foundry.

2.6. Isabel Desgagné-Penix

Lastly, our lab, at the Université du Québec à Trois-Rivières, has also made strides in medicinal plants specialized metabolism bioactivity and pathway elucidation, focusing on *Amaryllidaceae* alkaloids (AAs), cannabinoids, and vanillin. Early and more recent de novo transcriptomic studies of *Narcissus pseudonarcissus* "King Alfred", *N. papyraceus*, *Leucojum aestivum*, and *Crinum x powellii*, led to the identification of candidate genes involved in the biosynthesis of AAs (Singh and Desgagne-Penix 2017; Hotchandani et al. 2019; Tousignant et al. 2022; Koirala et al. 2023). Following identification of candidates, our strategy involves in-depth study of each specific step of the AA pathway. This led to the characterization of norbelladine synthase from *N. pseudonarcissus* (Singh et al. 2018), noroxomaritidine/norcrugsodine reductase in *N. pseudonarcissus* and *L. aestivum* (Majhi et al. 2023), as well as of cytochrome P450s (CYPs) cinnamate 4-hydroxylase, *p*-coumaroyl 3'-hydroxylase, and ascorbate peroxidase/4-coumarate 3-hydroxylase from *L. aestivum* (Karimzadegan et al. 2024). In addition to plants, we also assembled the transcriptome of two fungal species, the plant pathogen *Armillaria sinapina* and the medicinal fungus *Inonotus obliquus*, and identified terpenoid biosynthesis genes,

including squalene epoxidase and lupeol synthase (Fradj et al. 2019, 2020).

The research group also made strides in plant metabolite synthesis in heterologous systems. In addition to using yeast and bacteria (Majhi et al. 2023; Karimzadegan et al. 2024), the team bioengineered microalgae as platforms for specialized metabolite production and enzyme characterization. Although the process may be more tedious, it could lead to a more sustainable platform (Slattery et al. 2018; Diamond et al. 2023). The group succeeded in achieving stable expression of different *Cannabis sativa* and bacterial enzymes in the diatom *Phaeodactylum tricornutum* for the production of olivetolic acid, cannabigerolic acid and other cannabinoids (Awwad et al. 2023; Fantino et al. 2024). They have also optimized techniques to bioengineer other microalgae, including *Chlamydomonas reinhardtii* and *Chlorella vulgaris* (Carscallen et al. 2023; Desgagné-Penix et al. 2023; Beauchemin et al. 2024).

3. Investigating terpenes


Terpenes, derived from dimethylallyl pyrophosphate and isopentenyl pyrophosphate produced via the mevalonate and methylerythritol phosphate pathways, play critical roles in plant development, stress responses, and pollinator attraction (Singh and Sharma 2015; Pichersky and Raguso 2018; Boncan et al. 2020). Many of these compounds, such as paclitaxel, menthol, and artemisinin, are utilized in the food, cosmetic, and pharmaceutical industries (Wagner and Elmadfa 2003; da Silva-Santosa et al. 2005; Yang et al. 2020). Given the identification of thousands of terpenes and terpenoids, numerous research teams focus on characterizing the enzymes responsible for their synthesis (Kajikawa et al. 2005; Chen et al. 2011; Cornish 2014; Boutanaev et al. 2015; Singh and Sharma 2015; Lau et al. 2016; Huang and Osbourn 2019; Kopaczky et al. 2020; Zhou and Pichersky 2020; Amerik et al. 2021; Long et al. 2021; Jaramillo-Madrid et al. 2022).

This section details the contributions of professors Dae-Kyun Ro, Soheil Mahmoud, Jörg Bohlmann, Tariq A. Akhtar, and Reinhard Jetter in characterizing terpene synthases in gymnosperms and angiosperms. Additionally, it covers the research by professors Ro, Bohlmann, Akhtar, and Valerie C.A. Ward on the production of terpenoids in heterologous systems, including yeast, microalgae, and plants (Fig. 3).

3.1. Dae-Kyun Ro

At the University of Calgary, Pr. Dae-Kyun Ro studies the biosynthesis of terpenes, proanthocyanidins and natural rubber. His group used transcriptomics to identify *Pisum sativum* anthocyanidin, leucoanthocyanidin, and dihydroflavonol reductases, and characterized these enzymes in vitro, showing that they synthesize *cis*- and *trans*-flavan-3-ols (Ferraro et al. 2014). His research on sesquiterpene biosynthesis led to the identification and characterization of several enzymes in different pathways. In *Lippia dulcis* (Aztec sweet herb), they characterized (+)-*epi*- α -bisabolol synthase, which catalyzes the first step in the biosynthesis of hernandulcin, and bornyl diphosphate synthase, involved in the formation of camphor (Attia et al. 2012; Hurd et al. 2017). In *Valeriana officinalis* (valerian), the group characterized three sesquiterpene synthases

Fig. 3. Variety of terpenoids, and the plants that produce them, investigated in Canada and their main precursors, dimethylallyl phosphate and isopentenyl phosphate. Numbers in parenthesis indicate the principal investigators that research these topics. (2.1) Dae-Kyun Ro; (2.2) Soheil Mahmoud; (2.3) Jörg Bohlmann; (2.4) Tariq Akhtar; (2.5) Valerie CA Ward; (2.6) Reinhard Jetter. *Parthenium argentatum*'s photo was modified from Rousset et al. (2021). The following images were obtained from Wikimedia commons: *Adonis aestivalis*: By Osnat Ravid Amir (CC BY-SA 4.0) <https://commons.wikimedia.org/w/index.php?curid=151799697>; *Arabidopsis thaliana*: By Charles Andr  s (CC BY-SA 3.0) <https://commons.wikimedia.org/w/index.php?curid=22510218>; *Centella asiatica*: By Md. Siddiq Hasan (CC BY-SA 4.0) <https://commons.wikimedia.org/w/index.php?curid=114909062>; *Lavandula x intermedia*: By Photo by David J. Stang (First published at ZipcodeZoo.com, CC BY-SA 4.0) <https://commons.wikimedia.org/w/index.php?curid=61171226>; *Silene latifolia*: By Loz (L. B. Tettenborn, CC BY-SA 3.0) <https://commons.wikimedia.org/w/index.php?curid=4506347>; *Valeriana officinales*: By Ivar Leidus (CC BY-SA 4.0) <https://commons.wikimedia.org/w/index.php?curid=106650217>.



which synthesize germacrene C and D, valerana-4,7(11)-diene and (-)-drimenol (Pyle et al. 2012; Kwon et al. 2014). They also identified and characterized two germacrene A synthases from *Barnadesia spinosa* (espino santo or "holy thorn"), and one from *Lactuca sativa* (lettuce) (Nguyen et al. 2016). While studying two sesquiterpene oxidases from *Artemisia annuum*, they showed that germacrene A and amorphadiene oxidases are involved in the biosynthesis of a widespread and species-specific sesquiterpenes (Nguyen et al. 2019). They also identified and characterized a CYP that converts kolavenol to crotonolide G, a savorin-related compound (Kwon et al. 2022b). Finally, they characterized the promoters of *L. sativa*'s

Germacrene A Synthase 1 and 2, showing that these genes are expressed in parenchyma cells adjacent to laticifers (Kwon et al. 2022a).

The group's research on natural rubber biosynthesis includes works with *Parthenium argentatum* (guayule) and *L. sativa*, from which they identified and characterized cis-prenyltransferases (CPT) and CPT binding proteins (Lakusta et al. 2019; Barnes et al. 2021). In *L. sativa*, they show that CPT binding protein 2 (LsCBP2, previously named CPTL2) tethers CPT3 to the endoplasmic reticulum and synthesize cis-polyisoprenes in vitro (Qu et al. 2015b). Additionally, they identified the promoter region of these genes responsible for

their laticifer-specific expression (Barnes et al. 2021). Lastly, by expressing CPT3 from *P. argentatum* and *Solidago canadensis* (goldenrod) in natural rubber-deficient lettuce, they showed that these proteins can synthesize better quality natural rubber in the heterologous cell context, suggesting the quality of this compound is not dependent on CPT alone (Kwon et al. 2023).

In addition to the terpenoid pathway elucidation *in planta*, Pr. Ro has made important advances in the metabolic engineering of yeast as a platform to produce SMs during his graduate studies, and as a group leader or a collaborator. For instance, he succeeded in enhancing the carbon flux in mevalonate pathway through the expression of acetyl-CoA synthetase variant from *Salmonella enterica* with the aim of producing amorphaadiene, precursor of the high-value sesquiterpene artemesin (Shiba et al. 2007). Pr. Ro's group designed modular multiplex genome-edit-CRISPR platform to enhance feruloyl-CoA pathway in the aim of producing curcumin and other alkaloids in yeast (Utomo et al. 2024). Finally, his team's achievements *in planta* include works in *Nicotiana benthamiana* (a tobacco relative), and pea plants. In *N. benthamiana*, his group was able to increase the yield of sesqui- and triterpenes (α -bisabolol, amorphaadiene, valerenadiene and β -amyrin) in tobacco (Lee et al. 2019). This was achieved by discovering enzyme crosstalk and using push-pull strategies between different pathways. They recently targeted the pea plant gene β -amyrin synthase with CRISPR/Cas9 system to obtain saponin-free pea products, which are less bitter (Hodgins et al. 2024).

3.2. Soheil Mahmoud

An associate professor at UBC, Pr. Soheil Mahmoud has worked in the elucidation of monoterpene metabolism. From the transcriptome of *Coriandrum sativum* (coriander), his group identified and then characterized two monoterpene synthases, S-linalool synthase, and γ -terpinene synthase (Galata et al. 2014). Nevertheless, his research focuses on the biosynthesis of *Lavandula* spp. (lavender) essential oils. His efforts led to the publication of a draft genome for *L. angustifolia*, the first sequenced genome in the genus (Malli et al. 2019). He has used gene expression techniques, including expressed sequence tags, microarray, and RNA sequencing to identify and characterize terpene synthases in *Lavandula* spp. These include *L. angustifolia* β -phellandrene synthase (Demissie et al. 2011), borneol dehydrogenase, an enzyme in camphor biosynthesis pathway (Sarker et al. 2012), 1,8-cineole synthase (LiCINS) (Demissie et al. 2012), 9-epi-(E)-caryophyllene synthase (Sarker et al. 2013), lavandulyl diphosphate synthase (Demissie et al. 2013), two BAHD acyltransferases (Sarker and Mahmoud 2015), 3-carene synthase (Adal et al. 2017), S-linalool synthase (LiLINS) (Adal et al. 2019), geranyl diphosphate synthase, geranylgeranyl diphosphate synthase and farnesyl diphosphate synthase (Adal and Mahmoud 2020), and (+)-bornyl diphosphate synthase (Adal et al. 2023) from *L. x intermedia*. The team also identified and characterized several transcription factors from this hybrid that associate with the promoters of LiLINS and/or LiCINS (Sarker et al. 2019).

3.3. Jörg Bohlmann

Pr. Jörg Bohlmann started to investigate TPS while he was at the Max Planck Institute for Chemical Ecology in Germany, where he analyzed the expression profile of a *Nicotiana attenuata* gene encoding for a 5-*epi*-aristolochene synthase (an enzyme involved in capsidiol biosynthesis) (Bohlmann et al. 2002). He also identified and characterized a TPS from *Arabidopsis thaliana* (Bohlmann et al. 2000) that synthesizes myrcene, as well as (E)- β -ocimene. At UBC, his group performed *in silico* analysis of predicted TPSs from *A. thaliana* (Aubourg et al. 2002) and later characterized a (E)- β -ocimene synthase from this species (Faldt et al. 2003a).

Subsequent works of his group focused on terpene synthesis in gymnosperms. In addition to understanding the role of terpenes in conifer defense against pathogens and pests and how this response is affected by methyl jasmonate (Faldt et al. 2003b; Huber et al. 2005; Miller et al. 2005; Ro and Bohlmann 2006; Zulak et al. 2009, 2010; Hall et al. 2011b), they identified many genes and functionally characterized TPSs in *Pseudotsuga menziesii* (Douglas fir) (Huber et al. 2005), *Pinus taeda* (loblolly pine) (Ro and Bohlmann 2006), *Picea abies* (Norway spruce) (Faldt et al. 2003b; Martin et al. 2004; Keeling et al. 2011a), *Picea glauca* (white spruce) (Keeling et al. 2010), *Abies balsamea* (balsam fir) (Zerbe et al. 2012), and *Picea sitchensis* (Sitka spruce) (Hamberger et al. 2011; Roach et al. 2014). In a single study, his group identified and functionally characterized 15 monoterpene synthases from spruce, including 4 sesquiterpene synthases and 2 diterpene synthases of Sitka spruce, white spruce, and hybrid white spruce (Keeling et al. 2011b). They also characterized and performed evolutionary studies of cytochrome P450s involved in diterpene resin acid biosynthesis in different conifers, including enzymes of the CYP720B, CYP750B, and CYP76AA subfamilies (Ro and Bohlmann 2006; Hamberger et al. 2011; Gesell et al. 2015; Geisler et al. 2016).

Along with his work on conifers, Pr. Bohlmann's team also worked with angiosperms, characterizing (–)-germacrene D synthase from *Populus trichocarpa x deltoides* (hybrid poplar) (Arimura et al. 2004); characterizing TPSs from *Vitis vinifera* (grape) (Martin et al. 2010) and evaluating the relationship between wine aroma and terpene biosynthesis (Martin and Bohlmann 2004; Martin et al. 2009, 2012); identifying and characterizing CYPs that synthesize sesquiterpenols of *Santalum album* (sandalwood) oil (Celedon et al. 2016; Diaz-Chavez et al. 2013); and identifying and characterizing TPSs from *C. sativa* (Booth et al. 2017, 2020). Pr. Bohlmann also contributed to plant synthetic biology through the identification of cannabinoid glycosides and participating in reconstruction of pathways in yeast and *N. benthamiana* (Gülck et al. 2020).

Finally, his group elucidated the biosynthesis pathway of MontbretinA, identifying and characterizing several UDP-dependent glycosyltransferases, BAHD-acyltransferases, 4-coumaroyl-CoA ligases, and acyl activating enzyme (Irmisch et al. 2018, 2019, 2020; Sunstrum et al. 2021). They finally reconstructed the whole pathway in *N. benthamiana* (Irmisch et al. 2020).

3.4. Tariq A. Akhtar

Pr. Tariq Akhtar is an associate professor at University of Guelph, where he investigates plant metabolism. His work with specialized metabolism started with Pr. Eran Pichersky (University of Michigan) on the identification of the tomato CPT gene family (Akhtar et al. 2013). The investigation of polyprenols led to the characterization of CPT from tomato and *A. thaliana*, both located in the chloroplast stroma, which elongate geranylgeranyl diphosphate (Akhtar et al. 2017; Van Gelder et al. 2018). Furthermore, they showed that the biosynthesis of the polyprenol dolichol requires a CPT and an accessory protein which localizes in the ER and Golgi system in tomato (Brasher et al. 2015) and reconstructed the pathway in *N. benthamiana* (Van Gelder et al. 2021).

His group expanded on his initial work with Pr. Pichersky on veratrole biosynthesis in *Silene latifolia* (white campion) (Akhtar and Pichersky 2013; Gupta et al. 2012), showing that salicylic acid is catabolized into veratrole white champion, with catechol and guaiacol as intermediates (Van Gelder et al. 2020). Finally, his team also proposed the biosynthetic pathway of bibenzyls and cannflavins A and B in *C. sativa* (Boddington et al. 2022; Rea et al. 2019), along with the identification of the enzymes of both pathways.

3.5. Valerie CA Ward

Pr. Valerie CA Ward is an assistant professor at the University of Waterloo at the Chemical Engineering Department. She works on metabolic engineering of microorganisms for isoprenoid production (Edgar et al. 2021) and finding efficient process for high yield extraction from wet biomass. Dr Ward's team bioengineered *E. coli* to produce astaxanthin, expressing genes from the β -carotene pathway of the gram-negative bacteria *Pantoea agglomerans* and genes from the plant *Adonis aestivalis* (summer pheasant's eye) encoding for β -carotene hydroxylase and ketolase (Roth and Ward 2023). Dr Ward, in collaboration with Pr. Gregory Stephanopoulos from the Massachusetts Institute of Technology, has also worked with other organisms to produce fat-soluble isoprenoids, including the lipid-producing yeast *Yarrowia lipolytica* (Stephanopoulos et al. 2019; Edgar et al. 2021) and the microalga *Chlorella vulgaris* (Roushan et al. 2023).

3.6. Reinhard Jetter

Though his main focus is on plant cuticle composition and plant-insect interaction (Greer et al. 2007; Hegebarth et al. 2017; Sun et al. 2020, 2021, 2023), it is important to highlight Pr. Jetter's (UBC) contributions in the triterpenoid biosynthesis research. His group identified and characterized two CYPs from *Centella asiatica* (Indian/Asiatic pennywort), involved in the biosynthesis of the saponin asiaticoside (Kim et al. 2018), as well as oxidosqualene cyclases from *Ricinus communis* (castor bean) (Guhling et al. 2006), *Kalanchoe daigremontiana* (mother of thousands) (Wang et al. 2010), *Solanum lycopersicum* (tomato) (Wang et al. 2011), and *Quercus suber* (cork oak) (Busta et al. 2020).


4. Investigating polyketides and phenolic compounds

Phenolic compounds and polyketides have similar origins, with acyl molecules, from acyl-coenzyme A substrates, as precursors. Phenolic compounds, such as phenolic acids and lignans, are synthesized from the acetate-malonate or the shikimate pathway, with malonyl-coenzyme A, erythrose 4-phosphate, and phosphoenolpyruvate as primary metabolite precursors (Babenko et al. 2019; de la Rosa et al. 2019; Zagorskina et al. 2023). Meanwhile, polyketides, such as cannabinoids and flavonoids, are formed through the extension of an initial acyl molecules (such as acetyl, malonyl, and hexanoyl) by condensation of malonyl units by type III polyketide synthase, followed by the cyclization of the resulting chain (Abe and Morita 2010; Lussier et al. 2012). In plants, these compounds contribute to structure and defense, scavenging of reactive oxygen species, and attraction of pollinators (Di Ferdinando et al. 2012; Dias et al. 2020; Juca et al. 2020). With antioxidant and anti-inflammatory properties, these metabolites are of interest in medicine for targeting cancer, diabetes and other diseases (Al Jitan et al. 2018; Zhang et al. 2018b).

From 1983 to 2021, more than 20 chalcone synthase (CHS) genes had been characterized (Niu et al. 2021) and there have been many efforts to understand their evolution and that of CHS-like enzymes across the plant kingdom (Han et al. 2014; Naake et al. 2021; Guo et al. 2022). We described below the contributions of Pr. Dae-Yeon Suh's team to our understanding of polyketide synthase evolution and function in both bryophytes and angiosperms (Fig. 4). On a different front, the research on phytocannabinoid biosynthesis has included transcriptomics of *C. sativa*, evaluating transcription factors and stresses that regulate the production of cannabinoids and other specialized metabolites (Braich et al. 2019; Jalali et al. 2019; Zager et al. 2019; Apicella et al. 2022; Kim et al. 2022); and population studies, which have identified several markers related to morphology and productivity, cannabinoid content (Adamek et al. 2023; de Ronne et al. 2024; Dehnavi et al. 2025). This fundamental research is the basis for improving the production of the desired metabolites through breeding efforts, gene silencing, or synthetic biology (Geissler et al. 2018; Zirpel et al. 2018; Luo et al. 2019; Gülck et al. 2020; Thomas et al. 2020; Matchett-Oates et al. 2021; Ma et al. 2022; Awwad et al. 2023; Desgagné-Penix et al. 2023; Fantino et al. 2024). These studies were made possible by the pioneering contributions of Pr. Page's team on cannabis genomics, which are described below (Fig. 4).

Canadian researchers have identified genes and enzymes involved in the metabolism of phenolic compounds in crops like beans and poplar, focusing on transcription factors controlling their biosynthesis and catabolism. Global research has similarly elucidated mechanisms driving phenolic compound accumulation in these plants, including transcriptomic and genomic studies identifying phenolic compound-associated loci (Kosonen et al. 2012; Cho et al. 2016; James et al. 2017; Perez de Souza et al. 2019; He et al. 2021; Bryant et al. 2023; Campa et al. 2023; Celebioglu et al. 2023; Chen et al. 2023; Li et al. 2023; Ma et al. 2023, 2024; Tian et al.

Fig. 4. Polyketides and other phenolic compounds investigated in Canada, along with their precursors and plants that produce them. Numbers in parenthesis indicate principal investigators working in the subject. (3.1) Dae-Yeon Suh; (3.2) Jonathan E. Page; (3.3 and 3.4) Gale G. Bozzo and Sangeeta Dhaubhadel, respectively, who work on isoflavonoids and anthocyanins in beans; (3.5) C. Peter Constabel; (3.7) Nikola Kovic. *Cannabis sativa*'s drawing was done with Krita 5.2.9. Illustrations for *Glycine max* and *Phaseolus vulgaris* were obtained from Vecteezy; *Physcomitrium patens*'s drawing was obtained from Lang et al. (2018). *Populus tremuloides*' photo was Wikimedia Commons (By I, Hugo.arg (CC BY-SA 3.0) <https://commons.wikimedia.org/w/index.php?curid=2268194>).



2021; Tapia et al. 2024; Zhang et al. 2018a, 2023, 2024). For detailed insights, readers are referred to comprehensive reviews (Pourcel et al. 2007; Cheynier et al. 2013; Corso et al. 2020; Ma and Constabel 2021; Li et al. 2022a; Pratyusha and Sarada 2022; Sunil and Shetty 2022; Islam and Dhaubhadel 2023; Trush and Pal'ove-Balang 2023).

This section discusses the efforts of Professors Gale Bozzo, Sangeeta Dhaubhadel, and Mehran Dastmalchi on anthocyanin and flavonoid metabolism, and Professors C. Peter Constabel and Nikola Kovic on transcription factors in polyphenol metabolism (Fig. 4).

4.1. Dae-Yeon Suh

Having previously worked on chalcone synthases with Pr. Ushio Sankawa in Japan (Kojima et al. 2000; Okada et al. 2001; Suh et al. 2000a, 2000b, 2000c; Yamazaki et al. 2001), Pr. Dae-Yeon Suh established his research team at Univer-

sity of Regina, where they work on both bryophytes and angiosperms and their CHS enzymes. His group performed evolutionary studies on plant enzymes of the thiolase superfamily and CHS family (Jiang et al. 2008), collaborated with Pr. Sankawa in analysing the effects of mutating highly conserved arginine residues in CHS (Fukuma et al. 2007) and characterized a CHS from the spreading earthmoss (*Physcomitrium patens*) (Jiang et al. 2006). Later they used expressed sequence tags from the bryophyte and identified a gene encoding for Anther-Specific CHS-Like Enzyme, which was shown to play a role in sporopollenin biosynthesis, like its ortholog in arabidopsis, which is essential for reproduction (Colpitts et al. 2011; Daku et al. 2016). They also characterized 2-oxoalkylresorcinol synthase in *P. patens*, an enzyme proposed to be similar to ancestral type III polyketide synthases, showing that it plays a role in cuticle formation, and therefore is necessary for resistance to dehydration (Kim et al. 2013; Li et al. 2018; Aslam et al. 2022). His group has also

performed genome-wide analysis of spreading earthmoss' class III peroxidase gene family, with 49 members, and CHS gene superfamily, with 17 putative members (Aparato et al. 2024; Koduri et al. 2010). While the CHS gene superfamily's expansion was proposed to be through segmental duplication (Koduri et al. 2010), the group hypothesized that the expansion of class III peroxidase gene family was due to whole-genome duplication (Aparato et al. 2024). Finally, another CHS involved in sporopollenin biosynthesis that the group characterized was *Polyketide Synthase 1*, from the angiosperm St John's wort's (*Hypericum perforatum*), showing that the enzyme uses medium to very-long chain fatty acyl-coenzyme A substrates to produce triketide and tetraketide alkylpyrones and that, in nonanther tissues, it could be involved in the biosynthesis of components of cell wall (Jepson et al. 2014).

4.2. Jonathan E. Page

A professor at the Department of Botany of UBC, Jonathan E. Page, has carried out works on tobacco, *Humulus lupulus* (hop) and *Cannabis* spp., investigating the biosynthesis of alkaloids, flavonoids, and cannabinoids. In *N. benthamiana*, his group analyzed expressed sequence tags from methyl-jasmonate-treated roots, gene silencing, and gene overexpression to identify two basic Helix-Loop-Helix transcription factors acting as positive regulators of nicotine biosynthesis in response to jasmonate (Todd et al. 2010). This work led to two patents, one on the uses of *N*-methylputrescine (Page and Liu 2023), and the other on the transcription factors impacting the production of alkaloids in this plant (Page and Todd 2023). In hop, the group applied expressed sequence tags and transcriptome sequencing for the identification of a methyltransferase that synthesizes xanthohumol (Nagel et al. 2008), and of genes involved in branched-chain amino acid biosynthesis and catabolism (Clark et al. 2013). Pr. Page's team also filed two patents on specialized metabolism of hops plants (Nagel and Page 2010; Page et al. 2013).

Pr. Page's team has contributed significantly to the cannabinoid metabolism research, starting with the draft genome and transcriptome assembly of *C. sativa* "Purple Kush" in 2011 (van Bakel et al. 2011). Later, they generated a genetic map of the species by crossing this strain with hemp "Finola" strain, finding the genes cannabidiolic acid and tetrahydrocannabinolic acid synthases (*CBDAS* and *THCAS*, respectively) in a retrotransposon-rich region. Moreover, studying strains of marijuana and hemp using genotyping-by-sequencing, his group showed the greater heterozygosity in hemp, and the differences in many loci, not only in the *CBDAS/THCAS* genes (Sawler et al. 2015). Finally, they identified and characterized two acyl-activating enzymes able to synthesize hexanoyl-CoA (Stout et al. 2012) and olivetolic acid cyclase (Gagne et al. 2012) from expressed sequence tags obtained from glandular trichomes of cannabis flowers, and identified three transcription factors that regulate *THCAS* expression (Liu et al. 2021). Pr. Page has three patents related to cannabinoid biosynthesis enzyme characterization (Page et al. 2020; Page and Stout 2020; Page and Gagne 2024). These patents were widely used by companies (including Intrexon, Gingko Bioworks, Hyacinth, and Demetrix) or research insti-

tutes to produce cannabinoid compounds in vivo or in vitro (Awwad et al. 2023).

4.3. Gale G. Bozzo

Pr. Gale G. Bozzo is an associate professor at the University of Guelph interested in controlled atmosphere storage and specialized metabolism, mainly phenylpropanoids. His work with cranberry beans (*Phaseolus vulgaris* L.) showed that darkening of seed coat postharvest was associated with increased expression of proanthocyanidins biosynthesis genes, including *Anthocyanidin Reductase 1*, which converts cyanidin to epicatechin and catechin (Freixas Coutin et al. 2017). With common beans, the team showed that expression of isoflavone biosynthetic genes and concentration of isoflavones were up-regulated in a line resistant to common bacterial blight (caused by the bacterium *Xanthomonas axonopodis*), but not in the susceptible line (Cox et al. 2021). In addition, they discovered a β -glucosidase from *Arabidopsis* (BGLU15) which catabolized flavonol rhamnosides in nitrogen-deprived plants exposed to low temperatures (Roepke and Bozzo 2015), and later they showed that flavonol rhamnosides are also catabolized in radish stored in cold temperature and that this could be associated with the activity of an α -rhamnosidase (Unterlander et al. 2022).

4.4. Sangeeta Dhaubhadel

Pr. Sangeeta Dhaubhadel, a researcher at the Agriculture and Agri-Food Canada and adjunct professor at the University of Western Ontario, works on the metabolism of phenylpropanoids, isoflavonoids, and anthocyanins in legume species, including common bean and soybean (*P. vulgaris* and *Glycine max*, respectively), along with her research on soybean defense (Khatri et al. 2022; Scott et al. 2021, 2022). Her team showed that soybean embryos can synthesize isoflavonoids de novo (Dhaubhadel et al. 2003) and that the genes encoding for chalcone synthases 7 and 8 are more expressed in a cultivar that accumulates more isoflavonoids (Dhaubhadel et al. 2003). Later, they identified the transcription factor *GmMYB176*, which regulates the expression of *GmCS8* (Anguraj Vadivel et al. 2019; Yi et al. 2010), showing that it interacts with 14-3-3 proteins, affecting its localization in the cell (Li and Dhaubhadel 2012), and with *GmbZIP5*, involved in isoflavonoid and phytoalexin metabolism and accumulation (Anguraj Vadivel et al. 2021). In another study, they identified two genes encoding flavonoid-metabolizing enzymes, flavonoid 3'-hydroxylase and dihydroflavonol 4-reductase, which compete for substrates with isoflavonoid biosynthesis pathway and were highly down-regulated in isoflavonoid-rich cultivars (Dastmalchi et al. 2017). In addition, her group identified and characterized three soybean genes encoding for cinnamate 4-hydroxylases (Khatri et al. 2023).

Similar to what was shown by Pr. Bozzo's team (Freixas Coutin et al. 2017), her team revealed a correlation between the slow darkening of the seed coat of pinto beans, and the decrease in biosynthesis and accumulation of proanthocyanidins (Duwadi et al. 2018). Later, they found that the vacuolar transporter *PmMATE8* is down-regulated in the slow dark-

ening cultivar compared to the regular darkening cultivar, and that there is possibly a feedback regulation between the biosynthesis of proanthocyanidins and their accumulation in the vacuole (Islam et al. 2022).

4.5. C. Peter Constabel

Pr. C. Peter Constabel, from University of Victoria, has performed extensive research on poplar's defense mechanisms against herbivores and fungal pathogens. His group evaluated the impact of wound and herbivory on the expression of a hybrid poplar polyphenol oxidase (Constabel et al. 2000) and later showed, in *Populus tremuloides* (quaking aspen), that this enzyme contributes to plant defense by oxidizing catechol (Haruta et al. 2001). They also identified a UDP-glycosyltransferase, which co-expressed with proanthocyanidin biosynthesis genes, able to glycosylate the flavonols quercetin and kaempferol (Veljanovski and Constabel 2013) and a gene, encoding a dihydroflavonol reductase whose induced expression by herbivory led to increased accumulation of condensed tannins (Peters and Constabel 2002). Moreover, Pr. Constabel's team identified several MYB transcription factors that regulate proanthocyanidin biosynthesis genes, among others such as shikimate pathway genes, including the activators MYB134 (Mellway et al. 2009), which was shown to act mainly in the leaves (Gourlay et al. 2020), MYB115 (James et al. 2017; Liu et al. 2023), MYB117 (Ma et al. 2021), and the repressors MYB182 (Yoshida et al. 2015), MYB165 and MYB194 (Ma et al. 2018). His group also showed that the apple transcription factor MYB9 has a similar function to poplar's MYB134 (Gesell et al. 2014).

Besides his work on flavonoids, he also works on salicoids, which are phenolic glycosides of the *Salicaceae* family. The team characterized several enzymes involved in the biosynthesis of these metabolites, including: a salicyl benzoate acyltransferase and a benzyl benzoate acyltransferase (Chedgy et al. 2015); two UDP-glycosyltransferases involved in the synthesis of salicortin, tremulacin, tremuloidin, and salicin (Fellenberg et al. 2020); and a sulfotransferase that is able to synthesize these newly identified sulfated salicinoids (Lackus et al. 2020).

4.6. Mehran Dastmalchi

Pr. Mehran Dastmalchi previously worked with Pr. Sangeeta Dhaubhadel on bean flavonoids (Dastmalchi and Dhaubhadel 2015; Dastmalchi et al. 2019a), with Pr. Vincenzo de Luca on *Catharanthus* metabolites, and Pr. Peter J. Facchini on BIA metabolism (Morris et al. 2016; Dastmalchi et al. 2018, 2019a, 2019b). After opening his own laboratory at McGill University focusing on isoflavonoids biosynthesis in *Fabaceae* and *Apocyanaceae*, he published several reviews on his expertise on plant specialized metabolism (Dastmalchi 2020a, 2020b, 2021a, 2021b). Furthermore, his team showed that the accumulation of the alkaloid akuammicine (a weak opioid agonist and potential anti-diabetic drug) can be increased in *C. roseus* using methyljasmonate, and that naturally formed lesions in a mutant had more of this alkaloid (Li et al. 2022b). Pr. Dastmalchi's team also identified and characterized a transporter Multidrug And Toxic compound

Extrusion 1, which imports secologanin (a key intermediate in the synthesis of MIAs) into the vacuole, and revealed that in its absence this monoterpenoid is reduced to secologanol in *C. roseus* (Li et al. 2024).

4.7. Nikola Kovinich

Nikola Kovinich was appointed assistant professor at York University in 2019, where he works on soybean responses to biotic stress. Specifically, Pr. Kovinich is interested in understanding how biotic stresses affect soybean's metabolism (Yousefi-Taemeh et al. 2021) and the regulation of phytoalexin biosynthesis mediated by MYB and NAC transcription factors (Farrell et al. 2017; Kovinich and Durkin 2018; Jahan and Kovinich 2019; Jahan et al. 2020; Lin et al. 2023b). In addition, Pr. Kovinich has broadened his research on plant metabolites to include the semi-synthesis of anticancer natural products using soybean seed coats (Gary et al. 2018) and he has contributed with methods for genetic transformation of soybean (Lin et al. 2023a) and cannabis (Ahmed et al. 2021).

5. Biofoundries and incubators

In addition to the uplisted research teams, Canada is home to two biofoundries: Centre for Applied Synthetic Biology (Concordia Genome Foundry) and the BioFoundry at UBC. Both offer metabolic and enzyme engineering incubators in an academic environment to achieve more sustainable and profitable pharmaceutical production of molecules of interest. Concordia Genome Foundry allowed companies to use high-throughput genomic techniques and develop metabolic engineering host cells producing isoprenoids and cannabinoids (Keasling et al. 2007; Keasling et al. 2011).

The BioFoundry at UBC is led by Pr. Vikramaditya G. Yadav, an Associate Professor in the Department of Chemical & Biological Engineering, and the School of Biomedical Engineering UBC. The group collaborates with several companies and works on enzyme optimization and drug development. Among the fruits of these collaborations are innovations in the synthesis of cannabinoids (Ayakar et al. 2020; Yadav et al. 2022).

6. Conclusion

Canada is a fertile environment for advancing plant metabolomics and platforms to produce plant specialized compounds due to active researchers and their previous mentors. These include one of the pioneers of the Canadian research on plant specialized metabolism, Pr. G.H. Neil Towers that mentored Pr. Ragai K. Ibrahim, who then trained Pr. Vincenzo de Luca. Though this review focused on researchers directly working on pathway elucidation and metabolic engineering of plant specialized metabolites, many advances have also been supported by interdisciplinary collaborations. Researchers such as Pr. John T. Arnason (Binns et al. 2001), Pr. Jürgen Ehrling (Carrington et al. 2018; Adams et al. 2019; Alber et al. 2019), Pr. Abdelali Hannoufa (Yu et al. 2007, 2008; Bhinu et al. 2009; Ben-Abdallah et al. 2018; Carrington et al. 2018), and Pr. Charles Goulet (Bulut et al. 2022; Liscombe et al. 2022), to name a few, have significantly advanced our un-

derstanding of specialized metabolism and its applications across various fields, including plant stress responses, ecological interactions, and agricultural innovations.

Canadian research institutes are offering new insights into plant biochemistry and synthetic biology every year. A pan-Canadian research collaboration could allow major advances in this field and help young researchers to obtain funding, which would help Canada to retain talent of the new scientific generation.

Article information

History dates

Received: 10 July 2024

Accepted: 30 January 2025

Version of record online: 25 April 2025

Notes

This paper is part of a collection entitled “Current status of Plant Biotechnology in Canada.” <https://cdnsiencepub.com/topic/gen-plant-biotechnology-canada>.

Copyright

© 2025 The Author(s). This work is licensed under a [Creative Commons Attribution 4.0 International License](#) (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Data availability

Data generated or analyzed during this study are provided in full within the published article.

Author information

Author ORCIDs

Isabel Desgagné-Penix <https://orcid.org/0000-0002-4355-5503>

Author contributions

Conceptualization: KCGdS, FA, ID

Supervision: ID

Validation: NM

Writing – original draft: KCGdS, FA, NM

Writing – review & editing: KCGdS, FA, NM, ID

Competing interests

The authors declare there are no competing interests.

Funding information

This work was supported by grants from the Natural Sciences and Engineering Research Council of Canada award number RGPIN-2021-03218 and the Canada Research Chair on Plant Specialized Metabolism award number 950-232164. Many thanks are extended to the Canadian taxpayers and to the Canadian government for supporting the Canada Research Chairs Program.

References

- Abe, I., and Morita, H. 2010. Structure and function of the chalcone synthase superfamily of plant type III polyketide synthases. *Nat. Prod. Rep.* 27(6): 809–838. doi:10.1039/b909988n.
- Adal, A.M., and Mahmoud, S.S. 2020. Short-chain isoprenyl diphosphate synthases of lavender (*Lavandula*). *Plant Mol. Biol.* 102(4-5): 517–535. doi:10.1007/s11103-020-00962-8.
- Adal, A.M., Sarker, L.S., Lemke, A.D., and Mahmoud, S.S. 2017. Isolation and functional characterization of a methyl jasmonate-responsive 3-carene synthase from *Lavandula x intermedia*. *Plant Mol. Biol.* 93(6): 641–657. doi:10.1007/s11103-017-0588-6.
- Adal, A.M., Najafianashrafi, E., Sarker, L.S., and Mahmoud, S.S. 2023. Cloning, functional characterization and evaluating potential in metabolic engineering for lavender (+)-bornyl diphosphate synthase. *Plant Mol. Biol.* 111(1-2): 117–130. doi:10.1007/s11103-022-01315-3.
- Adal, A.M., Sarker, L.S., Malli, R.P.N., Liang, P., and Mahmoud, S.S. 2019. RNA-seq in the discovery of a sparsely expressed scent-determining monoterpene synthase in lavender (*Lavandula*). *Planta*, 249(1): 271–290. doi:10.1007/s00425-018-2935-5.
- Adamek, K., Grainger, C., Jones, A.M.P., and Torkamaneh, D. 2023. Genotyping-by-sequencing (GBS) reveals greater somatic mutations than simple sequence repeats (SSRs) in micropropagated cannabis plants. *In Vitro Cell. Dev. Biol. Plant*, 59: 756–766. doi:10.1007/s11627-023-10377-x.
- Adams, Z.P., Ehlting, J., and Edwards, R. 2019. The regulatory role of shikimate in plant phenylalanine metabolism. *J. Theor. Biol.* 462: 158–170. doi:10.1016/j.jtbi.2018.11.005.
- Aghaali, Z., and Naghavi, M.R. 2024. Developing benzylisoquinoline alkaloid-enriched opium poppy via CRISPR-directed genome editing: a review. *BMC Plant Biol.* 24(1). doi:10.1186/s12870-024-05412-x.
- Aghaali, Z., Naghavi, M.R., and Zargar, M. 2024. Promising approaches for simultaneous enhancement of medicinally significant benzylisoquinoline alkaloids in opium poppy. *Front. Plant Sci.* 15. doi:10.3389/fpls.2024.1377318.
- Ahmed, S., Gao, X., Jahan, M.A., Adams, M., Wu, N., and Kovich, N. 2021. Nanoparticle-based genetic transformation of *Cannabis sativa*. *J. Biotechnol.* 326: 48–51. doi:10.1016/j.jbiotec.2020.12.014.
- Akhtar, T.A., and Pichersky, E. 2013. Veratrole biosynthesis in white campion. *Plant Physiol.* 162(1): 62. doi:10.1104/pp.113.214346.
- Akhtar, T.A., Matsuba, Y., Schauvinhold, I., Yu, G., Lees, H.A., Klein, S.E., and Pichersky, E. 2013. The tomato cis-prenyltransferase gene family. *Plant J.* 73(4): 652. doi:10.1111/tpj.12063.
- Akhtar, T.A., Surowiecki, P., Siekierska, H., Kania, M., Van Gelder, K., Rea, K.A., et al. 2017. Polyphenols are synthesized by a plastidial cis-prenyltransferase and influence photosynthetic performance. *Plant Cell*, 29(7): 1709–1725. doi:10.1105/tpc.16.00796.
- Al Jitan, S., Alkhoori, S.A., and Yousef, L.F. 2018. Phenolic acids from plants: extraction and application to human health. In *Studies in natural products chemistry*. Edited by A. Rahman. Elsevier, Amsterdam, Netherlands. pp. 389–417.
- Alber, A.V., Renault, H., Babilio-Lopes, A., Bassard, J.E., Liu, Z., Ullmann, P., et al. 2019. Evolution of coumaroyl conjugate 3-hydroxylases in land plants: lignin biosynthesis and defense. *Plant J.* 99(5): 924–936. doi:10.1111/tpj.14373.
- Amerik, A.Y., Martirosyan, Y.T., Martirosyan, L.Y., Goldberg, V.M., Uteulin, K.R., and Varfolomeev, S.D. 2021. Molecular genetic analysis of natural rubber biosynthesis. *Russ. J. Plant Physiol.* 68: 31–45. doi:10.1134/S1021443721010039.
- Anguraj Vadivel, A.K., Renaud, J., Kagale, S., and Dhaubhadel, S. 2019. GmMYB176 regulates multiple steps in isoflavonoid biosynthesis in soybean. *Front. Plant Sci.* 10: 562. doi:10.3389/fpls.2019.00562.
- Anguraj Vadivel, A.K., McDowell, T., Renaud, J.B., and Dhaubhadel, S. 2021. A combinatorial action of GmMYB176 and GmZIP5 controls isoflavonoid biosynthesis in soybean (*Glycine max*). *Commun. Biol.* 4(1): 356. doi:10.1038/s42003-021-01889-6.
- Aparato, V.P.M., Rabbi, F., Madarash, T., Brisbourne, W.A., Barker, E.I., and Suh, D.Y. 2024. Genome-wide analysis of the class III peroxidase gene family in *Phiscomitrium patens* and search for clues to ancient class III peroxidase functions. *Int. J. Plant Biol.* 15(4): 1141–1161. doi:10.3390/ijpb15040079.
- Apicella, P.V., Sands, L.B., Ma, Y., and Berkowitz, G.A. 2022. Delineating genetic regulation of cannabinoid biosynthesis during female flower

- development in *Cannabis sativa*. *Plant Direct*, 6(6): e412. doi:10.1002/pld3.412.
- Arimura, G., Huber, D.P., and Bohlmann, J. 2004. Forest tent caterpillars (*Malacosoma disstria*) induce local and systemic diurnal emissions of terpenoid volatiles in hybrid poplar (*Populus trichocarpa* x *deltoides*): cDNA cloning, functional characterization, and patterns of gene expression of (-)-germacrene D synthase, PtdTPS1. *Plant J.* 37(4): 603–616. doi:10.1111/j.1365-3113x.2003.01987.x.
- Asada, K., Salim, V., Masada-Atsumi, S., Edmunds, E., Nagatoshi, M., Terasaka, K., et al. 2013. A 7-deoxyloganetic acid glucosyltransferase contributes a key step in secologanin biosynthesis in *Madagascar periwinkle*. *Plant Cell*, 25(10): 4134. doi:10.1105/tpc.113.115154.
- Aslam, M., Aparato, V.P.M., and Suh, D.Y. 2022. (2'-Oxo)alkylresorcinols restore dehydration tolerance in a knockout line of PpORS, a bryophyte-specific type III polyketide synthase in *Physcomitrium* (*Physcomitrella*) *patens*. *Planta*, 255(6): 129. doi:10.1007/s00425-022-03909-z.
- Attia, M., Kim, S.U., and Ro, D.K. 2012. Molecular cloning and characterization of (+)-*epi*- α -bisabolol synthase, catalyzing the first step in the biosynthesis of the natural sweetener, *hernandulcin*, in *Lippia dulcis*. *Arch. Biochem. Biophys.* 527(1): 37–44. doi:10.1016/j.abb.2012.07.010.
- Aubourg, S., Lecharny, A., and Bohlmann, J. 2002. Genomic analysis of the terpenoid synthase (*At*TPS) gene family of *Arabidopsis thaliana*. *Mol. Genet. Genomics*, 267(6): 730–745. doi:10.1007/s00438-002-0709-y.
- Awwad, F., Fantino, E.L., Héneault, M., Diaz-Garza, A.M., Merindol, N., Custeau, A., et al. 2023. Bioengineering of the marine diatom *Phaeodactylum tricornutum* with cannabis genes enables the production of the cannabinoid precursor, olivetolic acid. *Int. J. Mol. Sci.* 24(23). doi:10.3390/ijms242316624.
- Ayakar, S.R., Pawar, S.V., Hallam, S.J., Hossain, S., Yadav, V.G., Roy, P.R., and Srivastava, S.K. 2020. Metabolic engineering of *E. coli* for the biosynthesis of cannabinoid products. *Edited by USPTO*. Inmed Pharmaceuticals Inc, USA.
- Babenko, L.M., Smirnov, O.E., Romanenko, K.O., Trunova, O.K., and Kosakivska, I.V. 2019. Phenolic compounds in plants: biogenesis and functions. *Ukr. Biochem. J.* 91(3): 5–18. doi:10.15407/ubj91.03.005.
- Barnes, E.K., Kwon, M., Hodgins, C.L., Qu, Y., Kim, S.W., Yeung, E.C., and Ro, D.K. 2021. The promoter sequences of lettuce di-*prenyltransferase* and its binding protein specify gene expression in laticifers. *Planta*, 253(2): 51. doi:10.1007/s00425-021-03566-8.
- Beauchemin, R., Merindol, N., Fantino, E., Lavoie, P., Nouemssi, S.B., Meddeb-Mouelhi, F., and Desgagné-Penix, I. 2024. Successful reversal of transgene silencing in *Chlamydomonas reinhardtii*. *Biotechnol. J.* 19(1): doi:10.1002/biot.202300232.
- Beaudoin, G.A., and Facchini, P.J. 2013. Isolation and characterization of a cDNA encoding (S)-*cis*-N-methylstylopine 14-hydroxylase from opium poppy, a key enzyme in sanguinarine biosynthesis. *Biochem. Biophys. Res. Commun.* 431(3): 603. doi:10.1016/j.bbrc.2012.12.129.
- Ben-Abdallah, S., Zorrig, W., Amyot, L., Renaud, J., Hannoufa, A., Lachâal, M., and Karray-Bourauoi, N. 2018. Potential production of polyphenols, carotenoids and glycoalkaloids in *Solanum villosum* Mill. Under salt stress. *Biologia (Bratisl)*, 74: 309–324. doi:10.2478/s11756-018-00166-y.
- Bhinu, V.S., Schafer, U.A., Li, R., Huang, J., and Hannoufa, A. 2009. Targeted modulation of sinapine biosynthesis pathway for seed quality improvement in *Brassica napus*. *Transgenic Res.* 18(1): 31–44. doi:10.1007/s11248-008-9194-3.
- Binns, S.E., Inparajah, I., Baum, B.R., and Arnason, J.T. 2001. Methyl jasmonate increases reported alkaloids and ketoalkenyls in *Echinacea pallida* (Asteraceae). *Phytochemistry*, 57(3): 417–420. doi:10.1016/s0031-9422(00)00444-1.
- Bird, D.A., Franceschi, V.R., and Facchini, P.J. 2003. A tale of three cell types: alkaloid biosynthesis is localized to sieve elements in opium poppy. *Plant Cell*, 15(11): 2626–2635. doi:10.1105/tpc.015396.
- Boddington, K.F., Soubeyrand, E., Van Gelder, K., Casaretto, J.A., Perrin, C., Forrester, T.J.B., et al. 2022. Bibenzyl synthesis in *Cannabis sativa* L. *Plant J.* 109(3): 693–707. doi:10.1111/tpj.15588.
- Bohlmann, J., Martin, D., Oldham, N.J., and Gershenzon, J. 2000. Terpenoid secondary metabolism in *Arabidopsis thaliana*: cDNA cloning, characterization, and functional expression of a myrcene/(E)- β -ocimene synthase. *Arch. Biochem. Biophys.* 375(2): 261–269. doi:10.1006/abbi.1999.1669.
- Bohlmann, J., Stauber, E.J., Krock, B., Oldham, N.J., Gershenzon, J., and Baldwin, I.T. 2002. Gene expression of 5-*epi*-aristolochene synthase and formation of capsidiol in roots of *Nicotiana attenuata* and *N. sylvestris*. *Phytochemistry*, 60(2): 109–116. doi:10.1016/s0031-9422(02)00080-8.
- Boncan, D.A.T., Tsang, S.S.K., Li, C., Lee, I.H.T., Lam, H.M., Chan, T.F., and Hui, J.H.L. 2020. Terpenes and terpenoids in plants: interactions with environment and insects. *Int. J. Mol. Sci.* 21(19). doi:10.3390/ijms21197382.
- Booth, J.K., Page, J.E., and Bohlmann, J. 2017. Terpene synthases from *Cannabis sativa*. *PLoS One*, 12(3): e0173911. doi:10.1371/journal.pone.0173911.
- Booth, J.K., Yuen, M.M.S., Jancsik, S., Madilao, L.L., Page, J.E., and Bohlmann, J. 2020. Terpene synthases and terpene variation in *Cannabis sativa*. *Plant Physiol.* 184(1): 130–147. doi:10.1104/pp.20.00593.
- Boutanaev, A.M., Moses, T., Zi, J., Nelson, D.R., Mugford, S.T., Peters, R.J., and Osbourn, A. 2015. Investigation of terpene diversification across multiple sequenced plant genomes. *Proc. Natl. Acad. Sci.* 112(1): E81–E88. doi:10.1073/pnas.1419547112.
- Braich, S., Baillie, R.C., Jewell, L.S., Spangenberg, G.C., and Cogan, N.O.I. 2019. Generation of a comprehensive transcriptome atlas and transcriptome dynamics in medicinal cannabis. *Sci. Rep.* 9(1): 16583. doi:10.1038/s41598-019-53023-6.
- Brasher, M.I., Surmacz, L., Leong, B., Pitcher, J., Swiezewska, E., Pichersky, E., and Akhtar, T.A. 2015. A two-component enzyme complex is required for dolichol biosynthesis in tomato. *Plant J.* 82(6): 903–914. doi:10.1111/tpj.12859.
- Bryant, N., Zhang, J., Feng, K., Shu, M., Ployet, R., Chen, J.G., et al. 2023. Novel candidate genes for lignin structure identified through genome-wide association study of naturally varying *Populus trichocarpa*. *Front. Plant Sci.* 14: 1153113. doi:10.3389/fpls.2023.1153113.
- Bui, V.H., Rodriguez-Lopez, C.E., and Dang, T.T. 2023. Integration of discovery and engineering in plant alkaloid research: recent developments in elucidation, reconstruction, and repurposing biosynthetic pathways. *Curr. Opin. Plant Biol.* 74: 102379. doi:10.1016/j.pbi.2023.102379.
- Bulot, B., Isabelle, S., Montoya, R., Nadeau, L.F., Tremblay, J., and Goulet, C. 2022. Structural variations in the phytoene synthase 1 gene affect carotenoid accumulation in tomato fruits and result in bicolor and yellow phenotypes (PREPRINT). *Biorxiv*. doi:10.1101/2022.10.29.514357. bioRxiv.
- Busta, L., Serra, O., Kim, O.T., Molinas, M., Pere-Fossoul, I., Figueras, M., and Jetter, R. 2020. Oxidosqualene cyclases involved in the biosynthesis of triterpenoids in *Quercus suber* cork. *Sci. Rep.* 10(1): doi:10.1038/s41598-020-64913-5.
- Campa, A., Rodriguez Madrera, R., Jurado, M., Garcia-Fernandez, C., Suarez Valles, B., and Ferreira, J.J. 2023. Genome-wide association study for the extractable phenolic profile and coat color of common bean seeds (*Phaseolus vulgaris* L.). *BMC Plant Biol.* 23(1): 158. doi:10.1186/s12870-023-04177-z.
- Carqueijeiro, I., Brown, S., Chung, K., Dang, T.T., Walia, M., Besseau, S., et al. 2018. Two tabersonine 6,7-epoxidases initiate lochnericine-derived alkaloid biosynthesis in *Catharanthus roseus*. *Plant Physiol.* 177(4): 1473–1486. doi:10.1104/pp.18.00549.
- Carr, S.C., Torres, M.A., Morris, J.S., Facchini, P.J., and Ng, K.K.S. 2021. Structural studies of codeinone reductase reveal novel insights into aldo-keto reductase function in benzyloisoquinoline alkaloid biosynthesis. *J. Biol. Chem.* 297(4): 101211. doi:10.1016/j.jbc.2021.101211.
- Carrington, Y., Guo, J., Le, C.H., Fillo, A., Kwon, J., Tran, L.T., and Ehltling, J. 2018. Evolution of a secondary metabolic pathway from primary metabolism: shikimate and quinate biosynthesis in plants. *Plant J.* 95. doi:10.1111/tpj.13990.
- Carscallen, W.M.A., Desgagné-Penix, I., and Lemay, J.-f. 2023. Engineered microorganism for the production of cannabinoid biosynthetic pathway products. *Edited by W.I.P. Organization*. Algae-C Inc., WO.
- Cazares-Flores, P., Levac, D., and De Luca, V. 2016. *Rauvolfia serpentina* N-methyltransferases involved in ajmaline and *n*- β -methylajmaline biosynthesis belong to a gene family derived from γ -tocopherol C-methyltransferase. *Plant J.* 87(4): 335–342. doi:10.1111/tpj.13186.
- Celebioglu, B., Hart, J.P., Porch, T., Griffiths, P., and Myers, J.R. 2023. Genome-wide association study to identify possible candidate genes

- of snap bean leaf and pod color. *Genes*, 14(12). doi:10.3390/genes14122234.
- Celedon, J.M., Chiang, A., Yuen, M.M., Diaz-Chavez, M.L., Madilao, L.L., Finnegan, P.M., et al. 2016. Heartwood-specific transcriptome and metabolite signatures of tropical sandalwood (*Santalum album*) reveal the final step of (Z)-santalol fragrance biosynthesis. *Plant J.* 86(4): 289–299. doi:10.1111/tj.13162.
- Chakrabarty, R., Qu, Y., and Ro, D.K. 2015. Silencing the lettuce homologs of small rubber particle protein does not influence natural rubber biosynthesis in lettuce (*Lactuca sativa*). *Phytochemistry*, 113: 121–129. doi:10.1016/j.phytochem.2014.12.003.
- Chedgy, R.J., Kollner, T.G., and Constabel, C.P. 2015. Functional characterization of two acyltransferases from *Populus trichocarpa* capable of synthesizing benzyl benzoate and salicyl benzoate, potential intermediates in salicinoid phenolic glycoside biosynthesis. *Phytochemistry*, 113: 149–159. doi:10.1016/j.phytochem.2014.10.018.
- Chen, F., Tholl, D., Bohlmann, J., and Pichersky, E. 2011. The family of terpene synthases in plants: a mid-size family of genes for specialized metabolism that is highly diversified throughout the kingdom. *Plant J.* 66(1): 212–229. doi:10.1111/j.1365-3113.2011.04520.x.
- Chen, J., Zhan, J., Wang, H., Zhao, Y., Zhang, D., Chen, X., et al. 2023. VrMYB90 functions synergistically with VrbHLHA and VrMYB3 to regulate anthocyanin biosynthesis in mung bean. *Plant Cell Physiol.* 64(2): 221–233. doi:10.1093/pcp/pcac160.
- Chen, X., Hagel, J.M., Chang, L., Tucker, J.E., Shiigi, S.A., Yelapaala, Y., et al. 2018. A pathogenesis-related 10 protein catalyzes the final step in thebaïne biosynthesis. *Nat. Chem. Biol.* 14(7): 738–743. doi:10.1038/s41589-018-0059-7.
- Cheyrier, V., Comte, G., Davies, K.M., Lattanzio, V., and Martens, S. 2013. Plant phenolics: recent advances on their biosynthesis, genetics, and ecophysiology. *Plant Physiol. Biochem.* 72: 1–20. doi:10.1016/j.plaphy.2013.05.009.
- Cho, J.S., Nguyen, V.P., Jeon, H.W., Kim, M.H., Eom, S.H., Lim, Y.J., et al. 2016. Overexpression of *PtMYB119*, a R2R3-MYB transcription factor from *Populus trichocarpa*, promotes anthocyanin production in hybrid poplar. *Tree Physiol.* 36(9): 1162–1176. doi:10.1093/treephys/tpw046.
- Chung, M.K., and Campbell, J.N. 2016. Use of capsaicin to treat pain: mechanistic and therapeutic considerations. *Pharmaceuticals*, 9(4). doi:10.3390/ph9040066.
- Clark, S.M., Vaitheeswaran, V., Ambrose, S.J., Purves, R.W., and Page, J.E. 2013. Transcriptome analysis of bitter acid biosynthesis and precursor pathways in hop (*Humulus lupulus*). *BMC Plant Biol.* 13: 12. doi:10.1186/1471-2229-13-12.
- Colpitts, C.C., Kim, S.S., Posehn, S.E., Jepson, C., Kim, S.Y., Wiedemann, G., et al. 2011. PpASCL, a moss ortholog of anther-specific chalcone synthase-like enzymes, is a hydroxyalkylpyrone synthase involved in an evolutionarily conserved sporopollenin biosynthesis pathway. *New Phytol.* 192(4): 855–868. doi:10.1111/j.1469-8137.2011.03858.x.
- Constabel, C.P., Yip, L., Patton, J.J., and Christopher, M.E. 2000. Polyphenol oxidase from hybrid poplar. Cloning and expression in response to wounding and herbivory. *Plant Physiol.* 124(1): 285–295. doi:10.1104/pp.124.1.285.
- Cornish, K. 2014. Biosynthesis of natural rubber (NR) in different rubber-producing species. In *Chemistry, manufacture, and applications of natural rubber*. Edited by S. Kohjiya and Y. Ikeda. Woodhead Publishing. pp. 3–29.
- Corso, M., Perreau, F., Mouille, G., and Lepiniec, L. 2020. Specialized phenolic compounds in seeds: structures, functions, and regulations. *Plant Sci.* 296: 110471. doi:10.1016/j.plantsci.2020.110471.
- Costa, J., Sepulveda, M., Gallardo, V., Cayun, Y., Santander, C., Ruiz, A., et al. 2022. Antifungal potential of capsaicinoids and capsinoids from the *Capsicum* genus for the safeguarding of agrifood production: advantages and limitations for environmental health. *Microorganisms*, 10(12). doi:10.3390/microorganisms10122387.
- Cox, L.D., Munholland, S., Mats, L., Zhu, H., Crosby, W.L., Lukens, L., et al. 2021. The induction of the isoflavone biosynthesis pathway is associated with resistance to common bacterial blight in *Phaseolus vulgaris* L. *Metabolites*, 11(7). doi:10.3390/metabo11070433.
- da Silva-Santosa, A., Antunes, A., D'Avila, L., Bizzo, H., and Souza-Santos, L. 2005. Material study: the use of essential oils and terpenics/terpenoids in cosmetics and perfumery. Available from https://img.perfumerflavorist.com/files/base/allured/all/do-cument/2005/10/pf.PF_30_08_050_06.pdf [accessed 25 November 2024].
- Daku, R.M., Rabbi, F., Buttigieg, J., Coulson, I.M., Horne, D., Martens, G., et al. 2016. PpASCL, the *Physcomitrella patens* Anther-specific Chalcone synthase-Like Enzyme implicated in sporopollenin biosynthesis, is needed for integrity of the moss spore wall and spore viability. *PLoS One*, 11(1): e0146817. doi:10.1371/journal.pone.0146817.
- Dang, T.-T.T., Nguyen, T.-d., Tuan-ahn, Nguyen, M., and Leung, Y.Y. 2021. Cytochrome p450 monooxygenases and uses thereof. Edited by USPTO. University of British Columbia. US. p. 82.
- Dang, T.T., and Facchini, P.J. 2012. Characterization of three O-methyltransferases involved in noscapine biosynthesis in opium poppy. *Plant Physiol.* 159(2): 618–631. doi:10.1104/pp.112.194886.
- Dang, T.T., Chen, X., and Facchini, P.J. 2015. Acetylation serves as a protective group in noscapine biosynthesis in opium poppy. *Nat. Chem. Biol.* 11(2): 104–106. doi:10.1038/nchembio.1717.
- Dang, T.T., Onoyovwi, A., Farrow, S.C., and Facchini, P.J. 2012. Biochemical genomics for gene discovery in benzyloisoquinoline alkaloid biosynthesis in opium poppy and related species. *Methods Enzymol.* 515: 231–266. doi:10.1016/B978-0-12-394290-6.00011-2.
- Dang, T.T., Franke, J., Tatsis, E., and O'Connor, S.E. 2017. Dual catalytic activity of a cytochrome P450 controls bifurcation at a metabolic branch point of alkaloid biosynthesis in *Rauwolfia serpentina*. *Angew. Chem. Int. Ed.* 56(32): 9440–9444. doi:10.1002/anie.201705010.
- Dastmalchi, M. 2020a. A role for melatonin in the defense of sweet oranges against *Citrus* greening disease. *Plant Physiol.* 184(4): 1633–1634. doi:10.1104/pp.20.01463.
- Dastmalchi, M. 2020b. A well-oiled machine: two fatty acid exporters involved in seed oil biosynthesis. *Plant Physiol.* 182(4): 1798–1799. doi:10.1104/pp.20.00263.
- Dastmalchi, M. 2021a. Late-stage maturation erodes lipid content in soybean seeds. *Plant Physiol.* 186(2): 816–818. doi:10.1093/plphys/kiab129.
- Dastmalchi, M. 2021b. Elusive partners: a review of the auxiliary proteins guiding metabolic flux in flavonoid biosynthesis. *Plant J.* 108(2): 314–329. doi:10.1111/tj.15446.
- Dastmalchi, M., and Dhaubhadel, S. 2015. Soybean chalcone isomerase: evolution of the fold, and the differential expression and localization of the gene family. *Planta* 241(2): 507–523. doi:10.1007/s00425-014-2200-5.
- Dastmalchi, M., Chapman, P., Yu, J., Austin, R.S., and Dhaubhadel, S. 2017. Transcriptomic evidence for the control of soybean root isoflavonoid content by regulation of overlapping phenylpropanoid pathways. *BMC Genomics [Electronic Resource]*, 18(1): 70. doi:10.1186/s12864-016-3463-y.
- Dastmalchi, M., Chang, L., Torres, M.A., Ng, K.K.S., and Facchini, P.J. 2018. Codeinone reductase isoforms with differential stability, efficiency and product selectivity in opium poppy. *Plant J.* 95. doi:10.1111/tj.13975.
- Dastmalchi, M., Chang, L., Chen, R., Yu, L., Chen, X., Hagel, J.M., and Facchini, P.J. 2019a. Purine permease-type benzyloisoquinoline alkaloid transporters in opium poppy. *Plant Physiol.* 181(3): 916–933. doi:10.1104/pp.19.00565.
- Dastmalchi, M., Chen, X., Hagel, J.M., Chang, L., Chen, R., Ramasamy, S., et al. 2019b. Neopinone isomerase is involved in codeine and morphine biosynthesis in opium poppy. *Nat. Chem. Biol.* 15(4): 384–390. doi:10.1038/s41589-019-0247-0.
- De Carolis, E., and De Luca, V. 1993. Purification, characterization, and kinetic analysis of a 2-oxoglutarate-dependent dioxygenase involved in vindoline biosynthesis from *Catharanthus roseus*. *J. Biol. Chem.* 268(8): 5504–5511. doi:10.1016/S0021-9258(18)53349-4.
- De Carolis, E., and De Luca, V. 1994. A novel 2-oxoglutarate-dependent dioxygenase involved in vindoline biosynthesis: characterization, purification and kinetic properties. In *Primary and Secondary Metabolism of Plants and Cell Cultures III*. Edited by J. Schripsema and R. Verpoorte. Springer, Dordrecht. pp. 281–287.
- De Carolis, E., Chan, F., Balsevich, J., and De Luca, V. 1990. Isolation and characterization of a 2-oxoglutarate dependent dioxygenase involved in the second-to-last step in vindoline biosynthesis. *Plant Physiol.* 94(3): 1329. doi:10.1104/pp.94.3.1323.
- de la Rosa, L.A., Moreno-Escamilla, O., Rodrigo-García, J., and Alvarez-Parrilla, E. 2019. Phenolic compounds. In *Postharvest physiology and*

- biochemistry of fruits and vegetables. Edited by E.M. Yahia. Woodhead Publishing, Sawston, UK. pp. 253–271.
- De Luca, V., Marineau, C., and Brisson, N. 1989. Molecular cloning and analysis of cDNA encoding a plant tryptophan decarboxylase: comparison with animal dopa decarboxylases. *Proc. Natl. Acad. Sci.* 86(8): 2586. doi:10.1073/pnas.86.8.2582.
- De Luca, V., Salim, V., Thamm, A., Masada, S.A., and Yu, F. 2014. Making iridoids/secoiridoids and monoterpenoid indole alkaloids: progress on pathway elucidation. *Curr. Opin. Plant Biol.* 19: 42. doi:10.1016/j.pbi.2014.03.006.
- de Ronne, M., Lapierre, E., and Torkamaneh, D. 2024. Genetic insights into agronomic and morphological traits of drug-type cannabis revealed by genome-wide association studies. *Sci. Rep.* 14(1): 9162. doi:10.1038/s41598-024-58931-w.
- Dehnavi, M.M., Damerum, A., Taheri, S., Ebadi, A., Panahi, S., Hodgins, G., et al. 2025. Population genomics of a natural *Cannabis sativa* L. collection from Iran identifies novel genetic loci for flowering time, morphology, sex and chemotyping. *BMC Plant Biol.* 25(1): 80. doi:10.1186/s12870-025-06045-4.
- Demissie, Z., Woolfson, K.N., Yu, F., Qu, Y., and De Luca, V. 2017. The ATP binding cassette transporter, *VmTPT2/VmABCG1*, is involved in export of the monoterpenoid indole alkaloid, vincamine in *Vinca minor* leaves. *Phytochemistry*, 140: 118–124. doi:10.1016/j.phytochem.2017.04.019.
- Demissie, Z.A., Sarker, L.S., and Mahmoud, S.S. 2011. Cloning and functional characterization of β -phellandrene synthase from *Lavandula angustifolia*. *Planta*, 233(4): 685–696. doi:10.1007/s00425-010-1332-5.
- Demissie, Z.A., Erland, L.A., Rheault, M.R., and Mahmoud, S.S. 2013. The biosynthetic origin of irregular monoterpenes in *Lavandula*: isolation and biochemical characterization of a novel *cis*-prenyl diphosphate synthase gene, *lavandulyl diphosphate synthase*. *J. Biol. Chem.* 288(9): 6333–6341. doi:10.1074/jbc.M112.431171.
- Demissie, Z.A., Cella, M.A., Sarker, L.S., Thompson, T.J., Rheault, M.R., and Mahmoud, S.S. 2012. Cloning, functional characterization and genomic organization of 1,8-cineole synthases from *Lavandula*. *Plant Mol. Biol.* 79(4-5): 393–411. doi:10.1007/s11103-012-9920-3.
- Desgagné-Penix, I. 2021. Biosynthesis of alkaloids in Amaryllidaceae plants: a review. *Phytochem. Rev.* 20: 409–431. doi:10.1007/s11101-020-09678-5.
- Desgagné-Penix, I., Awwad, F., and Ghribi, M. 2023. Engineered microorganism for the production of cannabinoids.
- Dhaubhadel, S., McGarvey, B.D., Williams, R., and Gijzen, M. 2003. Isoflavonoid biosynthesis and accumulation in developing soybean seeds. *Plant Mol. Biol.* 53(6): 733–743. doi:10.1023/B:PLAN.0000023666.30358.ae.
- Dhyani, P., Quispe, C., Sharma, E., Bahukhandi, A., Sati, P., Attri, D.C., et al. 2022. Anticancer potential of alkaloids: a key emphasis to colchicine, vinblastine, vincristine, vindesine, vinorelbine and vincamine. *Cancer Cell Int.* 22(1). doi:10.1186/s12935-022-02624-9.
- Di Ferdinando, M., Brunetti, C., Fini, A., and Tattini, M. 2012. Flavonoids as antioxidants in plants under abiotic stresses. In *Abiotic Stress Responses in Plants*. Edited by P. Ahmad and M. Prasad. Springer, New York, NY.
- Diamond, A., Diaz-Garza, A.M., Li, J., Slaterry, S.S., Merindol, N., Fantino, E., et al. 2023. Instability of extrachromosomal DNA transformed into the diatom *Phaeodactylum tricornutum*. *Algal Res.* 70. doi:10.1016/j.algal.2023.102998.
- Dias, M.C., Pinto, D., Freitas, H., Santos, C., and Silva, A.M.S. 2020. The antioxidant system in *Olea europaea* to enhanced UV-B radiation also depends on flavonoids and secoiridoids. *Phytochemistry*, 170. doi:10.1016/j.phytochem.2019.112199.
- Diaz-Chavez, M.L., Moniodis, J., Madilao, L.L., Jancsik, S., Keeling, C.I., Barbour, E.L., et al. 2013. Biosynthesis of sandalwood oil: *Santalum album* CYP76F cytochromes P450 produce santalols and bergamotol. *PLoS One*, 8(9): e75053. doi:10.1371/journal.pone.0075053.
- Duwardi, K., Austin, R.S., Mainali, H.R., Bett, K., Marsolais, F., and Dhaubhadel, S. 2018. Slow darkening of pinto bean seed coat is associated with significant metabolite and transcript differences related to proanthocyanidin biosynthesis. *BMC Genomics* [Electronic Resource], 19(1): 260. doi:10.1186/s12864-018-4550-z.
- Dykstra, C.B., Pyne, M.E., and Martin, V.J. 2023. CRAPS: chromosomal-repair-assisted pathway shuffling in yeast. *ACS Synth. Biol.* 12(9): 2578–2587. doi:10.1021/acssynbio.3c00170.
- Edgar, S.M., Chatzivasileiou, A.O., Ward, V., and Stephanopoulos, G. 2021. Microbial engineering for the production of isoprenoids. Massachusetts Institute of Technology.
- Elmer, W.H., Mattina, M.J.I., and MacEachern, G.J. 1994. Sensitivity of plant pathogenic fungi to taxane extracts from ornamental yews. *Phytopathology*, 84(10): 1179–1185. doi:10.1094/Phyto-84-1179.
- Eng, J.G.M., Shahsavarani, M., Smith, D.P., Hajicek, J., De Luca, V., and Qu, Y. 2022. A *Catharanthus roseus* Fe(II)/ α -ketoglutarate-dependent dioxygenase catalyzes a redox-neutral reaction responsible for vindolinine biosynthesis. *Nat. Commun.* 13(1). doi:10.1038/s41467-022-31100-1.
- Facchini, P.J., and De Luca, V. 1994. Differential and tissue-specific expression of a gene family for tyrosine/dopa decarboxylase in opium poppy. *J. Biol. Chem.* 269(43): 26684–26690. Available from <https://www.ncbi.nlm.nih.gov/pubmed/7929401>. doi:10.1016/S0021-9258(18)47073-1.
- Facchini, P.J., and De Luca, V. 1995a. Phloem-specific expression of tyrosine/DOPA decarboxylase genes and the biosynthesis of isoquinoline alkaloids in opium poppy. *Plant Cell*, 7(11): 1811–1821. doi:10.1105/tpc.7.11.1811.
- Facchini, P.J., and De Luca, V. 1995b. Expression in *Escherichia coli* and partial characterization of two tyrosine/DOPA decarboxylases from opium poppy. *Phytochemistry*, 38(5): 1119–1126. doi:10.1016/0031-9422(94)00814-a.
- Facchini, P.J., Johnson, A.G., Poupart, J., and de Luca, V. 1996a. Uncoupled defense gene expression and antimicrobial alkaloid accumulation in elicited opium poppy cell cultures. *Plant Physiol.* 111(3): 687–697. doi:10.1104/pp.111.3.687.
- Facchini, P.J., Penzes, C., Johnson, A.G., and Bull, D. 1996b. Molecular characterization of berberine bridge enzyme genes from opium poppy. *Plant Physiol.* 112(4): 1669–1677. doi:10.1104/pp.112.4.1669.
- Facchini, P.J., Penzes-Yost, C., Samanani, N., and Kowalchuk, B. 1998. Expression patterns conferred by tyrosine/dihydroxyphenylalanine decarboxylase promoters from opium poppy are conserved in transgenic tobacco. *Plant Physiol.* 118(1): 69–81. doi:10.1104/pp.118.1.69.
- Faldt, J., Arimura, G., Gershenzon, J., Takabayashi, J., and Bohlmann, J. 2003a. Functional identification of *AtTPS03* as (E)- β -ocimene synthase: a monoterpene synthase catalyzing jasmonate- and wound-induced volatile formation in *Arabidopsis thaliana*. *Planta*, 216(5): 745–751. doi:10.1007/s00425-002-0924-0.
- Faldt, J., Martin, D., Miller, B., Rawat, S., and Bohlmann, J. 2003b. Traumatic resin defense in Norway spruce (*Picea abies*): methyl jasmonate-induced terpene synthase gene expression, and cDNA cloning and functional characterization of (+)-3-carene synthase. *Plant Mol. Biol.* 51(1): 119–133. doi:10.1023/a:1020714403780.
- Fantino, E., Awwad, F., Merindol, N., Diaz Garza, A.M., Gélinas, S.-E., Gajón Robles, G.C., et al. 2024. Bioengineering *phaeodactylum tricornutum*, a marine diatom, for cannabinoid biosynthesis. *Algal Res.* 77. doi:10.1016/j.algal.2023.103379.
- Farrell, K., Jahan, M.A., and Kovinich, N. 2017. Distinct mechanisms of biotic and chemical elicitors enable additive elicitation of the anticancer phytoalexin glyceollin I. *Molecules*, 22(8). doi:10.3390/molecules22081261.
- Fellenberg, C., Corea, O., Yan, L.H., Archinuk, F., Piirtola, E.M., Gordon, H., et al. 2020. Discovery of salicyl benzoate UDP-glycosyltransferase, a central enzyme in poplar salicinoid phenolic glycoside biosynthesis. *Plant J.* 102(1): 99–115. doi:10.1111/tjp.14615.
- Ferraro, K., Jin, A.L., Nguyen, T.D., Reinecke, D.M., Ozga, J.A., and Ro, D.K. 2014. Characterization of proanthocyanidin metabolism in pea (*Pisum sativum*) seeds. *BMC Plant Biol.* 14: 238. doi:10.1186/s12870-014-0238-y.
- Fossati, E., Ekins, A., Narcross, L., Zhu, Y., Falgoutyret, J.-P., Beaudoin, G.A., et al. 2014. Reconstitution of a 10-gene pathway for synthesis of the plant alkaloid dihydrosanguinarine in *Saccharomyces cerevisiae*. *Nat. Commun.* 5(1). doi:10.1038/ncomms4283.
- Fradj, N., Gonçalves Dos Santos, K.C., de Montigny, N., Awwad, F., Boumghar, Y., Germain, H., and Desgagne-Penix, I. 2019. RNA-seq de novo assembly and differential transcriptome analysis of Chaga (*Inonotus obliquus*) cultured with different betulin sources and the regulation of genes involved in terpenoid biosynthesis. *Int. J. Mol. Sci.* 20(18). doi:10.3390/ijms20184334.
- Fradj, N., de Montigny, N., Merindol, N., Awwad, F., Boumghar, Y., Germain, H., and Desgagne-Penix, I. 2020. A first insight into north amer-

- ican plant pathogenic fungi *Armillaria sinapina* transcriptome. *Biology*, 9(7). doi:10.3390/biology9070153.
- Freixas Coutin, J.A., Munholland, S., Silva, A., Subedi, S., Lukens, L., Crosby, W.L., et al. 2017. Proanthocyanidin accumulation and transcriptional responses in the seed coat of cranberry beans (*Phaseolus vulgaris* L.) with different susceptibility to postharvest darkening. *BMC Plant Biol.* 17(1): 89. doi:10.1186/s12870-017-1037-z.
- Fukuma, K., Neuls, E.D., Ryberg, J.M., Suh, D.Y., and Sankawa, U. 2007. Mutational analysis of conserved outer sphere arginine residues of chalcone synthase. *J. Biochem.* 142(6): 731–739. doi:10.1093/jb/mvm188.
- Gagne, S.J., Stout, J.M., Liu, E., Boubakir, Z., Clark, S.M., and Page, J.E. 2012. Identification of olivetolic acid cyclase from *Cannabis sativa* reveals a unique catalytic route to plant polyketides. *Proc. Natl. Acad. Sci.* 109(31): 12811–12816. doi:10.1073/pnas.1200330109.
- Galata, M., Sarker, L.S., and Mahmoud, S.S. 2014. Transcriptome profiling, and cloning and characterization of the main monoterpene synthases of *Coriandrum sativum* L. *Phytochemistry*, 102: 64–73. doi:10.1016/j.phytochem.2014.02.016.
- Gary, S., Adegboye, J., Popp, B., Cocuron, J.C., Woodrum, B., and Kovich, N. 2018. Combining semi-synthesis with plant and microbial biocatalysis: new frontiers in producing a chemical arsenal against cancer. *RSC Adv.* 8(38): 21332–21339. doi:10.1039/c8ra02184h.
- Geisler, K., Jensen, N.B., Yuen, M.M., Madilao, L., and Bohlmann, J. 2016. Modularity of conifer diterpene resin acid biosynthesis: P450 enzymes of different CYP720B clades use alternative substrates and converge on the same products. *Plant Physiol.* 171(1): 152–164. doi:10.1104/pp.16.00180.
- Geissler, M., Volk, J., Stehle, F., Kayser, O., and Warzecha, H. 2018. Subcellular localization defines modification and production of Δ^9 -tetrahydrocannabinolic acid synthase in transiently transformed *Nicotiana benthamiana*. *Biotechnol. Lett.* 40(6): 981–987. doi:10.1007/s10529-018-2545-0.
- Georgiev, V., Ivanov, I., and Pavlov, A. 2020. Recent progress in Amaryllidaceae biotechnology. *Molecules*, 25(20). doi:10.3390/molecules25204670.
- Gesell, A., Yoshida, K., Tran, L.T., and Constabel, C.P. 2014. Characterization of an apple TT2-type R2R3 MYB transcription factor functionally similar to the poplar proanthocyanidin regulator PtMYB134. *Planta*, 240(3): 497–511. doi:10.1007/s00425-014-2098-y.
- Gesell, A., Blaukopf, M., Madilao, L., Yuen, M.M., Withers, S.G., Mattsson, J., et al. 2015. The gymnosperm cytochrome P450 CYP750B1 catalyzes stereospecific monoterpene hydroxylation of (+)-sabinene in thujone biosynthesis in western redcedar. *Plant Physiol.* 168(1): 94–106. doi:10.1104/pp.15.00315.
- Gourlay, G., Ma, D., Schmidt, A., and Constabel, C.P. 2020. MYB134-RNAi poplar plants show reduced tannin synthesis in leaves but not roots, and increased susceptibility to oxidative stress. *J. Exp. Bot.* 71(20): 6601–6611. doi:10.1093/jxb/eraa371.
- Greer, S., Wen, M., Bird, D., Wu, X., Samuels, L., Kunst, L., and Jetter, R. 2007. The cytochrome P450 enzyme CYP96A15 is the midchain alkane hydroxylase responsible for formation of secondary alcohols and ketones in stem cuticular wax of *Arabidopsis*. *Plant Physiol.* 145(3): 653–667. doi:10.1104/pp.107.107300.
- Guhling, O., Hobl, B., Yeats, T., and Jetter, R. 2006. Cloning and characterization of a lupeol synthase involved in the synthesis of epicuticular wax crystals on stem and hypocotyl surfaces of *Ricinus communis*. *Arch. Biochem. Biophys.* 448(1-2): 60–72. doi:10.1016/j.abb.2005.12.013.
- Gülck, T., Booth, J.K., Carvalho, Â., Khakimov, B., Crocoll, C., Motawia, M.S., et al. 2020. Synthetic biology of cannabinoids and cannabinoid glucosides in *Nicotiana benthamiana* and *Saccharomyces cerevisiae*. *J. Nat. Prod.* 83(10): 2877–2893. doi:10.1021/acs.jnatprod.0c00241.
- Guo, D., Wang, H., Zhang, S., and Lan, T. 2022. The type III polyketide synthase supergene family in plants: complex evolutionary history and functional divergence. *Plant J.* 112(2): 414–428. doi:10.1111/tjp.15953.
- Guo, J., Gao, D., Lian, J., and Qu, Y. 2024. De novo biosynthesis of antiarrhythmic alkaloid ajmaline. *Nat. Commun.* 15(1): 457. doi:10.1038/s41467-024-44797-z.
- Gupta, A.K., Akhtar, T.A., Widmer, A., Pichersky, E., and Schiestl, F.P. 2012. Identification of white campion (*Silene latifolia*) guaiacol O-methyltransferase involved in the biosynthesis of veratrole, a key volatile for pollinator attraction. *BMC Plant Biol.* 12: 158. doi:10.1186/1471-2229-12-158.
- Hagel, J.M., and Facchini, P.J. 2010. Dioxygenases catalyze the O-demethylation steps of morphine biosynthesis in opium poppy. *Nat. Chem. Biol.* 6(4): 273–275. doi:10.1038/nchembio.317.
- Hagel, J.M., Beaudoin, G.A., Fossati, E., Ekins, A., Martin, V.J., and Facchini, P.J. 2012. Characterization of a flavoprotein oxidase from opium poppy catalyzing the final steps in sanguinarine and papaverine biosynthesis. *J. Biol. Chem.* 287(51): 42983. doi:10.1074/jbc.M112.420414.
- Hall, D., and De Luca, V. 2007. Mesocarp localization of a bi-functional resveratrol/hydroxycinnamic acid glucosyltransferase of Concord grape (*Vitis labrusca*). *Plant J.* 49(4): 579–591. doi:10.1111/j.1365-313X.2006.02987.x.
- Hall, D., Kim, K.H., and De Luca, V. 2011a. Molecular cloning and biochemical characterization of three Concord grape (*Vitis labrusca*) flavonol 7-O-glucosyltransferases. *Planta*, 234(6): 1201–1214. doi:10.1007/s00425-011-1474-0.
- Hall, D.E., Robert, J.A., Keeling, C.I., Domanski, D., Quesada, A.L., Jancsik, S., et al. 2011b. An integrated genomic, proteomic and biochemical analysis of (+)-3-carene biosynthesis in Sitka spruce (*Picea sitchensis*) genotypes that are resistant or susceptible to white pine weevil. *Plant J.* 65(6): 936–948. doi:10.1111/j.1365-313X.2010.04478.x.
- Hall, D., Yuan, X.X., Murata, J., and De Luca, V. 2012. Molecular cloning and biochemical characterization of the UDP-glucose: flavonoid 3-O-glucosyltransferase from Concord grape (*Vitis labrusca*). *Phytochemistry*, 74: 90–99. doi:10.1016/j.phytochem.2011.10.007.
- Hamberger, B., Ohnishi, T., Hamberger, B., Seguin, A., and Bohlmann, J. 2011. Evolution of diterpene metabolism: Sitka spruce CYP720B4 catalyzes multiple oxidations in resin acid biosynthesis of conifer defense against insects. *Plant Physiol.* 157(4): 1677–1695. doi:10.1104/pp.111.185843.
- Han, Y., Zhao, W., Wang, Z., Zhu, J., and Liu, Q. 2014. Molecular evolution and sequence divergence of plant chalcone synthase and chalcone synthase-like genes. *Genetica*, 142(3): 215–225. doi:10.1007/s10709-014-9768-3.
- Haruta, M., Pedersen, J.A., and Constabel, C.P. 2001. Polyphenol oxidase and herbivore defense in trembling aspen (*Populus tremuloides*): cDNA cloning, expression, and potential substrates. *Physiol. Plant.* 112(4): 552–558. doi:10.1034/j.1399-3054.2001.1120413.x.
- He, F., Shi, Y.J., Zhao, Q., Zhao, K.J., Cui, X.L., Chen, L.H., et al. 2021. Genome-wide investigation and expression profiling of polyphenol oxidase (PPO) family genes uncover likely functions in organ development and stress responses in *Populus trichocarpa*. *BMC Genomics* [Electronic Resource], 22(1): 731. doi:10.1186/s12864-021-08028-9.
- Hegebarth, D., Buschhaus, C., Joubes, J., Thoraval, D., Bird, D., and Jetter, R. 2017. Arabidopsis ketoacyl-CoA synthase 16 (KCS16) forms C₃₆/C₃₈ acyl precursors for leaf trichome and pavement surface wax. *Plant Cell Environ.* 40(9): 1761–1776. doi:10.1111/pce.12981.
- Hodgins, C.L., Salama, E.M., Kumar, R., Zhao, Y., Roth, S.A., Cheung, I.Z., et al. 2024. Creating saponin-free yellow pea seeds by CRISPR/Cas9-enabled mutagenesis on β -amyrin synthase. *Plant Direct*, 8(1). doi:10.1002/pld3.563.
- Hotchandani, T., de Villiers, J., and Desgagne-Penix, I. 2019. Developmental regulation of the expression of amaryllidaceae alkaloid biosynthetic genes in *Narcissus papyraceus* [Research Support, Non-U.S. Gov't]. *Genes*, 10(8). doi:10.3390/genes10080594.
- Huang, A.C., and Osbourn, A. 2019. Plant terpenes that mediate below-ground interactions: prospects for bioengineering terpenoids for plant protection. *Pest Manage. Sci.* 75(9): 2368–2377. doi:10.1002/ps.5410.
- Huber, D.P., Philippe, R.N., Godard, K.A., Sturrock, R.N., and Bohlmann, J. 2005. Characterization of four terpene synthase cDNAs from methyl jasmonate-induced Douglas-fir, *Pseudotsuga menziesii*. *Phytochemistry*, 66(12): 1427–1439. doi:10.1016/j.phytochem.2005.04.030.
- Hurd, M.C., Kwon, M., and Ro, D.K. 2017. Functional identification of a *Lippia dulcis* bornyl diphosphate synthase that contains a duplicated, inhibitory arginine-rich motif. *Biochem. Biophys. Res. Commun.* 490(3): 963–968. doi:10.1016/j.bbrc.2017.06.147.
- Irmisch, S., Jancsik, S., Yuen, M.M.S., Madilao, L.L., and Bohlmann, J. 2019. Biosynthesis of the anti-diabetic metabolite montbretin A: glucosylation of the central intermediate mini-MbA. *Plant J.* 100(5): 879–891. doi:10.1111/tjp.14493.

- Irmisch, S., Jancsik, S., Man Saint Yuen, M., Madilao, L.L., and Bohlmann, J. 2020. Complete biosynthesis of the anti-diabetic plant metabolite montbretin A. *Plant Physiol.* 184(1): 97–109. doi:10.1104/pp.20.00522.
- Irmisch, S., Jo, S., Roach, C.R., Jancsik, S., Man Saint Yuen, M., Madilao, L.L., et al. 2018. Discovery of UDP-glycosyltransferases and BAHD-acyltransferases involved in the biosynthesis of the antidiabetic plant metabolite montbretin A. *Plant Cell*, 30(8): 1864–1886. doi:10.1105/tpc.18.00406.
- Islam, N.S., and Dhaubhadel, S. 2023. Proanthocyanidin biosynthesis and postharvest seed coat darkening in pinto bean. *Phytochem. Rev.* doi:10.1007/s11101-023-09895-8.
- Islam, N.S., Duwadi, K., Chen, L., Pajak, A., McDowell, T., Marsolais, F., and Dhaubhadel, S. 2022. Global analysis of common bean multidrug and toxic compound extrusion transporters (PvMATEs): PvMATE8 and pinto bean seed coat darkening. *Front. Plant Sci.* 13: 1046597. doi:10.3389/fpls.2022.1046597.
- Jahan, M.A., and Kavinich, N. 2019. Acidity stress for the systemic elicitation of glyceollin phytoalexins in soybean plants. *Plant. Signal. Behav.* 14(7): 1604018. doi:10.1080/15592324.2019.1604018.
- Jahan, M.A., Harris, B., Lowery, M., Infante, A.M., Percifield, R.J., and Kavinich, N. 2020. Glyceollin transcription factor GmMYB29A2 regulates soybean resistance to *Phytophthora sojae*. *Plant Physiol.* 183(2): 530–546. doi:10.1104/pp.19.01293.
- Jalali, S., Salami, S.A., Sharifi, M., and Sohrabi, S. 2019. Signaling compounds elicit expression of key genes in cannabinoid pathway and related metabolites in cannabis. *Ind. Crops Prod.* 133: 105–110. doi:10.1016/j.indcrop.2019.03.004.
- James, A.M., Ma, D., Mellway, R., Gesell, A., Yoshida, K., Walker, V., et al. 2017. Poplar MYB115 and MYB134 transcription factors regulate proanthocyanidin synthesis and structure. *Plant Physiol.* 174(1): 154–171. doi:10.1104/pp.16.01962.
- Jaramillo-Madrid, A.C., Lacchini, E., and Gossens, A. 2022. Within and beyond organelle engineering: strategies for increased terpene production in yeast and plants. *Curr. Opin. Green Sustain. Chem.* 33: 100572. doi:10.1016/j.cogsc.2021.100572.
- Jepson, C., Karppinen, K., Daku, R.M., Sterenberg, B.T., and Suh, D.Y. 2014. *Hypericum perforatum* hydroxyalkylpyrone synthase involved in sporopollenin biosynthesis—Phylogeny, site-directed mutagenesis, and expression in nonanther tissues. *FEBS J.* 281(17): 3855–3868. doi:10.1111/febs.12920.
- Jiang, C., Kim, S.Y., and Suh, D.Y. 2008. Divergent evolution of the thiolase superfamily and chalcone synthase family. *Mol. Phylogenet. Evol.* 49(3): 691–701. doi:10.1016/j.ympev.2008.09.002.
- Jiang, C., Schommer, C.K., Kim, S.Y., and Suh, D.Y. 2006. Cloning and characterization of chalcone synthase from the moss *Physcomitrella patens*. *Phytochemistry*, 67(23): 2531–2540. doi:10.1016/j.phytochem.2006.09.030.
- Juca, M.M., Cysne Filho, F.M.S., de Almeida, J.C., Mesquita, D.D.S., Bariga, J.R.M., Dias, K.C.F., et al. 2020. Flavonoids: biological activities and therapeutic potential. *Nat. Prod. Res.* 34(5): 705. doi:10.1080/14786419.2018.1493588.
- Kajikawa, M., Yamato, K.T., Fukuzawa, H., Sakai, Y., Uchida, H., and Ohyama, K. 2005. Cloning and characterization of a cDNA encoding β -amyrin synthase from petroleum plant *Euphorbia tirucalli* L. *Phytochemistry*, 66(15): 1759–1766. doi:10.1016/j.phytochem.2005.05.021.
- Karimzadegan, V., Koirala, M., Sobhanverdi, S., Merindol, N., Majhi, B.B., Gélina, S.-E., et al. 2024. Characterization of cinnamate 4-hydroxylase (CYP73A) and p-coumaroyl 3'-hydroxylase (CYP98A) from *Leucojum aestivum*, a source of Amaryllidaceae alkaloids. *Plant Physiol. Biochem.* 210. doi:10.1016/j.plaphy.2024.108612.
- Kaur, J., Melkani, I., Singh, A.P., Singh, A.P., and Bala, K. 2022. Galantamine: a review update. *J. Drug Deliv Ther.* 12(4): 173. doi:10.22270/jddt.v12i4.5426.
- Keasling, J.D., Martin, V.J., Pitera, D.J., Withers, S.T., and Newman, J. 2007. Biosynthesis of amorpho-4,11-diene. Edited by USPTO. University of California, USA.
- Keasling, J.D., Martin, V.J., Pitera, D.J., Kim, S.-W., Withers, S.T., Yoshikuni, Y., et al. 2011. Host cells for production of isoprenoid compounds. Edited by USPTO. University of California, USA.
- Keeling, C.I., Madilao, L.L., Zerbe, P., Dullat, H.K., and Bohlmann, J. 2011a. The primary diterpene synthase products of *Picea abies* levopimaradiene/abietadiene synthase (PaLAS) are epimers of a thermally unstable diterpenol. *J. Biol. Chem.* 286(24): 21145–21153. doi:10.1074/jbc.M111.245951.
- Keeling, C.I., Weisshaar, S., Ralph, S.G., Jancsik, S., Hamberger, B., Dullat, H.K., and Bohlmann, J. 2011b. Transcriptome mining, functional characterization, and phylogeny of a large terpene synthase gene family in spruce (*Picea* spp.). *BMC Plant Biol.* 11: 43. doi:10.1186/1471-2229-11-43.
- Keeling, C.I., Dullat, H.K., Yuen, M., Ralph, S.G., Jancsik, S., and Bohlmann, J. 2010. Identification and functional characterization of monofunctional *ent*-copalyl diphosphate and *ent*-kaurene synthases in white spruce reveal different patterns for diterpene synthase evolution for primary and secondary metabolism in gymnosperms. *Plant Physiol.* 152(3): 1197–1208. doi:10.1104/pp.109.151456.
- Kempthorne, C.J., St Pierre, M., Le, A., Livingstone, S., McNulty, J., Cadotte, M.W., and Liscombe, D.K. 2024. Mass spectrometry-based metabolomics for the elucidation of alkaloid biosynthesis and function in invasive *Vincetoxicum rossicum* populations. *Phytochemistry*, 221: 114051. doi:10.1016/j.phytochem.2024.114051.
- Khatri, P., Wally, O., Rajcan, I., and Dhaubhadel, S. 2022. Comprehensive analysis of cytochrome P450 monooxygenases reveals insight into their role in partial resistance against *Phytophthora sojae* in soybean. *Front. Plant Sci.* 13: 862314. doi:10.3389/fpls.2022.862314.
- Khatri, P., Chen, L., Rajcan, I., and Dhaubhadel, S. 2023. Functional characterization of cinnamate 4-hydroxylase gene family in soybean (*Glycine max*). *PLoS One*, 18(5): e0285698. doi:10.1371/journal.pone.0285698.
- Kidd, T., Easson, M.L., Qu, Y., and De Luca, V. 2019. Inter-organ transport of secologanin allows assembly of monoterpenoid indole alkaloids in a *Catharanthus roseus* mutant. *Phytochemistry*, 159: 119–126. doi:10.1016/j.phytochem.2018.12.017.
- Kilpatrick, K., Pajak, A., Hagel, J.M., Sumarah, M.W., Lewinsohn, E., Facchini, P.J., and Marsolais, F. 2016. Characterization of aromatic aminotransferases from *Ephedra sinica* Stapf. *Amino Acids*, 48(5): 1209–1220. doi:10.1007/s00726-015-2156-1.
- Kim, A.L., Yun, Y.J., Choi, H.W., Hong, C.H., Shim, H.J., Lee, J.H., and Kim, Y.C. 2022. Profiling cannabinoid contents and expression levels of corresponding biosynthetic genes in commercial cannabis (*Cannabis sativa* L.) cultivars. *Plants*, 11(22):. doi:10.3390/plants11223088.
- Kim, K., Shahsavari, M., Garza-Garcia, J.J.O., Carlisle, J.E., Guo, J., De Luca, V., and Qu, Y. 2023. Biosynthesis of kratom opioids. *New Phytol.* 240(2): 757–769. doi:10.1111/nph.19162.
- Kim, O.T., Um, Y., Jin, M.L., Kim, J.U., Hegebarth, D., Busta, L., et al. 2018. A novel multifunctional C-23 oxidase, CYP714E19, is involved in asiaticoside biosynthesis. *Plant Cell Physiol.* 59(6): 1200–1213. doi:10.1093/pcp/pcy055.
- Kim, S.Y., Colpitts, C.C., Wiedemann, G., Jepson, C., Rahimi, M., Rothwell, J.R., et al. 2013. *Physcomitrella* PpORS, basal to plant type III polyketide synthases in phylogenetic trees, is a very long chain 2'-oxoalkylresorcinol synthase. *J. Biol. Chem.* 288(4): 2767–2777. doi:10.1074/jbc.M112.430686.
- Koduri, P.K., Gordon, G.S., Barker, E.I., Colpitts, C.C., Ashton, N.W., and Suh, D.Y. 2010. Genome-wide analysis of the chalcone synthase superfamily genes of *Physcomitrella patens*. *Plant Mol. Biol.* 72(3): 247–263. doi:10.1007/s11103-009-9565-z.
- Koirala, M., Cristine Gonçalves Dos Santos, K., Gelinas, S.E., Ricard, S., Karimzadegan, V., Lamichhane, B., et al. 2023. Auxin and light-mediated regulation of growth, morphogenesis, and alkaloid biosynthesis in *Crinum x powellii* 'Album' callus. *Phytochemistry*, 216: 113883. doi:10.1016/j.phytochem.2023.113883.
- Kojima, N., Sitthithaworn, W., Viroonchatapan, E., Suh, D.Y., Iwanami, N., Hayashi, T., and Sankaw, U. 2000. Geranylgeranyl diphosphate synthases from *Scoparia dulcis* and *Croton sublyratus*. cDNA cloning, functional expression, and conversion to a farnesyl diphosphate synthase. *Chem. Pharm. Bull.* 48(7): 1101–1103. doi:10.1248/cpb.48.1101.
- Kopaczky, J.M., Wargula, J., and Jelonek, T. 2020. The variability of terpenes in conifers under developmental and environmental stimuli. *Environ. Exp. Bot.* 180: 104197. doi:10.1016/j.envexpbot.2020.104197.
- Kosonen, M., Keski-Saari, S., Ruuhola, T., Constabel, C.P., and Julkunen-Tiitto, R. 2012. Effects of overproduction of condensed tannins and elevated temperature on chemical and ecological traits of genetically modified hybrid aspens (*Populus tremula* x *P. tremuloides*). *J. Chem. Ecol.* 38(10): 1235–1246. doi:10.1007/s10886-012-0193-8.

- Kovinich, N., and Durkin, P. 2018. Hormone deficient mutants have distinct flavonoid proportion fingerprints in response to abiotic stress. *Plant. Signal. Behav.* 13(12): e1542241. doi:10.1080/15592324.2018.1542241.
- Kulagina, N., Meteignier, L.V., Papon, N., O'Connor, S.E., and Courdavault, V. 2022. More than a *Catharanthus* plant: a multicellular and pluri-organelle alkaloid-producing factory. *Curr. Opin. Plant Biol.* 67: 102200. doi:10.1016/j.pbi.2022.102200.
- Kwan, B.D., Seligmann, B., Nguyen, T.D., Franke, J., and Dang, T.T. 2023. Leveraging synthetic biology and metabolic engineering to overcome obstacles in plant pathway elucidation. *Curr. Opin. Plant Biol.* 71: 102330. doi:10.1016/j.pbi.2022.102330.
- Kwon, M., Cochrane, S.A., Vederas, J.C., and Ro, D.K. 2014. Molecular cloning and characterization of drimenol synthase from valerian plant (*Valeriana officinalis*). *FEBS Lett.* 588(24): 4597–4603. doi:10.1016/j.febslet.2014.10.031.
- Kwon, M., Hodgins, C.L., Haslam, T.M., Roth, S.A., Nguyen, T.D., Yeung, E.C., and Ro, D.K. 2022a. Germacrene synthases for sesquiterpene lactone biosynthesis are expressed in vascular parenchyma cells neighboring laticifers in lettuce. *Plants*, 11(9). doi:10.3390/plants11091192.
- Kwon, M., Utomo, J.C., Park, K., Pascoe, C.A., Chiorean, S., Ngo, I., et al. 2022b. Cytochrome P450-catalyzed biosynthesis of a dihydrofuran neoclerodane in magic mint (*Salvia divinorum*). *ACS Catalysis*, 12(1): 777–782. doi:10.1021/acscatal.1c03691.
- Kwon, M., Hodgins, C.L., Salama, E.M., Dias, K.R., Parikh, A., Mackey, A.V., et al. 2023. New insights into natural rubber biosynthesis from rubber-deficient lettuce mutants expressing goldenrod or guayule cis-prenyltransferase. *New Phytol.* 239(3): 1098–1111. doi:10.1111/nph.18994.
- Lackus, N.D., Muller, A., Krober, T.D.U., Reichelt, M., Schmidt, A., Nakamura, Y., et al. 2020. The occurrence of sulfated salicinoids in poplar and their formation by sulfotransferase. *Plant Physiol.* 183(1): 137–151. doi:10.1104/pp.19.01447.
- Lakusta, A.M., Kwon, M., Kwon, E.G., Stonebloom, S., Scheller, H.V., and Ro, D.K. 2019. Molecular studies of the protein complexes involving cis-prenyltransferase in guayule (*Parthenium argentatum*), an alternative rubber-producing plant. *Front. Plant Sci.* 10: 165. doi:10.3389/fpls.2019.00165.
- Lang, D., Ullrich, K.K., Murat, F., Fuchs, J., Jenkins, J., Haas, F.B., et al. 2018. The *Physcomitrella patens* chromosome-scale assembly reveals moss genome structure and evolution. *Plant J.* 93(3): 515–533. doi:10.1111/tpl.13801.
- Lang, D.E., Morris, J.S., Rowley, M., Torres, M.A., Maksimovich, V.A., Facchini, P.J., and Ng, K.K.S. 2019. Structure-function studies of tetrahydroprotoberberine N-methyltransferase reveal the molecular basis of stereoselective substrate recognition. *J. Biol. Chem.* 294(40): 14482–14498. doi:10.1074/jbc.RA119.009214.
- Lau, N.S., Makita, Y., Kawashima, M., Taylor, T.D., Kondo, S., Othman, A.S., et al. 2016. The rubber tree genome shows expansion of gene family associated with rubber biosynthesis. *Sci. Rep.* 6: 28594. doi:10.1038/srep28594.
- Lee, A.-R., Kwon, M., Kang, M.-K., Kim, J., Kim, S.-U., and Ro, D.-K. 2019. Increased sesqui- and triterpene production by co-expression of HMG-CoA reductase and biotin carboxyl carrier protein in tobacco (*Nicotiana benthamiana*). *Metab. Eng.* 52: 20–28. doi:10.1016/j.ymben.2018.10.008.
- Lee, E.J., and Facchini, P. 2010. Norcoclaurine synthase is a member of the pathogenesis-related 10/bet v1 protein family. *Plant Cell*, 22(10): 3489–3503. doi:10.1105/tpc.110.07958.
- Lee, E.J., and Facchini, P.J. 2011. Tyrosine aminotransferase contributes to benzyloquinoline alkaloid biosynthesis in opium poppy. *Plant Physiol.* 157(3): 1067–1078. doi:10.1104/pp.111.185512.
- Levac, D., Murata, J., Kim, W.S., and De Luca, V. 2008. Application of carborundum abrasion for investigating the leaf epidermis: molecular cloning of *Catharanthus roseus* 16-hydroxytabersonine-16-O-methyltransferase. *Plant J.* 53(2): 236. doi:10.1111/j.1365-313X.2007.03337.x.
- Levac, D., Cazares, P., Yu, F., and De Luca, V. 2016. A picrinine N-methyltransferase belongs to a new family of γ -tocopherol-like methyltransferases found in medicinal plants that make biologically active monoterpene indole alkaloids. *Plant Physiol.* 170(4): 1935–1944. doi:10.1104/pp.15.01813.
- Li, C., Yu, W., Xu, J., Lu, X., and Liu, Y. 2022a. Anthocyanin biosynthesis induced by MYB transcription factors in plants. *Int. J. Mol. Sci.* 23(19). doi:10.3390/ijms231911701.
- Li, F., Bordeleau, S., Kim, K.H., Turcotte, J., Davis, B., Liu, L., et al. 2022b. A lesion-mimic mutant of *Catharanthus roseus* accumulates the opioid agonist, akummicine. *Phytochemistry*, 203: 113422. doi:10.1016/j.phytochem.2022.113422.
- Li, C., Gao, Z., Hu, W., Zhu, X., Li, Y., Li, N., and Ma, C. 2023. Integration of comparative transcriptomics and WGCNA characterizes the regulation of anthocyanin biosynthesis in mung bean (*Vigna radiata* L.). *Front. Plant Sci.* 14: 1251464. doi:10.3389/fpls.2023.1251464.
- Li, F., Shahsavarani, M., Handy-Hart, C.-J., Montgomery, V., Beech, R.N., Liu, L., et al. 2024. Characterization of a vacuolar importer of secologanin in *Catharanthus roseus*. *Commun. Biol.* 7: 939. doi:10.1038/s42003-024-06624-5.
- Li, L., Aslam, M., Rabbi, F., Vanderwel, M.C., Ashton, N.W., and Suh, D.Y. 2018. PpORS, an ancient type III polyketide synthase, is required for integrity of leaf cuticle and resistance to dehydration in the moss *Physcomitrella patens*. *Planta*, 247(2): 527–541. doi:10.1007/s00425-017-2806-5.
- Li, Q., Ramasamy, S., Singh, P., Hagel, J.M., Dunemann, S.M., Chen, X., et al. 2020. Gene clustering and copy number variation in alkaloid metabolic pathways of opium poppy. *Nat. Commun.* 11(1): 1190. doi:10.1038/s41467-020-15040-2.
- Li, X., and Dhaubhadel, S. 2012. 14-3-3 proteins act as scaffolds for GmMYB62 and GmMYB176 and regulate their intracellular localization in soybean. *Plant. Signal. Behav.* 7(8): 965–968. doi:10.4161/psb.20940.
- Lin, J., Wi, D., Ly, M., Jahan, M.A., Pullano, S., Martirosyan, I., and Kovinich, N. 2023a. Soybean hairy root transformation for the analysis of gene function. *J. Vis. Exp.*(195). doi:10.3791/65485.
- Lin, J., Monsalvo, I., Ly, M., Jahan, M.A., Wi, D., Martirosyan, I., and Kovinich, N. 2023b. RNA-seq dissects incomplete activation of phytoalexin biosynthesis by the soybean transcription factors GmMYB29A2 and GmNAC42-1. *Plants*, 12(3). doi:10.3390/plants12030545.
- Liscombe, D.K., and Facchini, P.J. 2007. Molecular cloning and characterization of tetrahydroprotoberberine cis-N-methyltransferase, an enzyme involved in alkaloid biosynthesis in opium poppy. *J. Biol. Chem.* 282(20): 14741–14751. doi:10.1074/jbc.M611908200.
- Liscombe, D.K., Macleod, B.P., Loukanina, N., Nandi, O.I., and Facchini, P.J. 2005. Evidence for the monophyletic evolution of benzyloquinoline alkaloid biosynthesis in angiosperms. *Phytochemistry*, 66(11): 1374–1393. doi:10.1016/j.phytochem.2005.04.029.
- Liscombe, D.K., Ziegler, J., Schmidt, J., Ammer, C., and Facchini, P.J. 2009. Targeted metabolite and transcript profiling for elucidating enzyme function: isolation of novel N-methyltransferases from three benzyloquinoline alkaloid-producing species. *Plant J.* 60(4): 729–743. doi:10.1111/j.1365-313X.2009.03980.x.
- Liscombe, D.K., Kamiyoshihara, Y., Ghironzi, J., Kempthorne, C.J., Hooton, K., Bulot, B., et al. 2022. A flavin-dependent monooxygenase produces nitrogenous tomato aroma volatiles using cysteine as a nitrogen source. *Proc. Natl. Acad. Sci.* 119(7). doi:10.1073/pnas.2118676119.
- Liu, Y., Ma, D., and Constabel, C.P. 2023. CRISPR/Cas9 disruption of MYB134 and MYB115 in transgenic poplar leads to differential reduction of proanthocyanidin synthesis in roots and leaves. *Plant Cell Physiol.* 64(10): 1189–1203. doi:10.1093/pcp/pcad086.
- Liu, Y., Zhu, P., Cai, S., Haughn, G., and Page, J.E. 2021. Three novel transcription factors involved in cannabinoid biosynthesis in *Cannabis sativa* L. *Plant Mol. Biol.* 106(1-2): 49–65. doi:10.1007/s11103-021-01129-9.
- Liyana, N.S., Awwad, F., Goncalves Dos Santos, K.C., Jayawardena, T.U., Merindol, N., and Desgagne-Penix, I. 2024. Navigating Amaryllidaceae alkaloids: bridging gaps and charting biosynthetic territories—a comprehensive review. *J. Exp. Bot.* 76. doi:10.1093/jxb/erae187.
- Long, X., Fang, Y., Qin, Y., Yang, J., and Xiao, X. 2021. Latex-specific transcriptome analysis reveals mechanisms for latex metabolism and natural rubber biosynthesis in laticifers of *Hevea brasiliensis*. *Ind. Crops Prod.* 171: 113835. doi:10.1016/j.indcrop.2021.113835.
- Luk, L.Y., Bunn, S., Liscombe, D.K., Facchini, P.J., and Tanner, M.E. 2007. Mechanistic studies on norcoclaurine synthase of benzyloquinoline

- line alkaloid biosynthesis: an enzymatic Pictet-Spengler reaction. *Biochemistry*, 46(35): 10153–10161. doi:[10.1021/bi700752n](https://doi.org/10.1021/bi700752n).
- Luo, X., Reiter, M.A., d'Espaux, L., Wong, J., Denby, C.M., Lechner, A., et al. 2019. Complete biosynthesis of cannabinoids and their unnatural analogues in yeast. *Nature*, 567(7746): 123–126. doi:[10.1038/s41586-019-0978-9](https://doi.org/10.1038/s41586-019-0978-9).
- Lussier, F.X., Colatrigiano, D., Wiltshire, Z., Page, J.E., and Martin, V.J. 2012. Engineering microbes for plant polyketide biosynthesis. *Comput. Struct. Biotechnol. J.* 3: e201210020. doi:[10.5936/csbj.201210020](https://doi.org/10.5936/csbj.201210020).
- Ma, C., Feng, Y., Zhou, S., Zhang, J., Guo, B., Xiong, Y., et al. 2023. Metabolomics and transcriptomics provide insights into the molecular mechanisms of anthocyanin accumulation in the seed coat of differently colored mung bean (*Vigna radiata* L.). *Plant Physiol. Biochem.* 200: 107739. doi:[10.1016/j.plaphy.2023.107739](https://doi.org/10.1016/j.plaphy.2023.107739).
- Ma, D., and Constabel, C.P. 2021. Complex regulation of proanthocyanidin biosynthesis in plants by R2R3 MYB activators and repressors. In *Recent advances in polyphenol research*. Edited by J.D. Reed, V.A.P. de Freitas and S. Quideau. John Wiley & Sons, West Sussex, UK. pp. 207–225.
- Ma, D., Reichelt, M., Yoshida, K., Gershenzon, J., and Constabel, C.P. 2018. Two R2R3-MYB proteins are broad repressors of flavonoid and phenylpropanoid metabolism in poplar. *Plant J.* 96(5): 949–965. doi:[10.1111/tip.14081](https://doi.org/10.1111/tip.14081).
- Ma, D., Tang, H., Reichelt, M., Piirtola, E.M., Salminen, J.P., Gershenzon, J., and Constabel, C.P. 2021. Poplar MYB117 promotes anthocyanin synthesis and enhances flavonoid B-ring hydroxylation by up-regulating the flavonoid 3',5'-hydroxylase gene. *J. Exp. Bot.* 72(10): 3864–3880. doi:[10.1093/jxb/erab116](https://doi.org/10.1093/jxb/erab116).
- Ma, J., Gu, Y., and Xu, P. 2022. Biosynthesis of cannabinoid precursor olivetolic acid in genetically engineered *Yarrowia lipolytica*. *Commun. Biol.* 5(1): 1239. doi:[10.1038/s42003-022-04202-1](https://doi.org/10.1038/s42003-022-04202-1).
- Ma, R., Huang, W., Hu, Q., Tian, G., An, J., Fang, T., et al. 2024. Tandemly duplicated MYB genes are functionally diverged in the regulation of anthocyanin biosynthesis in soybean. *Plant Physiol.* 194(4): 2549–2563. doi:[10.1093/plphys/kiad019](https://doi.org/10.1093/plphys/kiad019).
- Majhi, B.B., Gelinas, S.E., Merindol, N., Ricard, S., and Desgagne-Penix, I. 2023. Characterization of norbelladine synthase and noroxomaritidine/norcratichedine reductase reveals a novel catalytic route for the biosynthesis of Amaryllidaceae alkaloids including the Alzheimer's drug galanthamine. *Front. Plant Sci.* 14: 1231809. doi:[10.3389/fpls.2023.1231809](https://doi.org/10.3389/fpls.2023.1231809).
- Malli, R.P.N., Adal, A.M., Sarker, L.S., Liang, P., and Mahmoud, S.S. 2019. *De novo* sequencing of the *Lavandula angustifolia* genome reveals highly duplicated and optimized features for essential oil production. *Planta*, 249(1): 251–256. doi:[10.1007/s00425-018-3012-9](https://doi.org/10.1007/s00425-018-3012-9).
- Martin, D.M., and Bohlmann, J. 2004. Identification of *Vitis vinifera* (-)- α -terpineol synthase by in silico screening of full-length cDNA ESTs and functional characterization of recombinant terpene synthase. *Phytochemistry*, 65(9): 1223–1229. doi:[10.1016/j.phytochem.2004.03.018](https://doi.org/10.1016/j.phytochem.2004.03.018).
- Martin, D.M., Faldt, J., and Bohlmann, J. 2004. Functional characterization of nine Norway Spruce TPS genes and evolution of gymnosperm terpene synthases of the TPS-d subfamily. *Plant Physiol.* 135(4): 1908–1927. doi:[10.1104/pp.104.042028](https://doi.org/10.1104/pp.104.042028).
- Martin, D.M., Chiang, A., Lund, S.T., and Bohlmann, J. 2012. Biosynthesis of wine aroma: transcript profiles of hydroxymethylbutenyl diphosphate reductase, geranyl diphosphate synthase, and linalool/nerolidol synthase parallel monoterpenol glycoside accumulation in Gewürztraminer grapes. *Planta*, 236(3): 919–929. doi:[10.1007/s00425-012-1704-0](https://doi.org/10.1007/s00425-012-1704-0).
- Martin, D.M., Toub, O., Chiang, A., Lo, B.C., Ohse, S., Lund, S.T., and Bohlmann, J. 2009. The bouquet of grapevine (*Vitis vinifera* L. cv. Cabernet Sauvignon) flowers arises from the biosynthesis of sesquiterpene volatiles in pollen grains. *Proc. Natl. Acad. Sci.* 106(17): 7245–7250. doi:[10.1073/pnas.0901387106](https://doi.org/10.1073/pnas.0901387106).
- Martin, D.M., Aubourg, S., Schouwey, M.B., Daviet, L., Schalk, M., Toub, O., et al. 2010. Functional annotation, genome organization and phylogeny of the grapevine (*Vitis vinifera*) terpene synthase gene family based on genome assembly, FcDNA cloning, and enzyme assays. *BMC Plant Biol.* 10: 226. doi:[10.1186/1471-2229-10-226](https://doi.org/10.1186/1471-2229-10-226).
- Matchett-Oates, L., Spangenberg, G.C., and Cogan, N.O.I. 2021. Manipulation of cannabinoid biosynthesis via transient RNAi expression. *Front. Plant Sci.* 12: 773474. doi:[10.3389/fpls.2021.773474](https://doi.org/10.3389/fpls.2021.773474).
- McLellan, T.M., Caldwell, J.A., and Lieberman, H.R. 2016. A review of caffeine's effects on cognitive, physical and occupational performance. *Neurosci. Biobehav. Rev.* 71: 312. doi:[10.1016/j.neubiorev.2016.09.001](https://doi.org/10.1016/j.neubiorev.2016.09.001).
- Mellway, R.D., Tran, L.T., Prouse, M.B., Campbell, M.M., and Constabel, C.P. 2009. The wound-, pathogen-, and ultraviolet B-responsive MYB134 gene encodes an R2R3 MYB transcription factor that regulates proanthocyanidin synthesis in poplar. *Plant Physiol.* 150(2): 924–941. doi:[10.1104/pp.109.139071](https://doi.org/10.1104/pp.109.139071).
- Menendez-Perdomo, I.M., and Facchini, P.J. 2018. Benzylisoquinoline alkaloids biosynthesis in sacred lotus. *Molecules*, 23(11). doi:[10.3390/molecules23112899](https://doi.org/10.3390/molecules23112899).
- Miller, B., Madilao, L.L., Ralph, S., and Bohlmann, J. 2005. Insect-induced conifer defense. White pine weevil and methyl jasmonate induce traumatic resinosis, de novo formed volatile emissions, and accumulation of terpenoid synthase and putative octadecanoid pathway transcripts in Sitka spruce. *Plant Physiol.* 137(1): 369–382. doi:[10.1104/pp.104.050187](https://doi.org/10.1104/pp.104.050187).
- Morris, J.S., Groves, R.A., Hagel, J.M., and Facchini, P.J. 2018. An N-methyltransferase from *Ephedra sinica* catalyzing the formation of ephedrine and pseudoephedrine enables microbial phenylalkylamine production. *J. Biol. Chem.* 293(35): 13364–13376. doi:[10.1074/jbc.RA118.004067](https://doi.org/10.1074/jbc.RA118.004067).
- Morris, J.S., Dastmalchi, M., Li, J., Chang, L., Chen, X., Hagel, J.M., and Facchini, P.J. 2016. Plug-and-play benzylisoquinoline alkaloid biosynthetic gene discovery in engineered yeast. *Methods Enzymol.* 575: 143–178. doi:[10.1016/bs.mie.2016.03.023](https://doi.org/10.1016/bs.mie.2016.03.023).
- Murata, J., and De Luca, V. 2005. Localization of tabersonine 16-hydroxylase and 16-OH tabersonine-16-O-methyltransferase to leaf epidermal cells defines them as a major site of precursor biosynthesis in the vindoline pathway in *Catharanthus roseus*. *Plant J.* 44(4): 581–594. doi:[10.1111/j.1365-313X.2005.02557.x](https://doi.org/10.1111/j.1365-313X.2005.02557.x).
- Naake, T., Maeda, H.A., Proost, S., Tohge, T., and Fernie, A.R. 2021. Kingdom-wide analysis of the evolution of the plant type III polyketide synthase superfamily. *Plant Physiol.* 185(3): 857–875. doi:[10.1093/plphys/kiab086](https://doi.org/10.1093/plphys/kiab086).
- Nagel, J., and Page, J.E. 2010. DNA sequences involved in prenylflavonoid biosynthesis in hops and their use in modifying prenylflavonoid production in plants. Edited by USPTO. National Research Council of Canada, USA.
- Nagel, J., Culley, L.K., Lu, Y., Liu, E., Matthews, P.D., Stevens, J.F., and Page, J.E. 2008. EST analysis of hop glandular trichomes identifies an O-methyltransferase that catalyzes the biosynthesis of xanthohumol. *Plant Cell*, 20(1): 186–200. doi:[10.1105/tpc.107.055178](https://doi.org/10.1105/tpc.107.055178).
- Nguyen, T.D., and Dang, T.T. 2021. Cytochrome P450 enzymes as key drivers of alkaloid chemical diversification in plants. *Front. Plant Sci.* 12: 682181. doi:[10.3389/fpls.2021.682181](https://doi.org/10.3389/fpls.2021.682181).
- Nguyen, T.D., and Dang, T.T. 2022. Old path, new frontier. *Nat. Chem. Biol.* 18(6): 582–583. doi:[10.1038/s41589-022-01045-z](https://doi.org/10.1038/s41589-022-01045-z).
- Nguyen, T.D., Kwon, M., Kim, S.U., Fischer, C., and Ro, D.K. 2019. Catalytic plasticity of germacrene A oxidase underlies sesquiterpene lactone diversification. *Plant Physiol.* 181(3): 945–960. doi:[10.1104/pp.19.00629](https://doi.org/10.1104/pp.19.00629).
- Nguyen, T.D., Faraldos, J.A., Vardakou, M., Salmon, M., O'Maille, P.E., and Ro, D.K. 2016. Discovery of germacrene A synthases in *Barnadesia spinosa*: the first committed step in sesquiterpene lactone biosynthesis in the basal member of the Asteraceae. *Biochem. Biophys. Res. Commun.* 479(4): 622–627. doi:[10.1016/j.bbrc.2016.09.165](https://doi.org/10.1016/j.bbrc.2016.09.165).
- Nguyen, T.M., Grzech, D., Chung, K., Xia, Z., Nguyen, T.D., and Dang, T.T. 2023. Discovery of a cytochrome P450 enzyme catalyzing the formation of spirooxindole alkaloid scaffold. *Front. Plant Sci.* 14: 1125158. doi:[10.3389/fpls.2023.1125158](https://doi.org/10.3389/fpls.2023.1125158).
- Nguyen, T.M., Nguyen, T.D., Leung, Y.Y., McConnachie, M., Sannikov, O., Xia, Z., and Dang, T.T. 2021. Discovering and harnessing oxidative enzymes for chemoenzymatic synthesis and diversification of anticancer camptothecin analogues. *Commun. Chem.* 4(1): 177. doi:[10.1038/s42004-021-00602-2](https://doi.org/10.1038/s42004-021-00602-2).
- Niu, M., Fu, J., Ni, R., Xiong, R.L., Zhu, T.T., Lou, H.X., et al. 2021. Functional and structural investigation of chalcone synthases based on integrated metabolomics and transcriptome analysis on flavonoids and anthocyanins biosynthesis of the fern *Cydosorus parasiticus*. *Front. Plant Sci.* 12: 757516. doi:[10.3389/fpls.2021.757516](https://doi.org/10.3389/fpls.2021.757516).

- Nunes, H.H.C., Nguyen, T., and Dang, T.-T.T. 2022. Chemoenzymatic synthesis of natural products using plant biocatalysts. *Curr. Opin. Green Sustain. Chem.* 35. doi:10.1016/j.cogsc.2022.100627.
- O'Connor, S.E., and Maresh, J.J. 2006. Chemistry and biology of monoterpene indole alkaloid biosynthesis. *Nat. Prod. Rep.* 23(4): 547. doi:10.1039/b512615k.
- Okada, Y., Yamazaki, Y., Suh, D.Y., Sankawa, U., and Ito, K. 2001. Bi-functional activities of valerophenone synthase in hop (*Humulus lupulus* L.). *J. Am. Soc. Brew. Chem.* 59(4): 163–166. doi:10.1094/ASBCJ-59-0163.
- Page, J.E., and Stout, J.M. 2020. Cannabichromenic acid synthase from *Cannabis sativa*. Edited by USPTO. National Research Council of Canada, USA.
- Page, J.E., and Liu, E. 2023. Nucleic acid encoding N-methylputrescine oxidase and uses thereof. Edited by USPTO. 22nd Century Limited LLC, USA.
- Page, J.E., and Todd, A.T. 2023. Nucleic acid sequences encoding transcription factors regulating alkaloid biosynthesis and their use in modifying plant metabolism. Edited by USPTO. 22nd Century Limited LLC, USA.
- Page, J.E., and Gagne, S.J. 2024. Genes and proteins for aromatic polyketide synthesis. Edited by USPTO. University of Saskatchewan, National Research Council of Canada, USA.
- Page, J.E., Liu, E., and Nagel, J. 2013. Aromatic prenyltransferase from hop. Edited by USPTO. National Research Council of Canada, USA.
- Page, J.E., Baute, G., Mackenzie, J., Williams, K., and Gilchrist, E. 2020. Grafted cannabis plants and related methods for producing grafted cannabis plants. Edited by CIPO. Anandia Laboratories Inc, Canada.
- Pan, Q., Mustafa, N.R., Tang, K., Choi, Y.H., and Verpoorte, R. 2015. Monoterpenoid indole alkaloids biosynthesis and its regulation in *Catharanthus roseus*: a literature review from genes to metabolites. *Phytochem. Rev.* 15: 250. doi:10.1007/s11101-015-9406-4.
- Perez de Souza, L., Scossa, F., Proost, S., Bitocchi, E., Papa, R., Tohge, T., and Fernie, A.R. 2019. Multi-tissue integration of transcriptomic and specialized metabolite profiling provides tools for assessing the common bean (*Phaseolus vulgaris*) metabolome. *Plant J.* 97(6): 1132–1153. doi:10.1111/tpj.14178.
- Peters, D.J., and Constabel, C.P. 2002. Molecular analysis of herbivore-induced condensed tannin synthesis: cloning and expression of dihydroflavonol reductase from trembling aspen (*Populus tremuloides*). *Plant J.* 32(5): 701–712. doi:10.1046/j.1365-3113x.2002.01458.x.
- Pichersky, E., and Raguso, R.A. 2018. Why do plants produce so many terpenoid compounds? *New Phytol.* 220(3): 692–702. doi:10.1111/nph.14178.
- Pourcel, L., Routaboul, J.M., Cheynier, V., Lepiniec, L., and Debeaujon, I. 2007. Flavonoid oxidation in plants: from biochemical properties to physiological functions. *Trends Plant Sci.* 12(1): 29–36. doi:10.1016/j.tplants.2006.11.006.
- Power, R., Kurz, W.G., and De Luca, V. 1990. Purification and characterization of acetylcoenzyme A:deacetylindoline 4-O-acetyltransferase from *Catharanthus roseus*. *Arch. Biochem. Biophys.* 279(2): 376. doi:10.1016/0003-9861(90)90504-r.
- Pratyusha, D.S., and Sarada, D.V.L. 2022. MYB transcription factors-master regulators of phenylpropanoid biosynthesis and diverse developmental and stress responses. *Plant Cell Rep.* 41(12): 2245–2260. doi:10.1007/s00299-022-02927-1.
- Pyle, B.W., Tran, H.T., Pickel, B., Haslam, T.M., Gao, Z., MacNevin, G., et al. 2012. Enzymatic synthesis of valerena-4,7(11)-diene by a unique sesquiterpene synthase from the valerian plant (*Valeriana officinalis*). *FEBS J.* 279(17): 3136–3146. doi:10.1111/j.1742-4658.2012.08692.x.
- Pyne, M.E., Gold, N.D., and Martin, V.J. 2023a. Pathway elucidation and microbial synthesis of proaporphine and bis-benzylisoquinoline alkaloids from sacred lotus (*Nelumbo nucifera*). *Metab. Eng.* 77: 162–173. doi:10.1016/j.ymben.2023.03.010.
- Pyne, M.E., Bagley, J.A., Narcross, L., Kevvai, K., Exley, K., Davies, M., et al. 2023b. Screening non-conventional yeasts for acid tolerance and engineering *Pichia occidentalis* for production of muconic acid. *Nat. Comm.* 14(1): 5294. doi:10.1038/s41467-023-41064-5.
- Pyne, M.E., Kevvai, K., Grewal, P.S., Narcross, L., Choi, B., Bourgeois, L., et al. 2020. A yeast platform for high-level synthesis of tetrahydroisoquinoline alkaloids. *Nat. Commun.* 11(1):-. doi:10.1038/s41467-020-17172-x.
- Qu, Y., Safonova, O., and De Luca, V. 2019. Completion of the canonical pathway for assembly of anticancer drugs vincristine/vinblastine in *Catharanthus roseus*. *Plant J.* 97(2): 257–266. doi:10.1111/tpj.14111.
- Qu, Y., Easson, M.L., Froese, J., Simionescu, R., Hudlicky, T., and De Luca, V. 2015a. Completion of the seven-step pathway from tabersonine to the anticancer drug precursor vindoline and its assembly in yeast. *Proc. Natl. Acad. Sci.* 112(19): 6224–6229. doi:10.1073/pnas.1501821112.
- Qu, Y., Chakrabarty, R., Tran, H.T., Kwon, E.J., Kwon, M., Nguyen, T.D., and Ro, D.K. 2015b. A lettuce (*Lactuca sativa*) homolog of human Nogo-B receptor interacts with *cis*-prenyltransferase and is necessary for natural rubber biosynthesis. *J. Biol. Chem.* 290(4): 1898–1914. doi:10.1074/jbc.M114.616920.
- Qu, Y., Easson, M., Simionescu, R., Hajicek, J., Thamm, A.M.K., Salim, V., and De Luca, V. 2018a. Solution of the multistep pathway for assembly of corynanthean, strychnos, iboga, and aspidosperma monoterpenoid indole alkaloids from 19E-geissoschizine. *Proc. Natl. Acad. Sci.* 115(12): 3180–3185. doi:10.1073/pnas.1719979115.
- Qu, Y., Thamm, A.M.K., Czerwinski, M., Masada, S., Kim, K.H., Jones, G., et al. 2018b. Geissoschizine synthase controls flux in the formation of monoterpenoid indole alkaloids in a *Catharanthus roseus* mutant. *Planta*, 247(3): 625–634. doi:10.1007/s00425-017-2812-7.
- Rea, K.A., Casaretto, J.A., Al-Abdul-Wahid, M.S., Sukumaran, A., Geddes-McAlister, J., Rothstein, S.J., and Akhtar, T.A. 2019. Biosynthesis of cannflavins A and B from *Cannabis sativa* L. *Phytochemistry*, 164: 162–171. doi:10.1016/j.phytochem.2019.05.009.
- Ro, D.K., and Bohlmann, J. 2006. Diterpene resin acid biosynthesis in loblolly pine (*Pinus taeda*): functional characterization of abietadiene/levopimaradiene synthase (PtTPS-LAS) cDNA and subcellular targeting of PtTPS-LAS and abietadienol/abietadienol oxidase (PtAO, CYP720B1). *Phytochemistry*, 67(15): 1572–1578. doi:10.1016/j.phytochem.2006.01.011.
- Roach, C.R., Hall, D.E., Zerbe, P., and Bohlmann, J. 2014. Plasticity and evolution of (+)-3-carene synthase and (–)-sabinene synthase functions of a Sitka spruce monoterpene synthase gene family associated with weevil resistance. *J. Biol. Chem.* 289(34): 23859–23869. doi:10.1074/jbc.M114.571703.
- Roepeke, J., and Bozzo, G.G. 2015. *Arabidopsis thaliana* β -glucosidase BGLU15 attacks flavonol 3-O- β -glucoside-7-O- α -rhamnosides. *Phytochemistry*, 109: 24. doi:10.1016/j.phytochem.2014.10.028.
- Roth, J.H., and Ward, V.C. 2023. Production of astaxanthin using CBFD1/HFBD1 from *Adonis aestivalis* and the isopentenol utilization pathway in *Escherichia coli*. *Bioeng.* 10(9). doi:10.3390/bioengineering10091033.
- Roushan, M.R., Chen, C., Ahmadi, P., and Ward, V.C. 2023. *Agrobacterium tumefaciens*-mediated genetic engineering of green microalgae, *Chlorella vulgaris*. *J. Vis. Exp.* 200. doi:10.3791/65382.
- Rousset, A., Amor, A., Punvichai, T., Perino, S., Palu, S., Dorget, M., et al. 2021. Guayule (*Parthenium argentatum* A. Gray), a renewable resource for natural polyisoprene and resin: composition, processes and applications. *Molecules*, 26(3). doi:10.3390/molecules26030664.
- Salim, V., Jarecki, S.A., Vick, M., and Miller, R. 2023. Advances in metabolic engineering of plant monoterpene indole alkaloids. *Biology*, 12(8). doi:10.3390/biology12081056.
- Salim, V., Wiens, B., Masada-Atsumi, S., Yu, F., and De Luca, V. 2014. 7-deoxyloganic acid synthase catalyzes a key 3 step oxidation to form 7-deoxyloganic acid in *Catharanthus roseus* iridoid biosynthesis. *Phytochemistry*, 101: 23–31. doi:10.1016/j.phytochem.2014.02.009.
- Samanani, N., and Facchini, P.J. 2001. Isolation and partial characterization of norcoclaurine synthase, the first committed step in benzyloisoquinoline alkaloid biosynthesis, from opium poppy. *Planta*, 213(6): 898–906. doi:10.1007/s004250100581.
- Samanani, N., and Facchini, P.J. 2002. Purification and characterization of norcoclaurine synthase. The first committed enzyme in benzyloisoquinoline alkaloid biosynthesis in plants. *J. Biol. Chem.* 277(37): 33878–33883. doi:10.1074/jbc.M203051200.
- Samanani, N., Liscombe, D.K., and Facchini, P.J. 2004. Molecular cloning and characterization of norcoclaurine synthase, an enzyme catalyzing the first committed step in benzyloisoquinoline alkaloid biosynthesis. *Plant J.* 40(2): 302–313. doi:10.1111/j.1365-3113.2004.02210.x.
- Samanani, N., Park, S.U., and Facchini, P.J. 2005. Cell type-specific localization of transcripts encoding nine consecutive enzymes involved

- in protoberberine alkaloid biosynthesis. *Plant Cell*, 17(3): 915–926. doi:10.1105/tpc.104.028654.
- Sarker, L.S., and Mahmoud, S.S. 2015. Cloning and functional characterization of two monoterpene acetyltransferases from glandular trichomes of *L. x intermedia*. *Planta*, 242(3): 709–719. doi:10.1007/s00425-015-2325-1.
- Sarker, L.S., Demissie, Z.A., and Mahmoud, S.S. 2013. Cloning of a sesquiterpene synthase from *lavandula x intermedia* glandular trichomes. *Planta*, 238(5): 983–989. doi:10.1007/s00425-013-1937-6.
- Sarker, L.S., Adal, A.M., and Mahmoud, S.S. 2019. Diverse transcription factors control monoterpene synthase expression in lavender (*Lavandula*). *Planta*, 251(1): 5. doi:10.1007/s00425-019-03298-w.
- Sarker, L.S., Galata, M., Demissie, Z.A., and Mahmoud, S.S. 2012. Molecular cloning and functional characterization of borneol dehydrogenase from the glandular trichomes of *Lavandula x intermedia*. *Arch. Biochem. Biophys.* 528(2): 163–170. doi:10.1016/j.abb.2012.09.013.
- Sawler, J., Stout, J.M., Gardner, K.M., Hudson, D., Vidmar, J., Butler, L., et al. 2015. The genetic structure of marijuana and hemp. *PLoS One*, 10(8): e0133292. doi:10.1371/journal.pone.0133292.
- Scott, I.M., McDowell, T., Renaud, J.B., Krolikowski, S.W., Chen, L., and Dhaubhadel, S. 2021. Soybean (*Glycine max* L. Merr) host-plant defenses and resistance to the two-spotted spider mite (*Tetranychus urticae* Koch). *PLoS One*, 16(10): e0258198. doi:10.1371/journal.pone.0258198.
- Scott, I.M., McDowell, T., Renaud, J.B., Krolikowski, S.W., Chen, L., and Dhaubhadel, S. 2022. Investigation of metabolic resistance to soybean aphid (*Aphis glycines matsumura*) feeding in soybean cultivars. *Insects*, 13(4). doi:10.3390/insects13040356.
- Shahsavarani, M., Utomo, J.C., Kumar, R., Paz-Galeano, M., Garza-García, J.J.O., Mai, Z., et al. 2023. Improved protein glycosylation enabled heterologous biosynthesis of monoterpenoid indole alkaloids and their unnatural derivatives in yeast. *Metab. Eng. Commun.* 16. doi:10.1016/j.mec.2022.e00215.
- Sharma, S.K., Vij, A.S., and Sharma, M. 2013. Mechanisms and clinical uses of capsaicin. *Eur. J. Pharmacol.* 720(1-3): 55–62. doi:10.1016/j.ejphar.2013.10.053.
- Shiba, Y., Paradise, E.M., Kirby, J., Ro, D.-K., and Keasling, J.D. 2007. Engineering of the pyruvate dehydrogenase bypass in *Saccharomyces cerevisiae* for high-level production of isoprenoids. *Metab. Eng.* 9(2): 160–168. doi:10.1016/j.ymben.2006.10.005.
- Singh, A., and Desgagne-Penix, I. 2017. Transcriptome and metabolome profiling of *Narcissus pseudonarcissus* 'King Alfred' reveal components of Amaryllidaceae alkaloid metabolism. *Scientific Rep.* 7(1): 17356. doi:10.1038/s41598-017-17724-0.
- Singh, A., Menéndez-Perdomo, I.M., and Facchini, P. 2019. Benzylisoquinoline alkaloid biosynthesis in opium poppy: an update. *Phytochem. Rev.* 18: 1482–1457. doi:10.1007/s11101-019-09644-w.
- Singh, A., Massicotte, M.A., Garand, A., Tousignant, L., Ouellette, V., Berube, G., and Desgagne-Penix, I. 2018. Cloning and characterization of norbelladine synthase catalyzing the first committed reaction in Amaryllidaceae alkaloid biosynthesis. *BMC Plant Biol.* 18(1): 338. doi:10.1186/s12870-018-1570-4.
- Singh, B., and Sharma, R.A. 2015. Plant terpenes: defense responses, phylogenetic analysis, regulation and clinical applications. 3 *Biotech*, 5(2): 129–151. doi:10.1007/s13205-014-0220-2.
- Slattery, S.S., Diamond, A., Wang, H., Therrien, J.A., Lant, J.T., Jazey, T., et al. 2018. An expanded plasmid-based genetic toolbox enables Cas9 genome editing and stable maintenance of synthetic pathways in *Phaeodactylum tricornutum*. *ACS Synth. Biol.* 7(2): 328–338. doi:10.1021/acssynbio.7b00191.
- St-Pierre, B., and De Luca, V. 1995. A cytochrome P-450 monooxygenase catalyzes the first step in the conversion of tabersonine to vindoline in *Catharanthus roseus*. *Plant Physiol.* 109(1): 131–139. doi:10.1104/pp.109.1.131.
- St-Pierre, B., Besseau, S., Clastre, M., Courdavault, V., Courtois, M., Crèche, J., et al. 2013. Deciphering the evolution, cell biology and regulation of monoterpene indole alkaloids. In *New light on alkaloid biosynthesis and future prospects*. Edited by N. Giglioli-Guivarc'h. Academic Press, Kidlington, United Kingdom. pp. 73–109.
- Stamp, L.K., Horsley, G., Te Karu, L., Dalbeth, N., and Barclay, M. 2024. Colchicine: the good, the bad, the ugly and how to minimize the risks. *Rheumatology*, 63(4): 944. doi:10.1093/rheumatology/kead625.
- Stavrinides, A.K., Tatsis, E.C., Dang, T.T., Caputi, L., Stevenson, C.E.M., Lawson, D.M., et al. 2018. Discovery of a short-chain dehydrogenase from *Catharanthus roseus* that produces a new monoterpene indole alkaloid. *Chembiochem*, 19(9): 940–948. doi:10.1002/cbic.201700621.
- Stephanopoulos, G., Edgar, S.M., Chatzivasilieuo, A.O., and Ward, V. 2019. Microbial engineering for the production of isoprenoids.
- Stout, J.M., Boubakir, Z., Ambrose, S.J., Purves, R.W., and Page, J.E. 2012. The hexanoyl-CoA precursor for cannabinoid biosynthesis is formed by an acyl-activating enzyme in *Cannabis sativa* trichomes. *Plant J.* 71(3): 353–365. doi:10.1111/j.1365-3113X.2012.04949.x.
- Suh, D.Y., Kagami, J., Fukuma, K., and Sankawa, U. 2000a. Evidence for catalytic cysteine-histidine dyad in chalcone synthase. *Biochem. Biophys. Res. Commun.* 275: 725–730. doi:10.1006/bbrc.2000.3368.
- Suh, D.Y., Fukuma, K., Kagami, J., Yamazaki, Y., Shibuya, M., Ebizuka, Y., and Sankawa, U. 2000b. Identification of amino acid residues important in the cyclization reactions of chalcone and stilbene synthases. *Biochem. J.* 350: 229–235. doi:10.1042/bj3500229.
- Suh, D.Y., Kagami, J., Fukuma, K., Iwanami, N., Yamazaki, Y., Yurimoto, H., et al. 2000c. Chalcone and stilbene synthases expressed in eucaryotes exhibit reduced cross-reactivity in vitro. *Chemical and Pharm. Bull.* 48(7): 1051–1054. doi:10.1248/cpb.48.1051.
- Sun, Y., Hegebarth, D., and Jetter, R. 2021. Acyl-CoA desaturase ADS4.2 is involved in the formation of characteristic wax alkenes in young *Arabidopsis* leaves. *Plant Physiol.* 186(4): 1812–1831. doi:10.1093/plphys/kiab182.
- Sun, Y., Ruiz Orduna, A., Zhang, Z., Feakins, S.J., and Jetter, R. 2023. Biosynthesis of barley wax β -diketones: a type-III polyketide synthase condensing two fatty acyl units. *Nature Commun.* 14(1): 7284. doi:10.1038/s41467-023-42917-9.
- Sun, Y., Yao, R., Ji, X., Wu, H., Luna, A., Wang, Z., and Jetter, R. 2020. Characterization of an alkylresorcinol synthase that forms phenolics accumulating in the cuticular wax on various organs of rye (*Secale cereale*). *Plant J.* 102(6): 1294–1312. doi:10.1111/tpj.14704.
- Sunil, L., and Shetty, N.P. 2022. Biosynthesis and regulation of anthocyanin pathway genes. *Appl. Microbiol. Biotechnol.* 106: 1783–1798. doi:10.1007/s00253-022-11835-z.
- Sunstrum, F.G., Liu, H.L., Jancsik, S., Madilao, L.L., Bohlmann, J., and Irmisch, S. 2021. 4-Coumaroyl-CoA ligases in the biosynthesis of the anti-diabetic metabolite montbretin A. *PLoS One*, 16(10): e0257478. doi:10.1371/journal.pone.0257478.
- Tapia, G., Gonzalez, M., Mendez, J., Schmida-Hirschmann, G., Arrey, O., Carrasco, B., et al. 2024. Transcriptome analysis reveals biosynthesis and regulation of flavonoid in common bean seeds during grain filling. *BMC Plant Biol.* 24(1): 916. doi:10.1186/s12870-024-05593-5.
- Tatsis, E.C., Carqueijeiro, I., Duge de Bernonville, T., Franke, J., Dang, T.T., Oudin, A., et al. 2017. A three enzyme system to generate the *Strychnos* alkaloid scaffold from a central biosynthetic intermediate. *Nature Commun.* 8(1): 316. doi:10.1038/s41467-017-00154-x.
- Tewksbury, J.J., and Nabhan, G.P. 2001. Seed dispersal. Directed deterrence by capsaicin in chilies. *Nature*, 412(6845): 403–404. doi:10.1038/35086653.
- Thomas, F., Schmidt, C., and Kayser, O. 2020. Bioengineering studies and pathway modeling of the heterologous biosynthesis of tetrahydrocannabinolic acid in yeast. *Appl. Microbiol. Biotechnol.* 104(22): 9551–9563. doi:10.1007/s00253-020-10798-3.
- Tian, Y., Li, Q., Rao, S., Wang, A., Zhang, H., Wang, L., et al. 2021. Metabolic profiling and gene expression analysis provides insights into flavonoid and anthocyanin metabolism in poplar. *Tree Physiol.* 41(6): 1046–1064. doi:10.1093/treephys/tpaa152.
- Todd, A.T., Liu, E., Polvi, S.L., Pammett, R.T., and Page, J.E. 2010. A functional genomics screen identifies diverse transcription factors that regulate alkaloid biosynthesis in *Nicotiana benthamiana*. *Plant J.* 62(4): 589–600. doi:10.1111/j.1365-3113X.2010.04186.x.
- Torres, M.A., Hoffarth, E., Eugenio, L., Savtchouk, J., Chen, X., Morris, J.S., et al. 2016. Structural and functional studies of pavin N-methyltransferase from *Thalictrum flavum* reveal novel insights into substrate recognition and catalytic mechanism. *J. Biol. Chem.* 291(45): 23403–23415. doi:10.1074/jbc.M116.747261.
- Tousignant, L., Diaz-Garza, A.M., Majhi, B.B., Gelinis, S.E., Singh, A., and Desgagne-Penix, I. 2022. Transcriptome analysis of *Leucojum aestivum* and identification of genes involved in norbelladine biosynthesis. *Planta*, 255(2): 30. doi:10.1007/s00425-021-03741-x.

- Trush, K., and Pal'ove-Balang, P. 2023. Biosynthesis and role of isoflavonoids in legumes under different environmental conditions. *Plant Stress*, 8. doi:10.1016/j.stress.2023.100153.
- Unterlander, N., Mats, L., McGary, L.C., Gordon, H.O.W., and Bozzo, G.G. 2022. Kaempferol rhamnoside catabolism in rosette leaves of senescing *Arabidopsis* and postharvest stored radish. *Planta*, 256(2): 36. doi:10.1007/s00425-022-03949-5.
- Utomo, J.C., Barrell, H.B., Kumar, R., Smith, J., Brant, M.S., De la Hoz Siegler, H., and Ro, D.-K. 2024. Reconstructing curcumin biosynthesis in yeast reveals the implication of caffeoyl-shikimate esterase in phenylpropanoid metabolic flux. *Metab. Eng.* 82: 286–296. doi:10.1016/j.ymben.2024.02.011.
- van Bakel, H., Stout, J.M., Cote, A.G., Tallon, C.M., Sharpe, A.G., Hughes, T.R., and Page, J.E. 2011. The draft genome and transcriptome of *Cannabis sativa*. *Genome Biol.* 12(10): R102. doi:10.1186/gb-2011-12-10-r102.
- Van Gelder, K., Forrester, T., and Akhtar, T.A. 2020. Evidence from stable-isotope labeling that catechol is an intermediate in salicylic acid catabolism in the flowers of *Silene latifolia* (white campion). *Planta*, 252(1): 3. doi:10.1007/s00425-020-03410-5.
- Van Gelder, K., Virta, L.K.A., Easlick, J., Prudhomme, N., McAlister, J.A., Geddes-McAlister, J., and Akhtar, T.A. 2021. A central role for polyprenol reductase in plant dolichol biosynthesis. *Plant Sci.* 303: 110773. doi:10.1016/j.plantsci.2020.110773.
- Van Gelder, K., Rea, K.A., Virta, L.K.A., Whitnell, K.L., Osborn, M., Vatta, M., et al. 2018. Medium-chain polyprenols influence chloroplast membrane dynamics in *Solanum lycopersicum*. *Plant Cell Physiol.* 59(11): 2350–2365. doi:10.1093/pcp/pcy157.
- Vazquez-Flota, F., De Carolis, E., Alarco, A.M., and De Luca, V. 1997. Molecular cloning and characterization of desacetoxyvindoline-4-hydroxylase, a 2-oxoglutarate dependent-dioxygenase involved in the biosynthesis of vindoline in *Catharanthus roseus* (L.) G. Don. *Plant Mol. Biol.* 34(6): 935–948. doi:10.1023/a:1005894001516.
- Veljanovski, V., and Constabel, C.P. 2013. Molecular cloning and biochemical characterization of two UDP-glycosyltransferases from poplar. *Phytochemistry*, 91: 148–157. doi:10.1016/j.phytochem.2012.12.012.
- Wagner, K.H., and Elmadfa, I. 2003. Biological relevance of terpenoids. Overview focusing on mono-, di- and tetraterpenes. *Ann. Nutr. Metab.* 47(3-4): 95–106. doi:10.1159/000070030.
- Wang, J., and De Luca, V. 2005. The biosynthesis and regulation of biosynthesis of Concord grape fruit esters, including 'foxy' methylanthranilate. *Plant J.* 44(4): 606–619. doi:10.1111/j.1365-313X.2005.02552.x.
- Wang, Z., Yeats, T., Han, H., and Jetter, R. 2010. Cloning and characterization of oxidosqualene cyclases from *Kalanchoe daigremontiana*: enzymes catalyzing up to 10 rearrangement steps yielding friedelin and other triterpenoids. *J. Biol. Chem.* 285(39): 29703–29712. doi:10.1074/jbc.M109.098871.
- Wang, Z., Guhling, O., Yao, R., Li, F., Yeats, T.H., Rose, J.K., and Jetter, R. 2011. Two oxidosqualene cyclases responsible for biosynthesis of tomato fruit cuticular triterpenoids. *Plant Physiol.* 155(1): 540–552. doi:10.1104/pp.110.162883.
- Watkins, J.L., Li, Q., Yeaman, S., and Facchini, P.J. 2023. Elucidation of the mescaline biosynthetic pathway in peyote (*Lophophora williamsii*). *Plant J.* 116(3): 635–649. doi:10.1111/tpj.16447.
- Weaver, B.A. 2014. How Taxol/paclitaxel kills cancer cells. *Mol. Biol. Cell.* 25(18): 2677–2681. doi:10.1091/mbc.E14-04-0916.
- Williams, D., Qu, Y., Simionescu, R., and De Luca, V. 2019. The assembly of (+)-vincadifformine and (-)-tabersonine-derived monoterpenoid indole alkaloids in *Catharanthus roseus* involves separate branch pathways. *Plant J.* 99(4): 626–636. doi:10.1111/tpj.14346.
- Xiao, M., Zhang, Y., Chen, X., Lee, E.J., Barber, C.J., Chakrabarty, R., et al. 2013. Transcriptome analysis based on next-generation sequencing of non-model plants producing specialized metabolites of biotechnological interest. *J. Biotechnol.* 166(3): 122–134. doi:10.1016/j.jbiotec.2013.04.004.
- Yadav, V.G., Roy, P., Ayakar, S., Chang, B., and Hsu, E. 2022. Compositions and methods for biosynthesis of terpenoids or cannabinoids in a heterologous system. Edited by USPTO. Inmed Pharmaceuticals Inc., USA.
- Yamazaki, Y., Suh, D.Y., Sitthithaworn, W., Ishiguro, K., Kobayashi, Y., Shibuya, M., et al. 2001. Diverse chalcone synthase superfamily enzymes from the most primitive vascular plant, *Psilotum nudum*. *Planta*, 214(1): 75–84. doi:10.1007/s004250100586.
- Yang, W., Chen, X., Li, Y., Guo, S., Wang, Z., and Yu, X. 2020. Advances in pharmacological activities of terpenoids. *Nat. Prod. Commun.* 15(3). doi:10.1177/1934578X20903555.
- Yi, J., Derynck, M.R., Li, X., Telmer, P., Marsolais, F., and Dhaubhadel, S. 2010. A single-repeat MYB transcription factor, *GmMYB176*, regulates CHS8 gene expression and affects isoflavonoid biosynthesis in soybean. *Plant J.* 62(6): 1019–1034. doi:10.1111/j.1365-313X.2010.04214.x.
- Yoshida, K., Ma, D., and Constabel, C.P. 2015. The MYB182 protein down-regulates proanthocyanidin and anthocyanin biosynthesis in poplar by repressing both structural and regulatory flavonoid genes. *Plant Physiol.* 167(3): 693–710. doi:10.1104/pp.114.253674.
- Yousefi-Taemeh, M., Lin, J., Ifa, D.R., Parrott, W., and Kovinich, N. 2021. Metabolomics differences of *glycine max* QTLs resistant to soybean looper. *Metabolites*, 11(10). doi:10.3390/metabo11100710.
- Yu, B., Lydiate, D.J., Schafer, U.A., and Hannoufa, A. 2007. Characterization of a β -carotene hydroxylase of *Adonis aestivalis* and its expression in *Arabidopsis thaliana*. *Planta*, 226(1): 181–192. doi:10.1007/s00425-006-0455-1.
- Yu, B., Lydiate, D.J., Young, L.W., Schafer, U.A., and Hannoufa, A. 2008. Enhancing the carotenoid content of *Brassica napus* seeds by down-regulating lycopene ϵ -cyclase. *Transgenic Res.* 17(4): 573–585. doi:10.1007/s11248-007-9131-x.
- Yu, M., and Facchini, P.J. 1999. Purification, characterization, and immunolocalization of hydroxycinnamoyl-CoA: tyramine N-(hydroxycinnamoyl)transferase from opium poppy. *Planta*, 209(1): 33–44. doi:10.1007/s004250050604.
- Zager, J.J., Lange, I., Srividya, N., Smith, A., and Lange, B.M. 2019. Gene networks underlying cannabinoid and terpenoid accumulation in cannabis. *Plant Physiol.* 180(4): 1877–1897. doi:10.1104/pp.18.01506.
- Zagoskina, N.V., Zubova, M.Y., Nechaeva, T.L., Kazantseva, V.V., Goncharuk, E.A., Katanskaya, V.M., et al. 2023. Polyphenols in plants: structure, biosynthesis, abiotic stress regulation, and practical applications (review). *Int. J. Mol. Sci.* 24(18). doi:10.3390/ijms241813874.
- Zerbe, P., Chiang, A., Yuen, M., Hamberger, B., Hamberger, B., Draper, J.A., et al. 2012. Bifunctional *cis*-abienol synthase from *Abies balsamea* discovered by transcriptome sequencing and its implications for diterpenoid fragrance production. *J. Biol. Chem.* 287(15): 12121–12131. doi:10.1074/jbc.M111.317669.
- Zhang, H., Zhang, W., Meng, S., Hui, L., Liu, X., Chen, W., et al. 2024. Construction of a high-density genetic map for yardlong bean and identification of ANT1 as a regulator of anthocyanin biosynthesis. *Hortic. Res.* 11(1): uhad247. doi:10.1093/hr/uhad247.
- Zhang, J., Yang, Y., Zheng, K., Xie, M., Feng, K., Jawdy, S.S., et al. 2018a. Genome-wide association studies and expression-based quantitative trait loci analyses reveal roles of HCT2 in caffeoylquinic acid biosynthesis and its regulation by defense-responsive transcription factors in *Populus*. *New Phytol.* 220(2): 516. doi:10.1111/nph.15297.
- Zhang, Y., Chen, S., Fu, Y., Yang, H., and Ye, X. 2018b. Proanthocyanidins from Chinese bayberry (*Myrica rubra* Sieb. et Zucc.) leaves: structure elucidation and bioactive functions. In *Advances in plant phenolics: from chemistry to human health*. Edited by G.K. Jayaprakasha, B.S. Patil and G. Gattuso. American Chemical Society, Washington DC, USA. pp. 185–199.
- Zhang, J., Hansen, L.G., Gudich, O., Viehriig, K., Lassen, L.M.M., Schrubbers, L., et al. 2022. A microbial supply chain for production of the anti-cancer drug vinblastine. *Nature*, 609(7926): 347. doi:10.1038/s41586-022-05157-3.
- Zhang, Q., Wang, S., Qin, B., Sun, H.-y., Yuan, X.-k., Wang, Q., et al. 2023. Analysis of the transcriptome and metabolome reveals phenylpropanoid mechanism in common bean (*Phaseolus vulgaris*) responding to salt stress at sprout stage. *Food Energy Secur.* 12(5). doi:10.1002/fes3.481.
- Zhou, F., and Pichersky, E. 2020. More is better: the diversity of terpene metabolism in plants. *Curr. Opin. Plant Biol.* 55: 1–10. doi:10.1016/j.pbi.2020.01.005.
- Zirpel, B., Degenhardt, F., Zammarelli, C., Wibberg, D., Kalinowski, J., Stehle, F., and Kayser, O. 2018. Optimization of Δ^9 -tetrahydrocannabinolic acid synthase production in *Komagataella phaffii* via post-translational bottleneck identification. *J. Biotechnol.* 272-273: 40–47. doi:10.1016/j.jbiotec.2018.03.008.

Zulak, K.G., Dullat, H.K., Keeling, C.I., Lippert, D., and Bohlmann, J. 2010. Immunofluorescence localization of levopimaradiene/abietadiene synthase in methyl jasmonate treated stems of Sitka spruce (*Picea sitchensis*) shows activation of diterpenoid biosynthesis in cortical and developing traumatic resin ducts. *Phytochemistry*, 71(14-15): 1695–1699. doi:[10.1016/j.phytochem.2010.07.002](https://doi.org/10.1016/j.phytochem.2010.07.002).

Zulak, K.G., Lippert, D.N., Kuzyk, M.A., Domanski, D., Chou, T., Borchers, C.H., and Bohlmann, J. 2009. Targeted proteomics using selected reaction monitoring reveals the induction of specific terpene synthases in a multi-level study of methyl jasmonate-treated Norway spruce (*Picea abies*). *Plant J.* 60(6): 1015–1030. doi:[10.1111/j.1365-313X.2009.04020.x](https://doi.org/10.1111/j.1365-313X.2009.04020.x).