



Canadian Journal of Pain

Revue canadienne de la douleur

ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/ucjp20>

Development of a National Pain Management Competency Profile to Guide Entry-level Physiotherapy Education in Canada

Nathan Augeard, Geoff Bostick, Jordan Miller, David Walton, Yannick Tousignant-Laflamme, Anne Hudon, André Bussi eres, Lynn Cooper, Nicol McNiven, Alik Thomas, Lesley Singer, Scott M. Fishman, Marie H. Bement, Julia M. Hush, Kathleen A. Sluka, Judy Watt-Watson, Lisa C. Carlesso, Sinead Dufour, Roland Fletcher, Katherine Harman, Judith Hunter, Suzy Ngomo, Neil Pearson, Kadija Perreault, Barbara Shay, Peter Stilwell, Susan Tupper & Timothy H. Wideman

To cite this article: Nathan Augeard, Geoff Bostick, Jordan Miller, David Walton, Yannick Tousignant-Laflamme, Anne Hudon, Andr  Bussi eres, Lynn Cooper, Nicol McNiven, Alik Thomas, Lesley Singer, Scott M. Fishman, Marie H. Bement, Julia M. Hush, Kathleen A. Sluka, Judy Watt-Watson, Lisa C. Carlesso, Sinead Dufour, Roland Fletcher, Katherine Harman, Judith Hunter, Suzy Ngomo, Neil Pearson, Kadija Perreault, Barbara Shay, Peter Stilwell, Susan Tupper & Timothy H. Wideman (2021): Development of a National Pain Management Competency Profile to Guide Entry-level Physiotherapy Education in Canada, Canadian Journal of Pain, DOI: [10.1080/24740527.2021.2004103](https://doi.org/10.1080/24740527.2021.2004103)

To link to this article: <https://doi.org/10.1080/24740527.2021.2004103>



  2021 The Author(s). Published with license by Taylor & Francis Group, LLC.



Accepted author version posted online: 17 Nov 2021.



Submit your article to this journal [↗](#)



Article views: 506



View related articles [↗](#)



View Crossmark data [↗](#)

Publisher: Taylor & Francis & The Author(s). Published with license by Taylor & Francis Group, LLC.

Journal: *Canadian Journal of Pain*

DOI: 10.1080/24740527.2021.2004103

Development of a National Pain Management Competency Profile to Guide Entry-level Physiotherapy Education in Canada.

Nathan Augeard¹, MSc, PT; Geoff Bostick², PhD, PT; Jordan Miller³, PhD, PT; David Walton⁴, PhD, PT; Yannick Tousignant-Laflamme⁵, PhD, PT; Anne Hudon⁶, PhD, PT; André Bussièrès¹, PhD, DC; Lynn Cooper⁷, BES; Nicol McNiven⁸, MSc, PT; Aliko Thomas¹, PhD, OT; Lesley Singer¹, MSc, PT; Scott M. Fishman⁹, MD; Marie H. Bement¹⁰, PhD, PT; Julia M. Hush¹¹, PhD, PT; Kathleen A. Sluka¹², PhD, PT; Judy Watt-Watson¹³, PhD, RN; Lisa C. Carlesso¹⁴, PhD, PT; Sinead Dufour¹⁴, PhD, PT; Roland Fletcher¹⁵, MSc, PT; Katherine Harman¹⁶, PhD, PT; Judith Hunter¹⁷, PhD, PT; Suzy Ngomo¹⁸, PhD, MD; Neil Pearson¹⁵, MSc, PT; Kadija Perreault¹⁹, PhD, PT; Barbara Shay²⁰, PhD, PT; Peter Stilwell¹, PhD, DC; Susan Tupper²¹, PhD, PT; Timothy H. Wideman¹, PhD, PT.

1. School of Physical and Occupational Therapy, McGill University, Montreal, Canada
2. Department of Physical Therapy, University of Alberta, Edmonton, Canada
3. School of Rehabilitation Therapy, Queen's University, Kingston, Canada
4. School of Physical Therapy, Western University, London, Canada
5. School of Rehabilitation, Université de Sherbrooke, Sherbrooke, Canada
6. School of Rehabilitation, Université de Montréal, Montreal, Canada
7. Canadian Injured Workers Alliance, Thunder Bay, Canada
8. Faculty of Health Sciences, University of Ottawa, Ottawa, Canada

9. Davis School of Medicine, University of California, Sacramento, US
10. Department of Physical Therapy, Marquette University, Milwaukee, US
11. Department of Health Professions, Macquarie University, Sydney, Australia
12. Department of Physical Therapy and Rehabilitation Science, University of Iowa, Iowa City, US
13. Lawrence S Bloomberg Faculty of Nursing, University of Toronto, Toronto, Canada
14. School of Rehabilitation Science, Faculty of Health Sciences, McMaster University, Hamilton, Canada
15. Department of Physical Therapy, University of British Columbia, Vancouver, Canada
16. School of Physiotherapy, Dalhousie University, Halifax, Canada
17. Department of Physical Therapy, University of Toronto, Toronto, Canada
18. Department of Health Science, Université du Québec à Chicoutimi, Chicoutimi, Canada
19. Department of Rehabilitation Science, Faculty of Medicine, Université Laval, Québec City, Canada
20. Department of Physical Therapy, University of Manitoba, Winnipeg, Canada
21. School of Rehabilitation Sciences, University of Saskatchewan, Saskatoon, Canada

Corresponding author: Dr. Timothy Wideman, School of Physical and Occupational Therapy, Faculty of Medicine, McGill University, 3654 Promenade Sir-William-Osler, Montreal, QC H3G 1Y5; timothy.wideman@mcgill.ca.

Funding: This project was funded by the Canadian Institutes of Health Research Project Grant and was supported by the Centre de Recherche Interdisciplinaire en Réadaptation du Montréal Métropolitain.

Conflict of interest:

The nature of participatory research is that stakeholders are participants and are expected to have investments in the area researched; this investment is part of the research process. Consistent with this approach to research, the majority of the authors are university lecturers or professors focusing on pain education and receive a salary for this work. In addition, the following authors report financial compensation and/or royalties related to their work in pain education outside of their academic positions: DW, YTL, LS, MHB, RF, NP and TW.

Disclosure statement: NA, GB, JM, AH, AB, LC, NM, AT, SMF, JH, KS, JWW, LC, SD, KH, JH, SN, KP, BS, PS and ST report no potential competing interests.

Informed Consent Statement: informed written consent to participate in this project was obtained prior to the start of this study.

Running title: Developing a Pain Management Competency Profile.

Word count: 4327 (excluding abstract)

Abstract

Background: National strategies from North America call for substantive improvements in entry-level pain management education to help reduce the burden of chronic pain. Past work has generated a valuable set of interprofessional pain management competencies to guide the education of future health professionals. However, there has been very limited work that has explored the development of such competencies for individual professions in different regions. Developing profession-specific competencies tailored to the local context is a necessary first step to integrate them within local regulatory systems. Our group is working toward this goal within the context of entry-level physiotherapy (PT) programs across Canada. **Aims:** This study aimed to create a consensus-based competency profile for pain management, specific to the Canadian

PT context. **Methods:** A modified Delphi was used to achieve consensus across Canadian university-based and clinical pain educators. **Results:** Representatives from 14 entry-level PT programs (93% of Canadian programs) and six clinical educators were recruited. After two rounds, a total of 15 competencies reached the pre-determined endorsement threshold (75%). Most participants (85%) reported being “very satisfied” with the process. **Conclusions:** This process achieved consensus on a novel pain management competency profile specific to the Canadian PT context. The resulting profile delineates the necessary abilities required by physiotherapists to manage pain upon entry-to-practice. Participants were very satisfied with the process. This study also contributes to the emerging literature on integrated research in pain management by profiling research methodology that can be used to inform related work in other health professions and regions.

Key words: pain management; competency profile; participatory research; knowledge translation; physiotherapy.

Introduction

Chronic pain is a major healthcare issue associated with significant burden in terms of personal suffering and low quality of life for millions of North Americans^{1,2}. The lack of comprehensive pain management education across entry-level healthcare programs has been recognized as a major barrier to alleviating the burden of chronic pain². Previous work has shown that there are widespread discrepancies in how healthcare providers are trained to manage pain, both within and across healthcare professions³⁻⁵. These inconsistencies likely contribute to healthcare providers feeling ill-prepared in caring for their patients suffering from pain⁶⁻⁸, as well as people living with pain feeling misunderstood by their healthcare providers⁸⁻¹¹.

National strategies from the US and Canada emphasize the central importance of entry-level education to address the identified inconsistencies in pain management education^{1,2}. These national strategies call for the development of pain management competencies to guide the education of future healthcare professionals^{1,2}. The integration of key stakeholders in this process, such as people living with pain, healthcare students and pain management educators, is essential to develop tailored tools that are relevant to the people that will be using them¹²⁻¹⁵. To date, there has been limited work on entry-level healthcare education about pain management that focuses on developing resources tailored to specific contexts (e.g., across different regions and/or professions) and that uses robust participatory research methods. For example, seminal work from Fishman and colleagues¹⁶ generated a valuable set of interprofessional pain management competencies; however, research is yet to explore the development of corresponding competencies for individual healthcare professions and include end-users (e.g., people living with pain, educators, or students) in the creation process. Past work has highlighted the need for profession-specific competencies to match the unique attributes of each profession^{17,18}, as well as the importance of including stakeholders in the research process to increase relevance of the output and facilitate its uptake by end users¹³. In Canada, many healthcare professions are governed at the provincial level, with their own regulatory systems and priorities. Developing profession-specific competencies that are tailored to the local context is a necessary first step in further integrating pain management competencies within local regulatory systems.

Our group has recently started working toward the goal of establishing nationally accepted competencies for pain assessment and management within the context of entry-level physiotherapy (PT) programs across Canada^{18,19}. Physiotherapists (PTs) play an essential role in

in the management of pain, through the use of biopsychosocial interventions to increase patient knowledge and self-management skills, to reduce pain and disability, as well as to empower individuals to manage and live well with pain. Work from our group has clearly highlighted the need for tailored competencies to guide the improvement of entry-level pain education for PTs across Canada¹⁸. The current lack of a national pain competency profile for PTs likely contributes to the major discrepancies observed in how pain management competencies are integrated across Canadian PT education programs¹⁹.

The primary aim of this study was to achieve consensus across stakeholders on a new competency profile for pain management education, specific to the entry-level Canadian PT context.

Methods

Design

A modified Delphi design was used to achieve consensus across relevant stakeholders on a competency profile for pain management education in Canadian PT programs. The modified Delphi is an iterative process that uses a systematic progression of repeated rounds of voting to achieve agreement²²⁻²⁴. This approach strategically uses existing literature, perspectives of stakeholders and the judgement of experts within a field to reach consensus^{22,25}. Ethical approval for this study was granted by the McGill University Institutional Review Board.

Scope of the competency profile and conceptual framework

Throughout the generation and the consensus-building process, the scope of the profile was anchored by the definition of a competency as “an observable ability of a health professional, integrating multiple components such as knowledge, skills, and attitudes”^{26(p641)}. This profile

aimed to offer a profile of PT-specific core competencies that delineate the necessary abilities required to manage pain upon entry-to-practice¹⁸. In addition to distinct competencies, key concepts (e.g., therapeutic alliance) were defined and included in the profile to facilitate their interpretation and application. The framing of these distinct competencies was explicitly intended to align with the highest level of Miller's four-level Pyramid of Competency Assessment (i.e., *does*). Miller's Pyramid is a framework for teaching and assessing clinical reasoning or competencies amongst health profession learners^{27,28}. It provides a four-level, hierarchical depiction of competencies and their assessment, moving from the lowest level of *knows* (acquisition of knowledge), to *knows how* (application of knowledge), to *shows how* (demonstration of competency) and, finally, to the highest level of *does* (performing the competency within a clinical setting)²⁸. The *does* level requires adequate mastery of the three-lower levels (i.e., *knows*, *knows how* and *shows*), making it an attractive level of competency for this profile²⁷. While all levels of the pyramid are important, we assumed that the lower levels can be subsequently detailed within more granular educational profiles or curriculum guidelines to facilitate curriculum development, once there is consensus on these higher-level competencies.

Creation of a steering group

The steering group supervised the entire Delphi process. This group was formed by integrating members of different stakeholder groups, which included two people living with pain (LC, LS), one recent PT graduate (NM) and six university-based pain educators from Canadian institutions (AH, DW, GB, JM, TW, YTL). In addition, two methodological experts (AB, AT) provided input on the Delphi and implementation processes of this study. Individuals were invited to join the steering group based on their relevant knowledge, expertise, and past engagement. All

members of the steering group had equal involvement in the planning and decision-making related to all aspects of this project.

Delphi participants

Throughout this article, we used the term Delphi participants to refer to people who voted in the Delphi, which includes both university-based pain educators and clinical pain educators. Aligned with existing methodological recommendations, Delphi participants were recruited purposively²⁹.

Each of the 15 university PT programs in Canada has designated one or two representatives who lead the entry-level pain education at their local program; this national network of university-based pain educators is named the Pain Education in Physiotherapy Curriculum Initiative (PEPCI). Building on previous work¹⁸, we invited PEPCI members to participate in the study via email. PEPCI members from the same university were asked to consult one another in order to submit one vote per institution.

Clinical educators were also recruited to acknowledge the role that clinical placements play in shaping pain management skills for PT students. A recruitment email was circulated to members of the Canadian Physiotherapy Association (CPA) Pain Science Division, recent graduates from McGill University (Montreal, QC), University of Alberta (Edmonton, AB) and Université de Sherbrooke (Sherbrooke, QC) certificate in chronic pain management programs, as well as clinical specialists in pain from the CPA. Clinical educators were eligible if they had a minimum of five years of clinical practice experience focused on pain management as well as completing one of the following: (1) a post-licensure certificate program in pain management offered by a Canadian university or equivalent continuing education training in pain management; and/or (2)

certified as a Pain Science Clinical Specialist by the Physiotherapy Specialty Certification Board of Canada. In addition, clinical educators had to be actively involved in clinical supervision of PT students.

To ensure diversity in the recruitment of participants, the steering group also agreed on the following minimum diversity inclusion criteria for composition of the voting group: a) at least one participant that identified as a woman and at least one participant that identified as a man; b) at least one native English speaker and one native French speaker (to represent Canada's bilingual population); c) at least one clinician from public practice and one clinician from private practice; and d) at least one person who self-identifies as a Person of Color or as a member of Indigenous or First Nations community.

Modified Delphi Procedures

Anchored in the methodological recommendations for modified Delphi studies^{23,25}, the following steps were completed: 1) generation of the preliminary competency profile; 2) external review of the preliminary competency profile; 3) stakeholder survey to achieve consensus; 4) survey analysis and response to participants; and 5) evaluation of process. An overview of the entire process is presented in Figure 1.

Step 1: Generation of the preliminary competency profile

The steering group developed the preliminary version of the competency profile, based on themes from a recent national workshop which used a nominal group technique to ask key Canadian stakeholders in pain management what PTs should be able to do upon completing entry-level education. These stakeholders included pain educators, people living with pain and recent PT graduates. In addition, a literature search and seminal work in the field were used to

generate this preliminary version^{16,30,31}. Throughout this initial process, the steering group ensured that all competencies were aligned with the scope of the profile, while maintaining clarity and limiting redundancy.

Step 2: External review of the preliminary competency profile

A group of international leaders in research on pain education (n=5) reviewed the preliminary version of the competency profile to ensure that the content and scope of the items were consistent with existing international recommendations and the broader literature on competencies^{16,30,31}. This expert consulting group included members of: the task force that developed the International Association for the Study of Pain (IASP) Inter-professional Core Competencies in Pain Management, the task force that developed the IASP PT Pain Education Curriculum Guidelines and a PT faculty that systematically implemented the IASP PT Pain Education Curriculum Guidelines within their entry-level education program.

Step 3: Stakeholder survey to achieve consensus

Interested Delphi participants were invited to attend an online orientation meeting detailing the processes involved in the Delphi, such as the voting scheme and the expected timeline. They were provided with background information regarding the rationale and the research supporting the development of the competency profile, prior to completing the Delphi survey. Any questions were answered at that time and participants were encouraged to seek clarification via email if needed. At the start of each round, participants were emailed a link to the online Delphi survey and provided consent before voting. Data were collected between July and September 2020 using SurveyMonkey (San Mateo, CA, USA).

Voting process

Delphi participants were asked to anonymously rate their level of agreement with each competency according to the following scale: a) favorable to its inclusion in the competency profile (green light); b) favorable to the inclusion of a revised version (yellow light); or c) favorable to removing it from the competency profile (red light). In addition, participants were asked to rate their level of agreement with the definition of key concepts related to the competencies (e.g., social justice or therapeutic alliance) using the same scale. They were also invited to provide a rationale for their response and suggest potential improvements for each item (i.e., competencies or definitions) using an open-text box. Aligned with Delphi methodology best practices, the threshold for consensus was defined a priori at $\geq 75\%$ agreement across all participants to retain or exclude an item^{29,32}. We expected this study to require two to four rounds to reach consensus on all competencies.

Step 4: Survey analysis and response to participants

Following each round, the level of participant agreement for each item was summarized using frequency counts. Participants' open-text responses for items not reaching consensus were summarized and used to inform the revision of the competency profile within each round. The steering group had access to the summary of feedback, as well as the anonymized comments, to enhance rigor and credibility. The steering group updated the competencies that did not reach consensus, based on the feedback provided. Participants were provided with an overview of the votes (i.e., the frequency of green light, yellow light and red light votes for each item), as well as a summary of the feedback received alongside the new version (revised iteration) of the items not reaching consensus. Modification of an item was clearly highlighted within the survey and both versions of the item were included to allow for comparison. Steps 3 and 4 were repeated

iteratively until all items reached consensus for inclusion or exclusion. Following which, a summary of the changes was circulated alongside the finalized competency profile.

Step 5: Evaluation of process

Participants were asked to complete an anonymous post-Delphi survey, to evaluate satisfaction with the overall process using a seven-point scale (ranging from “very unsatisfied” to “very satisfied”). In addition, the survey assessed participants’ interest in using a similar Delphi process when making future consensus decisions related to this topic. Feedback from participants regarding potential improvements was sought using open-text comments.

Results

Delphi participation

A total of 20 participants were recruited, including representatives from 14 Canadian PT universities (14/15, 93% of all Canadian entry-level PT programs) and six clinical educators; predefined diversity criteria were met (Table 1). One university representative was unable to participate in this study, being on leave during the period of data collection. Nonetheless, the number of participants was within the typical range for Delphi studies^{23,29}. All participants completed the two rounds for item evaluation and answered the evaluation survey to provide feedback on the process (response rate: 100%).

Competency profile

The Delphi process resulted in the generation and stakeholder endorsement of the Pain Education in Physiotherapy (PEP) competency profile, which consists of 15 competencies that delineate the necessary abilities required to manage pain upon entry-to-practice (Table 2). The steering group initially generated a preliminary version of the competency profile, which consisted of 15

competencies and 17 key concept definitions. Throughout the external review and the Delphi evaluation process, five new key concept definitions were added to the profile and existing items were revised to enhance clarity, ensure the use of adequate terminology (e.g., *developing a diagnosis*) and update item phrasing to better align it with the profile values (e.g., using “assessment *with* the person” rather than “assessment *of* the person” to further demonstrate person-centeredness). By the end of the second round of the Delphi process, all 15 competencies and 23 key concept definitions reached the required consensus threshold (75%). Most competencies (12/15, 80%) reached a high level of endorsement (85% and above). A breakdown of the final agreement rate for each competency and the number of rounds required for reaching consensus is provided in Table 3.

The competencies can be divided into two different domains (Table 2), which emerged iteratively during the generation of the preliminary profile. The first domain is composed of eight competencies that address specific aspects of pain management, such as performing a comprehensive assessment, using appropriate tools and strategies to monitor progress or collaborating with relevant professionals. The second domain consists of seven competencies that permeate all aspects of pain management, such as promoting autonomy, using an ethical approach to care or demonstrating active listening and empathy.

Post-Delphi evaluation

All 20 participants completed the post-Delphi questionnaire. The findings suggest a high level of satisfaction with the Delphi process, with nearly all respondents feeling very satisfied (n=17, 85%) or satisfied (n=2, 10%) with it. Only one respondent reported being partially satisfied with the process. All respondents (n=20) mentioned they would be interested in using a similar Delphi process when making future consensus decisions related to this project. Responses to the open-

ended questions were consistent with this high level of satisfaction and endorsement. Several participants stated that the Delphi process was clearly outlined and easy to follow. Potential improvements suggested by participants include discussing some of the modifications with other participants, through focus groups. The clarity of the results and the justification provided when adapting an item led participants to report feeling included throughout the process. Comments, regarding the final competency profile were encouraging, although it was highlighted that some competencies lacked the sufficient granularity to inform specific teaching and learning strategies, due to the high-level scope of the profile.

Discussion

This study provides a novel contribution to the literature on pain management education. The purpose of this study was to generate and gain stakeholder consensus on a pain management competency profile that is specific to the Canadian PT context. By doing so, this work aimed to address the identified need to develop core competencies in pain management that are specific to a region (i.e., Canada) and a profession (i.e., PT), and that could be used to inform entry-level curriculum design and assessment.

One of the novel aspects of the PEP competency profile lies in the use of robust and transparent consensus-building methods to guide the development of this competency profile. The concept of integrated knowledge translation (KT) research also informed some of the methodological choices, such as the inclusion of key stakeholders in the research process³³. Integrated KT research calls for involvement of end users in the creation of new knowledge and provides a strong foundation to support the resulting competency profile^{33,34}. More specifically, collaborating with people living with pain as active members of the steering group is likely to have contributed to the person-centeredness of the PEP competency profile by building on their

lived experience of pain and key aspects of care that often remain unaddressed^{9-11,35}. In addition, engaging clinical pain educators in the voting process helped ensure that the competencies were directly relevant to real-life settings in addition to their academic pertinence. The contribution of key stakeholders actively involved in the topic through their lived experience of pain and their practical experience in providing pain management may have led to the emergence of competencies dedicated to interpersonal skills (e.g., using adequate communication strategies or promoting autonomy) that related to all aspects of pain management. Including people living with pain in the steering group is an important and novel improvement on previous work. However, including a sample of people living with pain as part of the external review process would have permitted the integration of a broader range of perspectives (e.g., via national advocacy groups) and enabled members of this stakeholder group to serve as external evaluators. Future work related to consensus building processes in pain should consider this additional level of involvement of people living with pain.

An interesting characteristic of the PEP competency profile was the emergence of two competency domains during the generation of the preliminary profile (Table 2). The first domain delineates competencies that address specific aspects of pain management, such as performing a comprehensive pain assessment or facilitating appropriate transitions in care for the person living with pain. The second domain includes competencies relevant to more global aspects of pain management, such as using a person-centered approach and tailored communication strategies. Each competency in the second domain is intended to be relevant to the demonstration of competencies in the first domain (e.g., using a person-centered approach and tailored communication strategies while completing a comprehensive pain assessment). The second domain may be broad, but we believe it brings in a novel piece that our stakeholders felt was not

addressed in previous work. While the ideas captured by this domain are likely applicable to all approaches of care and may overlap with the existing Canadian physiotherapy competency profile³⁰, they are of particular importance in the context of pain management. Having a domain dedicated to competencies relevant to all aspects of pain management is valuable to facilitate the in-depth development of these skills, while limiting redundancy by not repeating each of the interpersonal aspects of care that are vital to each of the specific aspects of pain management. It is also useful to facilitate both teaching and evaluation strategies by emphasizing what common aspects of care can be taught and assessed in relation to specific aspects of pain management.

The final version of the PEP competency profile bears some resemblance with previous work in the field, such as the IASP curriculum guidelines³¹ and the European Pain Federation core curriculum³⁶. These guidelines describe curriculum content related to pain management but do not answer the identified need for high-level competencies that delineate the required abilities to apply such content successfully and provide adequate pain management upon entry-to-practice. Our work is also similar to the competency profile generated by Fishman and colleagues, which has a strong interprofessional focus¹⁶. Both competency profiles advocate for using a multi-dimensional and person-centered approach to pain management, while considering the various contextual factors likely to influence access to and provision of care. In addition, they emphasize the importance of using adequate tools and strategies in pain management, as well as including communication competencies such as expressing compassion and empathy. However, the PEP competency profile further extends the work from Fishman and colleagues¹⁶, as a discipline-specific resource, but also by expanding the breadth and depth of the competencies included. For example, Fishman's competency related to empathic and compassionate communication is anchored in the assessment of pain (domain 2, competency 4), whereas the PEP competency

profile frames these elements in relation to all aspects of care (competencies 11 and 15). Similarly, the notion of advocacy is anchored in the treatment process in Fishman's profile (domain 4, competency 5), whereas we included different contexts for advocacy, such as at the level of the family, care providers or institution (competency 8) without anchoring them in a specific aspect of care. While it could be argued that these differences are the results of having a different focus (i.e., interprofessional versus PT-specific), the unique structure of the PEP competency profile plays an important role in this increased depth. More specifically, it differentiates between competencies addressing specific aspects of pain management (first domain) and competencies relevant to all aspects of pain management (second domain). This allows competencies such as promoting autonomy to be explicitly linked to all specific aspects of pain management without excessive repetitions. To our knowledge, this is the first time that a competency profile in pain management includes a stand-alone domain focused on overarching competencies. This is particularly relevant as the lack of these competencies (e.g., adequate communication, demonstrating empathy or using a person-centered approach to care) has been linked with impaired therapeutic alliance and increased stigma for people living with pain^{9,11,35}. Having these competencies clearly identified in the PEP competency profile holds the potential to help address them better and improve their application to clinical practice.

Another strength of the PEP competency profile is its alignment with how the broader literature defines competencies^{26,27}. This is the result of the input from educational experts, as they helped to ensure that the preliminary version of the PEP competency profile was consistent with the existing literature on competencies. In addition, the scope of the competency profile was made explicit to Delphi participants from the beginning, as the orientation session provided them with the definition of a competency as “an observable ability of a health professional, integrating

multiple components such as knowledge, skills, and attitudes^{26(p641)} and clarified how the competencies would be anchored in this definition. Throughout the Delphi process, the steering group ensured that framing of the competencies was explicitly aligned with the highest level of Miller's four-level Pyramid of Competency Assessment (i.e., does). As a result, the final competencies are all located at the does level of Miller's pyramid, which describes the situation where the learner has acquired the desired skill and performs it appropriately in clinical practice²⁸. We expect that the PEP competency profile could be used to inform curriculum guidelines and detail the competencies located at lower levels. This choice allowed us to omit competencies from these lower levels of Miller's pyramid, as compared to the work from Fishman and colleagues¹⁶, which included content located on different levels. For example, the competencies of the first domain from Fishman and colleagues' profile detail what the learner should know rather than the practical application of the desired knowledge, skills and attitudes in clinical settings.

It should be acknowledged that due to our focus on the does level, some stakeholders have reported wanting more granularity to inform the practical applications of the PEP competency profile, such as detailing content (e.g., best practice recommendations), as well as teaching and learning strategies. This feedback is important and highlights the need for subsequent work around the practical implementation of the PEP competency profile. An important next step would therefore be to expand the PEP competency profile through developing curriculum guidelines, entrustable professional activities, and assessment strategies. We also acknowledge that important barriers to implementing these competencies are still possible, both at a local program level and at a national regulatory level, which might impede uptake of the PEP competency profile. Future work with relevant stakeholders (e.g., PT program directors,

university-based pain educators or national regulators) is needed to unpack the barriers and facilitators related to uptake of our competency profile. It is also important to point out that the protocol for this study was not registered prior to its implementation. While this is not standard practice for Delphi methodology, or for other consensus-building methods that are unrelated to hypothesis testing²²⁻²⁵, pre-registration would help further improve transparency.

Conclusion

A national Delphi process was used to achieve consensus on a novel competency profile specific to the Canadian PT context, that delineates the necessary competencies required to manage pain upon entry-to-practice. The steering group members and the review provided by the international leaders in research on pain education helped create a preliminary version of the profile, which was subsequently refined and endorsed by Delphi participants, which consisted of university-based (n=14) and clinical (n=6) pain educators. The final version of the PEP competency profile includes 15 competencies, which can be divided into two interdependent domains. The first domain focuses on specific aspects of pain management, whereas the second domain details competencies relevant to all aspects of pain management. Building on this profile, future research will need to detail the content relevant to each competency, develop curriculum guidelines for pain education in Canada and understand potential barriers to implementation. The consensus-building and participatory research methods used in this study provides a practical example of the processes involved in the creation and approval of a competency profile, and may be useful to inform future interventions to improve pain education across healthcare professions and geographic regions.

ACCEPTED MANUSCRIPT

References

1. Canadian Pain Task Force. Chronic Pain in Canada: Laying a Foundation for Action: Government of Canada; 2019 [Available from: <https://www.canada.ca/en/health-canada/corporate/about-health-canada/public-engagement/external-advisory-bodies/canadian-pain-task-force/report-2019.html>].
2. The National Institutes of Health. National Pain Strategy - A Comprehensive Population Health -Level Strategy for Pain. The National Institutes of Health; 2016.
3. National Academies of Sciences. The National Academies Collection: Reports funded by National Institutes of Health. In: Stroud C, Posey Norris SM, Bain L, editors. The Role of Nonpharmacological Approaches to Pain Management: Proceedings of a Workshop. Washington (DC): National Academies Press (US)
4. Watt-Watson J, McGillion M, Hunter J, et al. A survey of prelicensure pain curricula in health science faculties in Canadian universities. *Pain research & management*. 2009;14(6):439-44.
5. Thompson K, Johnson MI, Milligan J, Briggs M. Twenty-five years of pain education research-what have we learned? Findings from a comprehensive scoping review of research into pre-registration pain education for health professionals. *Pain*. 2018;159(11):2146-58.
6. Synnott A, O'Keeffe M, Bunzli S, Dankaerts W, O'Sullivan P, O'Sullivan K. Physiotherapists may stigmatise or feel unprepared to treat people with low back pain and psychosocial factors that influence recovery: a systematic review. *Journal of physiotherapy*. 2015;61(2):68-76.
7. Alexanders J, Anderson A, Henderson S. Musculoskeletal physiotherapists' use of psychological interventions: a systematic review of therapists' perceptions and practice. *Physiotherapy*. 2015;101(2):95-102.
8. Toye F, Seers K, Hannink E, Barker K. A mega-ethnography of eleven qualitative evidence

syntheses exploring the experience of living with chronic non-malignant pain. *BMC medical research methodology*. 2017;17(1):116.

9. De Ruddere L, Craig KD. Understanding stigma and chronic pain: a-state-of-the-art review. *Pain*. 2016;157(8):1607-10.

10. Wideman TH, Edwards RR, Walton DM, Martel MO, Hudon A, Seminowicz DA. The Multimodal Assessment Model of Pain: A Novel Framework for Further Integrating the Subjective Pain Experience Within Research and Practice. *Clin J Pain*. 2019;35(3):212-21.

11. Cohen M, Quintner J, Buchanan D, Nielsen M, Guy L. Stigmatization of Patients with Chronic Pain: The Extinction of Empathy. *Pain Medicine*. 2011;12(11):1637-43.

12. Birnie KA, Dib K, Ouellette C, et al. Partnering For Pain: a Priority Setting Partnership to identify patient-oriented research priorities for pediatric chronic pain in Canada. *CMAJ Open*. 2019;7(4):E654-e64.

13. Forsythe LP, Carman KL, Szydlowski V, et al. Patient Engagement In Research: Early Findings From The Patient-Centered Outcomes Research Institute. *Health Aff (Millwood)*. 2019;38(3):359-67.

14. Fitzcharles M-A, Brachaniec M, Cooper L, et al. A paradigm change to inform fibromyalgia research priorities by engaging patients and health care professionals. *Canadian Journal of Pain*. 2017;1(1):137-47.

15. Poulin P, Shergill Y, Romanow H, et al. Researching what matters to improve chronic pain care in Canada: A priority-setting partnership process to support patient-oriented research. *Canadian Journal of Pain*. 2018;2(1):191-204.

16. Fishman SM, Young HM, Arwood LE, et al. Core competencies for pain management: results of an interprofessional consensus summit. *Pain medicine (Malden, Mass)*.

2013;14(7):971-81.

17. van Lankveld W, Afram B, Staal JB, van der Sande R. The IASP pain curriculum for undergraduate allied health professionals: educators defining competence level using Dublin descriptors. *BMC Medical Education*. 2020;20(1):60.

18. Wideman TH, Miller J, Bostick G, Thomas A, Bussièrès A. Advancing Pain Education in Canadian Physiotherapy Programmes: Results of a Consensus-Generating Workshop. *Physiotherapy Canada Physiothérapie Canada*. 2018;70(1):24-33.

19. Wideman TH, Miller J, Bostick G, Thomas A, Bussières A, Wickens RH. The current state of pain education within Canadian physiotherapy programs: a national survey of pain educators. *Disabil Rehabil*. 2019:1-7.

20. Moseley GL, Butler DS. Fifteen Years of Explaining Pain: The Past, Present, and Future. *The journal of pain : official journal of the American Pain Society*. 2015;16(9):807-13.

21. Thacker L, Walsh RM, Song SG, et al. Exploring physiotherapy practice within hospital-based interprofessional chronic pain clinics in Ontario. *Canadian Journal of Pain*. 2021;5(1):96-106.

22. Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs*. 2000;32(4):1008-15.

23. Boukchedid R, Abdoul H, Loustau M, Sibony O, Alberti C. Using and reporting the Delphi method for selecting healthcare quality indicators: a systematic review. *PLoS One*. 2011;6(6):e20476.

24. Hauer KE, Cate OT, Boscardin CK, et al. Ensuring Resident Competence: A Narrative Review of the Literature on Group Decision Making to Inform the Work of Clinical Competency Committees. *J Grad Med Educ*. 2016;8(2):156-64.

25. Humphrey-Murto S, Varpio L, Wood TJ, et al. The Use of the Delphi and Other Consensus Group Methods in Medical Education Research: A Review. *Acad Med.* 2017;92(10):1491-8.
26. Frank JR, Snell LS, Cate OT, et al. Competency-based medical education: theory to practice. *Medical Teacher.* 2010;32(8):638-45.
27. Ten Cate O, Carraccio C, Damodaran A, et al. Entrustment Decision Making: Extending Miller's Pyramid. *Acad Med.* 2021;96(2):199-204.
28. Miller G. The assessment of clinical skills/competence/performance. *Academic Medicine.* 1990;65(9):A63-S7.
29. Taylor E. We Agree, Don't We? The Delphi Method for Health Environments Research. *Herd.* 2020;13(1):11-23.
30. NPAG. National Physiotherapy Advisory Group. Competency Profile for Physiotherapists in Canada <https://www.peac-aepec.ca/pdfs/Resources/Competency%20Profiles/Competency%20Profile%20for%20PTs%202017%20EN.pdf>: Physiotherapy Education Accreditation Canada; 2017 [
31. International Association for the Study of Pain. Curriculum outline on pain for physical therapy 2018 [cited 2018 Mar 29]. Available from: <https://www.iasp-pain.org/Education/CurriculaList.aspx>.
32. Diamond IR, Grant RC, Feldman BM, et al. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol.* 2014;67(4):401-9.
33. Jull J, Giles A, Graham ID. Community-based participatory research and integrated knowledge translation: advancing the co-creation of knowledge. *Implementation Science.* 2017;12(1):150.

34. Gagliardi AR, Berta W, Kothari A, Boyko J, Urquhart R. Integrated knowledge translation (IKT) in health care: a scoping review. *Implementation Science*. 2016;11(1):38.
35. Slade SC, Molloy E, Keating JL. Stigma experienced by people with nonspecific chronic low back pain: a qualitative study. *Pain medicine (Malden, Mass)*. 2009;10(1):143-54.
36. European Pain Federation. Core Curriculum for European Diploma in Pain Physiotherapy 2017 [Available from: <https://europeanpainfederation.org/wp-content/uploads/2018/10/EFIC-Pain-Physiotherapy-Curriculum1.pdf>].
37. Knights S. Reflection and learning: the importance of a listener. In: Boud D, Keogh R, Walker D, editors. *Reflection: turning experience into learning*. New York: Nichols Publishing Company; 1985.
38. Beauchamp T. Principlism in bioethics. In: Bermúdez P, Seoane J, editors. *Bioethical Decision Making and Argumentation*. 1st ed: Springer International Publishing; 2016.
39. Foster NE, Hill JC, O'Sullivan P, Hancock M. Stratified models of care. *Best Pract Res Clin Rheumatol*. 2013;27(5):649-61.
40. Setchell J, Dalziel B. Using Critical Reflexivity to Enhance Clinical Care: A Clinician Perspective. *Journal of Humanities in Rehabilitation*. 2019.
41. First Nation Health Authority. Cultural Humility 2020 [Available from: <https://www.fnha.ca/wellness/cultural-humility>].
42. Mercer SW, Reynolds WJ. Empathy and quality of care. *Br J Gen Pract*. 2002;52 Suppl(Suppl):S9-12.
43. World Health Organization. Health equity: World Health Organization; 2020 [Available from: https://www.who.int/topics/health_equity/en/].
44. Canadian Physiotherapy Association. Code of ethical conduct 2020 [Available from:

<https://physiotherapy.ca/sites/default/files/code-of-conduct-en.pdf>.

45. Maclachlan M, Khasnabis C, Mannan H. Inclusive health. *Trop Med Int Health*. 2012;17(1):139-41.

46. Oxford Reference. Lived Experience 2020 [Available from:

<https://www.oxfordreference.com/view/10.1093/oi/authority.20110803100109997>.

47. Okun S, Schoenbaum S, Andrews D, et al. Patients and Health Care Teams Forging Effective Partnerships. *NAM Perspectives*. 2014.

48. Dukhanin V, Searle A, Zwerling A, Dowdy DW, Taylor HA, Merritt MW. Integrating social justice concerns into economic evaluation for healthcare and public health: A systematic review. *Soc Sci Med*. 2018;198:27-35.

49. Kinney M, Seider J, Beaty AF, Coughlin K, Dyal M, Clewley D. The impact of therapeutic alliance in physical therapy for chronic musculoskeletal pain: A systematic review of the literature. *Physiother Theory Pract*. 2020;36(8):886-98.

50. Barlow J, Wright C, Sheasby J, Turner A, Hainsworth J. Self-management approaches for people with chronic conditions: a review. *Patient Education and Counseling*. 2002;48(2):177-187.

Figure legends.

Figure 1: Overview of modified Delphi process.

This figure provides an overview of the different steps involved in our modified Delphi process.




Table 1: Summary of recruited clinical educators.

Characteristics	Participants (percentage of sample)
Practice settings	
Private practice	3 (50%)
Public practice	3 (50%)
Expertise in pain management	
Graduate certificate	3 (50%)
Clinical specialist	1 (16.67%)
Other training	2 (33.33%)
Gender	
Woman	5 (83.33%)
Man	1 (16.67%)
Mother tongue	
English	4 (66.66%)
French	1 (16.67%)
Other	1 (16.67%)
Self-identifies as indigenous person, racialized person or visible minority?	
Yes	1 (16.67%)
No	5 (83.33%)
This table provides a summary of the pre-determined diversity characteristics used to recruit clinical educators.	

Table 2: Final version of the PEP competency profile (bolded words are defined in the glossary).

Domain 1: Competencies addressing specific aspects of pain management	
Competency 1	Facilitate the development of a therapeutic alliance with the person living with pain.
Competency 2	Perform a comprehensive assessment with the person living with pain that uses appropriate tools and strategies to explore and evaluate the lived experience of pain, as well as the mechanisms underlying pain and the physical, psychological and socio-environmental factors that influence pain.
Competency 3	Synthesize and interpret assessment findings to develop a pain-related diagnosis and/or classification , and to generate a prognosis .
Competency 4	Develop and implement an individualized treatment plan that is based on the assessment findings and goals of the person living with pain.
Competency 5	Facilitate appropriate transitions in care for the person living with pain.
Competency 6	Use appropriate tools and strategies to monitor and evaluate the effectiveness of the treatment plan for the person living with pain, and adapt care accordingly.
Competency 7	Collaborate with relevant professionals in a manner that fosters an integrated, patient-centered approach to pain management.
Competency 8	Advocate with, and on behalf of, people living with pain, at the level of the individual, family and/or care providers, institution and community.
Domain 2: Competencies that permeate all aspects of pain management	
Competency 9	Use a person-centered approach to pain management that addresses the complex, multidimensional and subjective nature of pain.
Competency 10	Support and promote the autonomy of the person living with pain and foster partnership in their care.
Competency 11	Communicate with people living with pain in a way that is tailored to their individual needs and abilities, demonstrates active listening and empathy , and validates their lived experience.
Competency 12	Make practice decisions that are informed by principles of social justice , inclusiveness and equity , and promote cultural safety .
Competency 13	Engage in critical self-reflection that fosters continuous professional growth and development.
Competency 14	Integrate best research evidence, clinical expertise and patient values when making practice decisions.
Competency 15	Use a safe, ethical and compassionate approach to care.
Glossary	
Active listening	Active listening involves “placing all of one’s attention and awareness at the disposal of another person, listening with interest and appreciating without interrupting”. ³⁷
Appropriate	Appropriate tools and strategies are those that are best aligned with research

tools and strategies	evidence, clinical expertise and patient values.
Autonomy	Autonomy refers to a person's capacity to self-govern and make and enact choices that are aligned with one's values and life goals. ³⁸
Classification	Classification refers to identifying the underlying set of mechanisms and/or prognostic factors that are suspected to contribute to the pain experience and/or recovery. ³⁹
Communicate	Communicate is understood to include oral, non-verbal and written forms of communication. ³⁰
Compassionate	A compassionate approach to care involves recognizing, acknowledging and working to alleviate suffering.
Comprehensive assessment	Comprehensive assessment refers to listening to the pain narrative and conducting a subjective interview and physical examination. This assessment aims to obtain information about pain-related impairments, activity limitations, participation restrictions as well as personal and environmental factors influencing pain, health and function. Comprehensive assessment also needs to accurately screen, identify and triage conditions that require immediate action, that would benefit from non-urgent referral to other health services, or that would benefit from modifications to the pain management plan.
Critical self-reflection	Critical self-reflection can be used interchangeably with the term "critical reflexivity" and involves examining the assumptions, beliefs, and values that underpin one's thinking and practice. ⁴⁰
Cultural safety	Cultural safety refers to creating "an environment free of racism and discrimination, where people feel safe when receiving health care". ⁴¹
Diagnosis	Diagnosis refers to a physiotherapy diagnosis, which is defined as "a conclusion about physical function based on a subjective and objective assessment and analysis by a physiotherapist to investigate the cause or nature of a client's condition or problem". ³⁰ Developing a diagnosis may not always be possible and is not required for effective pain management.
Empathy	Empathy refers to the recognition and understanding of another person's experience. ⁴²
Equity	Equity is "the absence of avoidable, unfair, or remediable differences among groups of people". ⁴³
Ethical	An ethical approach to care is characterized by actions that align with the core principles of the Canadian Physiotherapy Association's Code of Ethical conduct, namely: respect for autonomy, beneficence, least harm and justice. ⁴⁴
Inclusiveness	Inclusiveness implies "health services that work for all and are equitable". ⁴⁵
Lived experience	Lived experience refers to "personal knowledge about the world gained through direct, first-hand involvement in everyday events". ⁴⁶
Partnership	Partnership refers to health care providers working in concert with patients and their caregivers to achieve positive experiences and mutually agreed-upon outcomes. ⁴⁷
Person-centred approach	Person-centered approach is intended to emphasize the importance of placing the needs of the person living with pain at the centre of pain management.
Person living	A person living with pain refers to any person living with any type of pain, for

with pain	any period of time.
Prognosis	Prognosis refers to the probable course of a person's health and functional status. Prognosis can evolve over the course of treatment.
Social justice	Social justice is the "moral imperative to avoid and remediate unfair distributions of societal disadvantage". ⁴⁸
Therapeutic alliance	Therapeutic alliance is defined as "a collaborative relationship characterized by a reciprocal trust between the client and the clinician, as well as the development of mutually agreed-upon goals and interventions". ⁴⁹
Transitions in care	Transitions in care refers to the transfer of a patient between different settings, health professionals, or health services. Transitions in care include discharge and linking people living with pain to available resources.
Treatment plan	Consistent with a physiotherapy scope of practice, the treatment plan should aim to provide self-management support, ⁵⁰ facilitate engagement in physical activity or other meaningful activities, and support the effective management of pain-related symptoms.

ACCEPTED MANUSCRIPT

Table 3: Summary of final agreement rate (green light vote) for each competency.

Consensus	95%	90%	85%	80%	75%
First round	Competency 1	Competency 5 Competency 12	Competency 2 Competency 6 Competency 9 Competency 13		Competency 8 Competency 11
Second round (for revised items)	Competency 15	Competency 7	Competency 3 Competency 4 Competency 10	Competency 14	

This table summarizes the final agreement rate for each competency. Competencies that did not meet the pre-determined consensus threshold (75%) for inclusion or exclusion were modified by the steering group and resubmitted to the voting participants for a second round.